

A NEW MODULAR PLETHYSTIC $\mathrm{SL}_2(\mathbb{F})$ -ISOMORPHISM

$$\mathrm{Sym}^{N-1} E \otimes \bigwedge^{N+1} \mathrm{Sym}^{d+1} E \cong \Delta^{(2,1^{N-1})} \mathrm{Sym}^d E$$

ALVARO L. MARTINEZ AND MARK WILDON

ABSTRACT. Let \mathbb{F} be a field and let E be the natural representation of $\mathrm{SL}_2(\mathbb{F})$. Given a vector space V , let $\Delta^{(2,1^{N-1})} V$ be the kernel of the multiplication map $\bigwedge^N V \otimes V \rightarrow \bigwedge^{N+1} V$. We construct an explicit $\mathrm{SL}_2(\mathbb{F})$ -isomorphism $\mathrm{Sym}^{N-1} E \otimes \bigwedge^{N+1} \mathrm{Sym}^{d+1} E \cong \Delta^{(2,1^{N-1})} \mathrm{Sym}^d E$. This $\mathrm{SL}_2(\mathbb{F})$ -isomorphism is a modular lift of the q -binomial identity $q^{\frac{N(N-1)}{2}} [N]_q \begin{bmatrix} d+2 \\ N+1 \end{bmatrix}_q = s_{(2,1^{N-1})}(1, q, \dots, q^d)$, where $s_{(2,1^{N-1})}$ is the Schur function for the partition $(2, 1^{N-1})$. This identity, which follows from our main theorem, implies the existence of an isomorphism when \mathbb{F} is the field of complex numbers but it is notable, and not typical of the general case, that there is an explicit isomorphism defined in a uniform way for any field.

1. INTRODUCTION

Let \mathbb{F} be an arbitrary field and let E be the natural 2-dimensional representation of the special linear group $\mathrm{SL}_2(\mathbb{F})$. Let Δ^λ denote the Schur functor canonically labelled by the partition λ . Working over the field of complex numbers there is a rich theory of plethystic isomorphisms between the representations $\Delta^\lambda \mathrm{Sym}^d E$. These include Hermite reciprocity and the Wronskian isomorphism; we refer the reader to [PW21] for a comprehensive account and references to earlier results. In [McDW22] it was shown that both these classical isomorphisms hold over an arbitrary field, *provided that suitable dualities are introduced*. The modular version of Hermite reciprocity is $\mathrm{Sym}_M \mathrm{Sym}^d E \cong \mathrm{Sym}^d \mathrm{Sym}_M E$, where, given a $\mathrm{SL}_2(\mathbb{F})$ -representation V , $\mathrm{Sym}^r V$ is the symmetric power defined as a quotient of $V^{\otimes r}$ and $\mathrm{Sym}_r V$ is its dual defined as the subspace of invariant tensors in $V^{\otimes r}$. The modular Wronskian isomorphism is $\mathrm{Sym}_M \mathrm{Sym}^d E \cong \bigwedge^M \mathrm{Sym}^{d+M-1} E$. The purpose of this article is to add to the collection of such modular plethystic isomorphisms by proving the following theorem. The version of the Schur functor $\Delta^{(2,1^{N-1})}$ we require is defined in §1.1 immediately below.

Theorem 1.1. *Let $N \in \mathbb{N}$ and let $d \in \mathbb{N}_0$. The map φ defined in Definition 1.8 is an isomorphism of $\mathrm{SL}_2(\mathbb{F})$ -representations*

$$\mathrm{Sym}^{N-1} E \otimes \bigwedge^{N+1} \mathrm{Sym}^{d+1} E \cong \Delta^{(2,1^{N-1})} \mathrm{Sym}^d E.$$

1991 *Mathematics Subject Classification*. Primary: 20C20, Secondary: 05E05, 05E10, 17B10, 20G05.

This theorem is notable as the first explicit example of a modular plethystic isomorphism involving a Schur functor for a partition that is not one-row or one-column, and also for the unexpected tensor factorisation it exhibits.

This isomorphism is a modular lift of the q -binomial identity

$$q^{\frac{N(N-1)}{2}} [N]_q \begin{bmatrix} d+2 \\ N+1 \end{bmatrix}_q = s_{(2,1^{N-1})}(1, q, \dots, q^d) \quad (1.1)$$

where $s_{(2,1^{N-1})}$ denotes the Schur function labelled by the partition $(2, 1^{N-1})$. We structure our proof so that we can obtain (1.1) as a fairly routine corollary of Theorem 1.1: see Corollary 3.2, where we also give combinatorial interpretations of each side. As shown in [McDW22, Theorem 1.6] there exist representations of the form $\Delta^\lambda \text{Sym}^d E$ that have equal q -characters in the sense of (1.1), and so are isomorphic over \mathbb{C} , but fail to be isomorphic over arbitrary fields \mathbb{F} , even after considering all possible dualities. Indeed, the authors believe this is the generic case. This adds to the interest and importance of Theorem 1.1. We finish with Corollary 3.1, which lifts the isomorphism in Theorem 1.1 to an isomorphism of representations of $\text{GL}_2(\mathbb{F})$, and Conjecture 3.3 on a conjectured more general isomorphism.

1.1. Preliminaries. Fix a basis X, Y of the \mathbb{F} -vector space E . For each $r \in \mathbb{N}_0$, the symmetric power $\text{Sym}^c E$ has as a basis the monomials $X^{c-i} Y^i$ for $0 \leq i \leq c$.

Schur functor. It will be convenient to define the Schur functor $\Delta^{(2,1^{N-1})}$ on a vector space V by

$$\Delta^{(2,1^{N-1})} V = \ker \mu_N : \bigwedge^N V \otimes V \rightarrow \bigwedge^{N+1} V \quad (1.2)$$

where μ_N is the multiplication map $v_1 \wedge \dots \wedge v_N \otimes w \mapsto v_1 \wedge \dots \wedge v_N \wedge w$. Thus $\Delta^{(2,1^{N-1})} V$ is a subspace of $\bigwedge^N V \otimes V$. Since μ_N is a homomorphism of representations of $\text{GL}(V)$, for any fixed group G , $\Delta^{(2,1^{N-1})}$ is a functor on the category of \mathbb{F} -representations of G . This construction agrees with that in [EGS08, Ch. 5]; the dual module is constructed in [dBPW21, §2.3] and in [EGS08, Ch. 4].

Multi-indices. For $c, r \in \mathbb{N}_0$, let $I^{(c)}(r)$ denote the set $\{0, 1, \dots, c\}^r$. We say that the elements of $I^{(c)}(r)$ are *multi-indices*. We define the *sum* of a multi-index \mathbf{i} by $|\mathbf{i}| = \sum_{\alpha=1}^r i_\alpha$. Given $\mathbf{i} \in I^{(c)}(r)$ we define

$$F_\wedge^{(c)}(\mathbf{i}) = X^{c-i_1} Y^{i_1} \wedge \dots \wedge X^{c-i_r} Y^{i_r}.$$

Thus $\bigwedge^r \text{Sym}^c E$ has as a basis all $F_\wedge^{(c)}(\mathbf{i})$ for strictly increasing $\mathbf{i} \in I^{(c)}(r)$. We say that a pair $(\mathbf{i}, j) \in I^{(c)}(r) \times \{0, 1, \dots, c\}$ is *semistandard* if \mathbf{i} is strictly increasing and $i_1 \leq j$. Observe that (\mathbf{i}, j) is semistandard if and only if the $(2, 1^{N-1})$ -tableau $t_{(\mathbf{i}, j)}$ shown in the margin having entry i_α in box $(\alpha, 1)$ and entry j in box $(1, 2)$ is semistandard in the usual sense.

$$t_{(\mathbf{i}, j)} = \begin{array}{|c|c|} \hline i_1 & j \\ \hline i_2 & \\ \hline \vdots & \\ \hline i_N & \\ \hline \end{array}$$

Definition 1.2 (Content and Neighbour). Let $d \in \mathbb{N}_0$. Let $\mathbf{i} \in I^{(d)}(N)$ and $j \in \{0, 1, \dots, d\}$ and let (\mathbf{i}, j) be semistandard. We define the *content* of (\mathbf{i}, j) to be the multiset $\{i_1, \dots, i_N\} \cup \{j\}$. We define the *neighbour* of (\mathbf{i}, j) by

$$\mathcal{P}(\mathbf{i}, j) = ((i_1, \dots, i_{\alpha-1}, j, i_{\alpha+1}, \dots, i_N), i_\alpha) \quad (1.3)$$

where $\alpha \in \{1, \dots, N\}$ is maximal such that $i_\alpha \leq j$.

Observe that the neighbour map is well-defined because $i_1 \leq j$ and that it preserves content. Moreover, $\mathcal{P}(\mathbf{i}, j) = (\mathbf{i}, j)$ if and only if j is in the set $\{i_1, \dots, i_N\}$; in this case j is the unique repeated element of the multiset. For example, repeated applications of the neighbour map give

$$((0, 2, 3), 5) \xrightarrow{\mathcal{P}} ((0, 2, 5), 3) \xrightarrow{\mathcal{P}} ((0, 3, 5), 2) \xrightarrow{\mathcal{P}} ((2, 3, 5), 0) \quad (1.4)$$

in which all pairs have content $\{0, 2, 3, 5\}$ and the final pair is the only one that is not semistandard and so does not have a defined neighbour.

Definition 1.3. Let $d \in \mathbb{N}_0$. Let $\mathbf{i} \in I^{(d)}(N)$ and $j \in \{0, 1, \dots, d\}$. We define $F(\mathbf{i}, j) \in \bigwedge^N \text{Sym}^d E \otimes \text{Sym}^d E$ by

$$F(\mathbf{i}, j) = F_{\wedge}^{(d)}(\mathbf{i}) \otimes X^{d-j} Y^j. \quad (1.5)$$

If (\mathbf{i}, j) is semistandard we define

$$F_{\Delta}(\mathbf{i}, j) = \begin{cases} F(\mathbf{i}, j) & \text{if } j \in \{i_1, \dots, i_N\} \\ F(\mathbf{i}, j) + F(\mathcal{P}(\mathbf{i}, j)) & \text{otherwise.} \end{cases} \quad (1.6)$$

Whenever we use the notation $F(\mathbf{i}, j)$ in (1.5), the value of d will be clear from context. To give an example we take $N = 3$ and $d = 5$. Then, omitting some parentheses for readability, we have $F((0, 2, 5), 3) = X^5 \wedge X^3 Y^2 \wedge Y^5 \otimes X^2 Y^3$ and

$$\begin{aligned} F_{\Delta}((0, 2, 5), 3) &= F((0, 2, 5), 3) + F((0, 3, 5), 2) \\ &= X^5 \wedge X^3 Y^2 \wedge Y^5 \otimes X^2 Y^3 + X^5 \wedge X^2 Y^3 \wedge Y^5 \otimes X^3 Y^2. \end{aligned}$$

Semistandard basis. It is clear that $\bigwedge^N \text{Sym}^d E \otimes \text{Sym}^d E$ has as a canonical basis all $F(\mathbf{i}, j)$ for strictly increasing $\mathbf{i} \in I^{(d)}(N)$ and $j \in \{0, 1, \dots, d\}$. In this subsection we show that the $F_{\Delta}(\mathbf{i}, j)$ for semistandard (\mathbf{i}, j) form a basis for its submodule $\Delta^{(2, 1^{N-1})} \text{Sym}^d E$, which we defined to be the kernel of $\mu_N : \bigwedge^N V \otimes V \rightarrow \bigwedge^{N+1} V$ in (1.2).

We first show that these elements are in the kernel. Let $\mathbf{i} \in I^{(d)}(N)$ be strictly increasing and let $j \in \{0, 1, \dots, d\}$. Let α be maximal such that $i_\alpha \leq j$. Then

$$\begin{aligned} \mu_N F(\mathcal{P}(\mathbf{i}, j)) &= \mu_N F((i_1, \dots, i_{\alpha-1}, j, i_{\alpha+1}, \dots, i_N), i_\alpha) \\ &= X^{d-i_1} Y^{i_1} \wedge \dots \wedge X^{d-j} Y^j \wedge \dots \wedge X^{d-i_N} Y^{i_N} \wedge X^{d-i_\alpha} Y^{i_\alpha}. \end{aligned}$$

Up to a swap of the factor $X^{d-j}Y^j$ in position α and the final factor $X^{d-i_\alpha}Y^{i_\alpha}$, the right-hand side agrees with $\mu_N(F(\mathbf{i}, j))$. Hence, by the definition of F_Δ in (1.6), we have $\mu_N F_\Delta(\mathbf{i}, j) = 0$. In order to prove the basis theorem, we shall use the following lemmas.

Lemma 1.4. *Let $\{a_1, \dots, a_N, b\}$ be a subset of $\{0, 1, \dots, d\}$ with $a_1 < \dots < a_N < b$. The N distinct semistandard pairs with content $\{a_1, \dots, a_N, b\}$ are $\mathcal{P}^\alpha((a_1, \dots, a_N), b)$ for $\alpha \in \{0, 1, \dots, N-1\}$; the second elements of these pairs form the decreasing chain $b > a_N > \dots > a_2$. Furthermore, $\mathcal{P}^N((a_1, \dots, a_N), b)$ is not semistandard.*

Proof. We leave this to the reader as a routine generalisation of (1.4). \square

Lemma 1.5. *The $F_\Delta(\mathbf{i}, j)$ for semistandard $(\mathbf{i}, j) \in I^{(d)}(N) \times \{0, 1, \dots, d\}$ are linearly independent.*

Proof. It is clear from the canonical basis of $\bigwedge^N \text{Sym}^d E \otimes \text{Sym}^d E$ that if there is a non-trivial linear relation between the $F_\Delta(\mathbf{i}, j)$ then it may be assumed to involve only (\mathbf{i}, j) of the same content (in the sense of Definition 1.2). If $j \in \{i_1, \dots, i_N\}$ then there is a unique element $F_\Delta(\mathbf{i}, j)$ of content $\{i_1, \dots, i_N\} \cup \{j\}$. In the remaining case, the content multiset is a set, A say, and Lemma 1.4 applies. Since each $F_\Delta(\mathbf{i}, j)$ of content A is a sum of $F(\mathbf{i}, j)$ and its neighbour $F(\mathcal{P}(\mathbf{i}, j))$, they are linearly independent. \square

Lemma 1.6. *There are $N \binom{d+2}{N+1}$ semistandard Young tableaux of shape $(2, 1^{N-1})$ and entries in $\{0, 1, \dots, d\}$.*

Proof. Let S_α be the set of all Young tableaux $t_{(\mathbf{i}, j)}$ such that $i_\alpha \leq j < i_{\alpha+1}$ if $\alpha < N$, and such that $i_N \leq j$ if $\alpha = N$. It is clear that the set of semistandard Young tableaux of shape $(2, 1^{N-1})$ is partitioned into the N disjoint subsets S_1, \dots, S_N . We claim that each S_α has the same cardinality $\binom{d+2}{N+1}$. To see this, we define a bijection from S_α to the set of strictly increasing multi-indices in $I^{(d+1)}(N+1)$ by

$$(\mathbf{i}, j) \mapsto (i_1, \dots, i_\alpha, j+1, i_{\alpha+1}+1, i_{\alpha+2}+1, \dots, i_N+1).$$

The inverse of this map is easily shown to be

$$(k_1, \dots, k_{N+1}) \mapsto ((k_1, \dots, k_\alpha, k_{\alpha+2}-1, \dots, k_{N+1}-1), k_{\alpha+1}-1). \quad \square$$

Proposition 1.7. *The vector space $\Delta^{(2, 1^{N-1})} \text{Sym}^d E$ has dimension $N \binom{d+2}{N+1}$ and has as a basis*

$$\{F_\Delta(\mathbf{i}, j) : \mathbf{i} \in I^{(d)}(N), j \in \{0, 1, \dots, d\}, (\mathbf{i}, j) \text{ semistandard}\}.$$

Proof. By Lemma 1.5, the claimed basis is linearly independent. We use a dimension counting argument to show that it spans $\Delta^{(2, 1^{N-1})} \text{Sym}^d E$. By

Lemma 1.6, it suffices to show that $\dim \Delta^{(2,1^{N-1})} \text{Sym}^d E = N \binom{d+2}{N+1}$. This follows from the rank-nullity formula applied to (1.2):

$$\begin{aligned} \dim \ker \mu_N &= \binom{d+1}{N} (d+1) - \binom{d+1}{N+1} \\ &= \binom{d+1}{N} (d+2) - \left(\binom{d+1}{N} + \binom{d+1}{N+1} \right) \\ &= \binom{d+2}{N+1} (N+1) - \binom{d+2}{N+1} \\ &= N \binom{d+2}{N+1}. \end{aligned} \quad \square$$

1.2. Definition of φ . Given $0 \leq j < k$, set $[j, k] = \{j, j+1, \dots, k-1\}$. Given a strictly increasing multi-index $\mathbf{k} \in I^{(d+1)}(N+1)$, we define

$$\mathcal{B}(\mathbf{k}) = [k_1, k_2] \times [k_2, k_3] \times \dots \times [k_N, k_{N+1}] \subseteq I^{(d)}(N). \quad (1.7)$$

For example if $d = 5$ and $N = 3$, we have

$$\mathcal{B}((0, 2, 3, 6)) = [0, 2] \times [2, 3] \times [3, 6] = \{0, 1\} \times \{2\} \times \{3, 4, 5\} \subseteq I^{(5)}(3).$$

Definition 1.8. Fix $d \in \mathbb{N}_0$ and $N \in \mathbb{N}$. We define

$$\varphi : \text{Sym}^{N-1} E \otimes \bigwedge^{N+1} \text{Sym}^{d+1} E \rightarrow \bigwedge^N \text{Sym}^d E \otimes \text{Sym}^d E$$

by

$$\varphi(X^{N-1-s} Y^s \otimes F_{\wedge}^{(d+1)}(\mathbf{k})) = \sum_{\mathbf{i} \in \mathcal{B}(\mathbf{k})} F(\mathbf{i}, s + |\mathbf{k}| - N - |\mathbf{i}|) \quad (1.8)$$

where $s \in \{0, 1, \dots, N-1\}$ and $\mathbf{k} \in I^{(d+1)}(N+1)$ is strictly increasing.

We remind the reader that, by (1.5) in Definition 1.3, the summand on the right hand side in (1.8) is $F_{\wedge}^{(d)}(\mathbf{i}) \otimes X^{d-(s+|\mathbf{k}|-N-|\mathbf{i}|)} Y^{s+|\mathbf{k}|-N-|\mathbf{i}|}$, or written out in full,

$$X^{d-i_1} Y^{i_1} \wedge \dots \wedge X^{d-i_N} Y^{i_N} \otimes X^{d-(s+|\mathbf{k}|-N-|\mathbf{i}|)} Y^{s+|\mathbf{k}|-N-|\mathbf{i}|}.$$

If $\mathbf{i} \in \mathcal{B}(\mathbf{k})$ then

$$\begin{aligned} |\mathbf{i}| &\leq (k_2 - 1) + \dots + (k_{N+1} - 1) = |\mathbf{k}| - k_1 - N \\ |\mathbf{i}| &\geq k_1 + \dots + k_N = |\mathbf{k}| - k_{N+1} \end{aligned} \quad (1.9)$$

and so $s + |\mathbf{k}| - N - |\mathbf{i}| \geq s + k_1 \geq 0$ and $s + |\mathbf{k}| - N - |\mathbf{i}| \leq s + k_{N+1} - N \leq k_{N+1} - 1 \leq d$. Therefore φ is well-defined. As motivation and an aide-memoire, we note that a canonical basis element of Y -degree $s + |\mathbf{k}|$ maps under φ to a sum of canonical basis elements each of Y -degree $s + |\mathbf{k}| - N$. We refer to the space of elements of Y -degree l as the *weight space of weight l* . It is not obvious that φ has image in the subrepresentation $\Delta^{(2,1^{N-1})} \text{Sym}^d E$ of $\bigwedge^N V \otimes V$. We give a short proof of this fact in Lemma 2.1.

Example 1.9.

(a) By (1.8), the canonical basis element

$$X^{N-1} \otimes F_{\wedge}^{(d+1)}(0, 1, \dots, N) = X^{N-1} \otimes X^{d+1} \wedge X^d Y \wedge \dots \wedge X^{d-N+1} Y^N$$

in $\text{Sym}^{N-1} E \otimes \bigwedge^{N+1} \text{Sym}^{d+1} E$ of minimal Y -degree $0 + 1 + \dots + N$ maps under φ to the canonical basis element

$$F((0, 1, \dots, N-1), 0) = X^d \wedge X^{d-1} Y \wedge \dots \wedge X^{d-N+1} Y^{N-1} \otimes X^d$$

in $\Delta^{(2, 1^{N-1})} \text{Sym}^d E$ of minimal Y -degree $0 + 1 + \dots + (N-1)$. Working over \mathbb{C} , these vectors are highest weight for the action of the Lie algebra generator e (which may be thought of as $X \frac{d}{dY}$) in (2.2).

(b) More generally the image of $X^{N-1-s} Y^s \otimes F_{\wedge}^{(d+1)}(i, i+1, \dots, i+N)$ is $F((i, i+1, \dots, i+N-1), s+i)$. Note that since $s \in \{0, \dots, N-1\}$, it follows from (1.2) that this image is in $\Delta^{(2, 1^{N-1})} \text{Sym}^d E$.

(c) The image of a canonical basis element of $\text{Sym}^{N-1} E \otimes \bigwedge^{N+1} \text{Sym}^{d+1} E$ typically has many summands. For instance take $N = 3$ and $d = 5$ and $(s, \mathbf{k}) = (1, (0, 2, 3, 6))$. Then

$$\begin{aligned} & \varphi(XY \otimes X^6 \wedge X^4 Y^2 \wedge X^3 Y^3 \wedge Y^6) \\ &= \sum_{\mathbf{i} \in [0, 2] \times [2, 3] \times [3, 6]} F(\mathbf{i}, 1 + |(0, 2, 3, 6)| - 3 - |\mathbf{i}|) \\ &= F((0, 2, 3), 4) + F((1, 2, 3), 3) + F((0, 2, 4), 3) + F((1, 2, 4), 2) \\ & \quad + F((0, 2, 5), 2) + F((1, 2, 5), 1). \end{aligned}$$

We invite the reader to check that the right-hand side is in $\Delta^{(2, 1, 1)} \text{Sym}^5 E$.

(d) Taking $N = 1$ we may identify $\text{Sym}^0 E$ with \mathbb{F} and $\Delta^{(2)} \text{Sym}^d E$ with the symmetric tensors inside $\text{Sym}^d E \otimes \text{Sym}^d E$. The map $\varphi : \bigwedge^2 \text{Sym}^{d+1} E \rightarrow \Delta^{(2)} \text{Sym}^d E$ is then defined by

$$\varphi(X^{d+1-k} Y^k \wedge X^{d+1-\ell} Y^\ell) = \sum_{k \leq i < \ell} X^{d-i} Y^i \otimes X^{d-(k+\ell-1-i)} Y^{k+\ell-1-i}.$$

It is clear from the powers of Y in the tensor factors on the right-hand side that the right-hand side is a symmetric tensor lying in $\Delta^{(2)} \text{Sym}^d E$. This is an example of the Wronskian isomorphism mentioned at the start of the introduction.

2. THE MAP φ IS AN $\text{SL}_2(\mathbb{F})$ -ISOMORPHISM

2.1. The image of φ . As defined φ has codomain $\bigwedge^N \text{Sym}^d E \otimes \text{Sym}^d E$.

Lemma 2.1. *The image of φ is contained in $\Delta^{(2, 1^{N-1})} \text{Sym}^d E$.*

Proof. Let $s \in \{0, 1, \dots, N-1\}$ and let $\mathbf{k} \in I^{(d+1)}(N+1)$ be strictly increasing. Let Ω be the subset of

$$\mathcal{B}(\mathbf{k}) \times [0, d+1) = [k_1, k_2) \times \dots \times [k_N, k_{N+1}) \times \{0, 1, \dots, d\}$$

of all tuples (i_1, \dots, i_N, j) such that $i_1 + \dots + i_N + j = s + |\mathbf{k}| - N$. Writing elements of Ω as (\mathbf{i}, j) , we have, using Definition 1.8,

$$\mu_N \varphi(X^{N-1-s} Y^s \otimes F_{\wedge}^{(d+1)}(\mathbf{k})) = \sum_{(\mathbf{i}, j) \in \Omega} \mu_N F(\mathbf{i}, j). \quad (2.1)$$

By the definition of $\Delta^{(2, 1^{N-1})} \text{Sym}^d E$ from (1.2), it suffices to show that the right-hand side vanishes. Our proof uses an involution on Ω related to the partition used to prove Lemma 1.6. (See after the proof for the connection with the neighbour map.) First observe that if $(\mathbf{i}, j) \in \Omega$ then $j = s + |\mathbf{k}| - N - |\mathbf{i}|$ and by (1.9) we have

$$k_1 \leq j < k_{N+1}.$$

Thus, given $(\mathbf{i}, j) \in \Omega$, there exists a unique $1 \leq \alpha \leq N$ such that $k_\alpha \leq j < k_{\alpha+1}$. We send (\mathbf{i}, j) to $(i_1, \dots, j, \dots, i_N, i_\alpha)$ where j appears in position α , so j and i_α are swapped. It is clear this defines an involution in which (\mathbf{i}, j) is a fixed point if and only if $i_\alpha = j$. Since $(i_1, \dots, j, \dots, i_N, i_\alpha)$ and $(i_1, \dots, i_\alpha, \dots, i_N, j)$ are either equal or differ by a transposition, their contributions to the sum in (2.1) cancel. \square

To illustrate the neighbour map in Definition 1.2 we remark that the image of (\mathbf{i}, j) under the involution in the proof of Lemma 2.1 is $\mathcal{P}(\mathbf{i}, j)$ if $k_\alpha \leq i_\alpha \leq j < k_{\alpha+1}$ and $\mathcal{P}^{-1}(\mathbf{i}, j)$ if $k_\alpha \leq j < i_\alpha < k_{\alpha+1}$.

2.2. φ is an $\text{SL}_2(\mathbb{F})$ -homomorphism. For $\beta \in \mathbb{N}$ we denote by $\mathbf{u}^{(\beta)}$ the unit vector $(0, \dots, 1, \dots, 0)$ where the non-zero entry is in position β ; the length is always N or $N+1$ and will always be clear from context.

Reduction. We recall the technical trick in [McDW22, §4.2] used to pass from $\text{SL}_2(\mathbb{F})$ to $\text{SL}_2(\mathbb{C})$. First notice that φ is a map of vector spaces, but it is defined over the integers. Let $\gamma \in \mathbb{F}$ be an arbitrary element and $U_\gamma = \begin{pmatrix} 1 & \gamma \\ 0 & 1 \end{pmatrix}$. The elements U_γ and their transposes generate $\text{SL}_2(\mathbb{F})$.

Checking that φ intertwines the action of U_γ (or its transpose) amounts to an equality of polynomials in γ with coefficients in the image of \mathbb{Z} in \mathbb{F} . Clearly, it suffices to check that this equality holds over the polynomial ring $\mathbb{Z}[\gamma]$. For this, in turn, it suffices to prove the equality for *any* transcendental element γ in any field containing \mathbb{Z} as a subring. Proving the result for $\text{SL}_2(\mathbb{C})$ certainly implies the latter condition. A basic fact from Lie theory (see for instance [FH91, Ch. 8]) then reduces the question to proving that φ commutes with the Lie algebra generators e and f of $\mathfrak{sl}_2(\mathbb{C})$, defined on the X, Y basis of E by the matrices

$$e = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad f = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}. \quad (2.2)$$

Their action on $\text{Sym}^d E$ is given by $e \cdot g = X \frac{dg}{dY}$ and $f \cdot g = Y \frac{dg}{dX}$. Their action on $\bigwedge^R \text{Sym}^c E$ is then given by the usual multilinear rule for Lie algebra actions, coming ultimately from

$$x \cdot (u \otimes v) = (x \cdot u) \otimes v + u \otimes (x \cdot v). \quad (2.3)$$

We state it below using the unit vectors $\mathbf{u}^{(\gamma)}$ just defined:

$$\begin{aligned} e \cdot F_{\wedge}^{(c)}(\mathbf{k}) &= \sum_{\alpha=1}^R k_{\alpha} F_{\wedge}^{(c)}(\mathbf{k} - \mathbf{u}^{(\alpha)}) \\ f \cdot F_{\wedge}^{(c)}(\mathbf{k}) &= \sum_{\alpha=1}^R (c - k_{\alpha}) F_{\wedge}^{(c)}(\mathbf{k} + \mathbf{u}^{(\alpha)}) \end{aligned} \quad (2.4)$$

for $\mathbf{k} \in I^{(c)}(R)$. Here we use the convention that if $\mathbf{k} \pm \mathbf{u}^{(\alpha)} \notin I^{(c)}(R)$ then $F_{\wedge}^{(c)}(\mathbf{k} \pm \mathbf{u}^{(\alpha)}) = 0$. An application of (2.3) now gives the Lie algebra action on $\Delta^{(2,1^{N-1})} \text{Sym}^d E \subseteq \bigwedge^N \text{Sym}^d E \otimes \text{Sym}^d E$, which we use in the proof of Lemma 2.3 below.

Technical lemma. The following lemma isolates the key step in the calculation that φ commutes with the Lie algebra action of e . In it we write $\mathcal{B}(\mathbf{k}) - \mathbf{u}^{(\beta)}$ for $\{\mathbf{j} - \mathbf{u}^{(\beta)} : \mathbf{j} \in \mathcal{B}(\mathbf{k})\}$. The notation $F(\mathbf{i}, j)$ was introduced in Definition 1.3. The first paragraph of the proof below checks that, in every summand, the second component of the pair is in $\{0, 1, \dots, d\}$, and so the expression is well-defined. This technical check could be skipped by instead regarding the $F(\mathbf{i}, j)$ as formal symbols; the proof then goes through.

Lemma 2.2. *Let $s \in \{0, 1, \dots, N-1\}$. Let $\mathbf{k} \in I^{(d+1)}(N+1)$ be strictly increasing. Then for any t such that $|\mathbf{k}| - N - 1 \leq t \leq |\mathbf{k}| - 2$ we have*

$$\begin{aligned} &\sum_{\alpha=1}^{N+1} \sum_{\mathbf{i} \in \mathcal{B}(\mathbf{k} - \mathbf{u}^{(\alpha)})} k_{\alpha} F(\mathbf{i}, t - |\mathbf{i}|) \\ &= \sum_{\beta=1}^N \sum_{\mathbf{j} \in \mathcal{B}(\mathbf{k}) - \mathbf{u}^{(\beta)}} (j_{\beta} + 1) F(\mathbf{j}, t - |\mathbf{j}|) + \sum_{\mathbf{j} \in \mathcal{B}(\mathbf{k})} (|\mathbf{k}| - N - |\mathbf{j}|) F(\mathbf{j}, t - |\mathbf{j}|). \end{aligned}$$

Proof. If $\mathbf{i} \in \mathcal{B}(\mathbf{k} - \mathbf{u}^{(\alpha)})$ and $\alpha \neq 1, N+1$ then, by replacing \mathbf{k} with $\mathbf{k} - \mathbf{u}^{(\alpha)}$ in (1.9) we have $|\mathbf{k}| - k_{N+1} - 1 \leq |\mathbf{i}| \leq |\mathbf{k}| - k_1 - N - 1$ and so

$$|\mathbf{k}| - (d+1) - 1 \leq |\mathbf{i}| \leq |\mathbf{k}| - N - 1.$$

Hence $t - |\mathbf{k}| + N + 1 \leq t - |\mathbf{i}| \leq t - |\mathbf{k}| + d + 2$ and the hypothesis on t implies that $0 \leq t - |\mathbf{i}| \leq d$. Thus each $F(\mathbf{i}, t - |\mathbf{i}|)$ is well-defined. This also shows each $F(\mathbf{j}, t - |\mathbf{j}|)$ in the first summand on the right-hand side is well-defined; in the second summand we have instead $t - |\mathbf{k}| + N \leq t - |\mathbf{j}| \leq t - |\mathbf{k}| + d + 1$ and now if $t = |\mathbf{k}| - N - 1$ we may have $-1 = t - |\mathbf{j}|$, but in this case the coefficient is $|\mathbf{k}| - N - |\mathbf{j}| = |\mathbf{k}| - N - (1 + t) = 0$, so the ill-defined summand

can be ignored. We leave the similar verification when $\alpha = 1$ or $\alpha = N + 1$ to the reader. We are now ready to begin the main part of the proof.

Given $x \in \{0, 1, \dots, d\}$ and $1 \leq \alpha \leq N$, we set

$$\mathcal{C}_\alpha^{(x)}(\mathbf{k}) = [k_1, k_2) \times \cdots \times [k_{\alpha-1}, k_\alpha) \times \{x\} \times [k_{\alpha+1}, k_{\alpha+2}) \times \cdots \times [k_N, k_{N+1})$$

where $\{x\}$ in position α replaces the interval $[k_\alpha, k_{\alpha+1})$ in position α of the product defining $\mathcal{B}(\mathbf{k})$ in (1.7). Observe that

$$\begin{aligned} \mathcal{B}(\mathbf{k} - \mathbf{u}^{(1)}) &= [k_1 - 1, k_2) \times [k_2, k_3) \times \cdots \times [k_N, k_{N+1}) = \mathcal{B}(\mathbf{k}) \cup \mathcal{C}_1^{(k_1-1)}, \\ \mathcal{B}(\mathbf{k} - \mathbf{u}^{(N+1)}) &= [k_1, k_2) \times \cdots \times [k_{N-1}, k_N) \times [k_N, k_{N+1} - 1) = \mathcal{B}(\mathbf{k}) \setminus \mathcal{C}_N^{(k_{N+1}-1)} \end{aligned}$$

and, if $2 \leq \alpha \leq N$, then

$$\begin{aligned} \mathcal{B}(\mathbf{k} - \mathbf{u}^{(\alpha)}) &= [k_1, k_2) \times \cdots \times [k_{\alpha-1}, k_\alpha - 1) \times [k_\alpha - 1, k_\alpha) \times \cdots \times [k_N, k_{N+1}) \\ &= \mathcal{B}(\mathbf{k}) \cup \mathcal{C}_\alpha^{(k_\alpha-1)}(\mathbf{k}) \setminus \mathcal{C}_{\alpha-1}^{(k_\alpha-1)}(\mathbf{k}). \end{aligned} \quad (2.5)$$

Thus by setting $\mathcal{C}_0^{(x)}(\mathbf{k}) = \mathcal{C}_{N+1}^{(x)}(\mathbf{k}) = \emptyset$, we may unify the cases so that (2.5) holds for all $1 \leq \alpha \leq N + 1$. By (2.5), the left-hand side in the lemma is

$$\begin{aligned} |\mathbf{k}| \sum_{\mathbf{i} \in \mathcal{B}(\mathbf{k})} F(\mathbf{i}, t - |\mathbf{i}|) &+ \sum_{\alpha=1}^{N+1} \sum_{\mathbf{i} \in \mathcal{C}_\alpha^{(k_\alpha-1)}(\mathbf{k})} k_\alpha F(\mathbf{i}, t - |\mathbf{i}|) \\ &- \sum_{\alpha=1}^{N+1} \sum_{\mathbf{i} \in \mathcal{C}_{\alpha-1}^{(k_\alpha-1)}(\mathbf{k})} k_\alpha F(\mathbf{i}, t - |\mathbf{i}|). \end{aligned} \quad (2.6)$$

Similarly to (2.5) we have

$$\begin{aligned} \mathcal{B}(\mathbf{k}) - \mathbf{u}^{(\beta)} &= [k_1, k_2) \times \cdots \times [k_\beta - 1, k_{\beta+1} - 1) \times \cdots \times [k_N, k_{N+1}) \\ &= \mathcal{B}(\mathbf{k}) \cup \mathcal{C}_\beta^{(k_\beta-1)}(\mathbf{k}) \setminus \mathcal{C}_{\beta+1}^{(k_{\beta+1}-1)}(\mathbf{k}). \end{aligned} \quad (2.7)$$

By (2.7) the first summand in the right side in the lemma is

$$\begin{aligned} \sum_{\beta=1}^N \left(\sum_{\mathbf{j} \in \mathcal{B}(\mathbf{k})} (j_\beta + 1) F(\mathbf{j}, t - |\mathbf{j}|) + \sum_{\mathbf{j} \in \mathcal{C}_\beta^{(k_\beta-1)}(\mathbf{k})} (k_\beta - 1 + 1) F(\mathbf{j}, t - |\mathbf{j}|) \right. \\ \left. - \sum_{\mathbf{j} \in \mathcal{C}_{\beta+1}^{(k_{\beta+1}-1)}(\mathbf{k})} (k_{\beta+1} - 1 + 1) F(\mathbf{j}, t - |\mathbf{j}|) \right). \end{aligned}$$

Since $\sum_{\beta=1}^N (j_\beta + 1) = |\mathbf{j}| + N$, and the second summand on the right-hand side is $\sum_{\mathbf{j} \in \mathcal{B}(\mathbf{k})} (|\mathbf{k}| - N - |\mathbf{j}|) F(\mathbf{j}, t - |\mathbf{j}|)$, the right-hand side in the lemma simplifies to

$$\begin{aligned} |\mathbf{k}| \sum_{\mathbf{j} \in \mathcal{B}(\mathbf{k})} F(\mathbf{j}, t - |\mathbf{j}|) &+ \sum_{\beta=1}^N \sum_{\mathbf{j} \in \mathcal{C}_\beta^{(k_\beta-1)}(\mathbf{k})} k_\beta F(\mathbf{j}, t - |\mathbf{j}|) \\ &- \sum_{\beta=1}^N \sum_{\mathbf{j} \in \mathcal{C}_{\beta+1}^{(k_{\beta+1}-1)}(\mathbf{k})} k_{\beta+1} F(\mathbf{j}, t - |\mathbf{j}|). \end{aligned} \quad (2.8)$$

The lemma now follows by comparing (2.6) and (2.8); since $\mathcal{C}_0^{(x)}(\mathbf{k}) = \mathcal{C}_{N+1}^{(x)}(\mathbf{k}) = \emptyset$ the three summands agree in the order written. \square

The map φ commutes with e . The Lie algebra element $e \in \mathfrak{sl}_2(\mathbb{C})$ (which may be thought of as $X \frac{d}{dY}$) acts on $\text{Sym}^d E$ by $e \cdot X^{d-j} Y^j = j X^{d-j+1} Y^{j-1}$.

Lemma 2.3. *The map φ defined over the complex numbers commutes with the Lie algebra action of $e \in \mathfrak{sl}_2(\mathbb{C})$.*

Proof. We compare $e \cdot \varphi(x)$ and $\varphi(e \cdot x)$ for x in the canonical basis of $\text{Sym}^{N-1} E \otimes \bigwedge^{N+1} \text{Sym}^{d+1} E$. Let $s \in \{0, 1, \dots, N-1\}$ and let $\mathbf{k} \in I^{(d+1)}(N+1)$ be strictly increasing. For ease of notation we set $w = s + |\mathbf{k}| - N$. By (2.3) and (2.4) and the definition of φ in Definition 1.8, then the technical lemma to obtain the third equality, and finally (2.3) and (2.4) again we have

$$\begin{aligned}
& \varphi(e \cdot (X^{N-1-s} Y^s \otimes F_{\wedge}^{(d+1)}(\mathbf{k}))) \\
&= \varphi(s X^{N-s} Y^{s-1} \otimes F_{\wedge}^{(d+1)}(\mathbf{k}) + X^{N-1-s} Y^s \otimes \sum_{\alpha=1}^{N+1} k_{\alpha} F_{\wedge}^{(d+1)}(\mathbf{k} - \mathbf{u}^{(\alpha)})) \\
&= s \sum_{\mathbf{i} \in B(\mathbf{k})} F(\mathbf{i}, s + |\mathbf{k}| - 1 - N - |\mathbf{i}|) \\
&\quad + \sum_{\alpha=1}^{N+1} k_{\alpha} \sum_{\mathbf{i} \in B(\mathbf{k} - \mathbf{u}^{(\alpha)})} F(\mathbf{i}, s + |\mathbf{k}| - 1 - N - |\mathbf{i}|) \\
&= \sum_{\beta=1}^N \sum_{\mathbf{j} \in B(\mathbf{k}) - \mathbf{u}^{(\beta)}} (j_{\beta} + 1) F(\mathbf{j}, w - 1 - |\mathbf{j}|) \\
&\quad + \sum_{\mathbf{j} \in B(\mathbf{k})} (s + |\mathbf{k}| - N - |\mathbf{j}|) F(\mathbf{j}, w - 1 - |\mathbf{j}|) \\
&= \sum_{\beta=1}^N \sum_{\mathbf{i} \in B(\mathbf{k})} i_{\beta} F(\mathbf{i} - \mathbf{u}^{(\beta)}, w - |\mathbf{i}|) + \sum_{\mathbf{j} \in B(\mathbf{k})} (s + |\mathbf{k}| - N - |\mathbf{j}|) F(\mathbf{j}, w - 1 - |\mathbf{j}|) \\
&= e \cdot \sum_{\mathbf{j} \in B(\mathbf{k})} F(\mathbf{j}, s + |\mathbf{k}| - N - |\mathbf{j}|) \\
&= e \cdot \varphi(X^{N-1-s} Y^s \otimes F_{\wedge}^{(d+1)}(\mathbf{k})).
\end{aligned}$$

The hypothesis of the technical lemma that $|\mathbf{k}| - N - 1 \leq w - 1 \leq |\mathbf{k}| - 2$ follows easily from the definition of w . \square

Duality. To show that φ commutes with f we use a duality argument. This appears to the authors to be more conceptual and involve less calculation than adapting the proof already given for e , although this would also be possible. Let $\mathbf{e} = (d+1, \dots, d+1) \in I^{(d+1)}(N+1)$ and define $\tau \in \text{End}_{\mathbb{R}}(\text{Sym}^{N-1} E \otimes \bigwedge \text{Sym}^{N+1} \text{Sym}^{d+1}(E))$ by linear extension of

$$\tau(X^{N-1-j} Y^j \otimes F_{\wedge}^{(d+1)}(\mathbf{i})) = X^j Y^{N-1-j} \otimes F_{\wedge}^{(d+1)}(\mathbf{e} - \mathbf{i}).$$

Let $\mathbf{d} = (d, \dots, d) \in I^{(d)}(N)$ and define $\tau' \in \text{End}_{\mathbb{F}}(\bigwedge^N \text{Sym}^d \otimes \text{Sym}^d E)$ by linear extension of

$$\tau'(F_{\wedge}^{(d)}(\mathbf{j}) \otimes X^{d-\ell} Y^{\ell}) = F_{\wedge}^{(d)}(\mathbf{d} - \mathbf{j}) \otimes X^{\ell} Y^{d-\ell}.$$

Lemma 2.4. *We have $e\tau = \tau f$, $\tau'e = f\tau'$ and $\tau'\varphi = \pm\varphi\tau$.*

Proof. Observe that τ and τ' are defined by multilinear extension of the maps $\theta_c : \text{Sym}^c E \rightarrow \text{Sym}^c E$ defined on the canonical basis by

$$\theta_c(X^{c-j} Y^j) = X^j Y^{c-j}.$$

Since $\theta_c(e \cdot X^{c-j} Y^j) = \theta_c(j X^{c-j+1} Y^{j-1}) = j X^{j-1} Y^{c-j+1} = f \cdot X^j Y^{c-j} = f \cdot (\theta_c(X^{c-j} Y^j))$ we have $\theta_c e = f \theta_c$. By multilinearity, this implies the first two equations in the lemma. For the third, let ε_R denote the sign of the permutation of $\{1, \dots, R\}$ reversing the positions in an R -tuple. (Thus $\varepsilon_R = -1$ if $R \equiv 2, 3 \pmod{4}$, and otherwise $\varepsilon_R = 1$.) Let $s \in \{0, 1, \dots, N-1\}$, let $\mathbf{k} \in I^{(d+1)}(N+1)$ be strictly increasing, and let $\mathbf{k}^{\text{rev}} = (k_{N+1}, \dots, k_1)$ be the reverse of \mathbf{k} . Observe that $\mathbf{e} - \mathbf{k}^{\text{rev}}$ is strictly increasing and $|\mathbf{e} - \mathbf{k}^{\text{rev}}| = (d+1)(N+1) - |\mathbf{k}|$. Set

$$w = |\mathbf{e} - \mathbf{k}^{\text{rev}}| - 1 - s = dN + N + d - s - |\mathbf{k}|. \quad (2.9)$$

By (1.7),

$$\mathcal{B}(\mathbf{e} - \mathbf{k}^{\text{rev}}) = [d+1 - k_{N+1}, d+1 - k_N] \times \dots \times [d+1 - k_2, d+1 - k_1].$$

Thus $(i_1, \dots, i_N) \in \mathcal{B}(\mathbf{e} - \mathbf{k}^{\text{rev}})$ if and only if $(d+1 - i_N, \dots, d+1 - i_1) \in (k_1, k_2] \times \dots \times (k_N, k_{N+1}]$, so if and only if $(d - i_N, \dots, d - i_1) \in \mathcal{B}(\mathbf{k}) = [k_1, k_2] \times \dots \times [k_N, k_{N+1}]$. Using this to step from line 3 to line 4 below, and the definition of φ in (1.8) for the immediately preceding step, we have

$$\begin{aligned} & \varphi\tau(X^{N-1-s} Y^s \otimes F_{\wedge}^{(d+1)}(\mathbf{k})) \\ &= \varphi(X^s Y^{N-1-s} \otimes \varepsilon_{N+1} F_{\wedge}^{(d+1)}(\mathbf{e} - \mathbf{k}^{\text{rev}})) \\ &= \sum_{\mathbf{i} \in \mathcal{B}(\mathbf{e} - \mathbf{k}^{\text{rev}})} \varepsilon_{N+1} F(\mathbf{i}, w - |\mathbf{i}|) \\ &= \varepsilon_{N+1} \sum_{\mathbf{j} \in \mathcal{B}(\mathbf{k})} F((d - j_N, \dots, d - j_1), w - (dN - |\mathbf{j}|)) \\ &= \varepsilon_N \varepsilon_{N+1} \tau' \left(\sum_{\mathbf{j} \in \mathcal{B}(\mathbf{k})} F((j_1, \dots, j_N), d - (w - (dN - |\mathbf{j}|))) \right) \\ &= \varepsilon_N \varepsilon_{N+1} \tau' \left(\sum_{\mathbf{j} \in \mathcal{B}(\mathbf{k})} F((j_1, \dots, j_N), s + |\mathbf{k}| - N - |\mathbf{j}|) \right) \\ &= \varepsilon_N \varepsilon_{N+1} \tau' \varphi(X^{N-1-s} Y^s \otimes F_{\wedge}^{(d+1)}(\mathbf{k})) \end{aligned}$$

where the penultimate equality uses (2.9). Since $\varepsilon_N \varepsilon_{N+1} \in \{-1, 1\}$ only depends on N , this completes the proof. \square

Proposition 2.5. *The map φ defined over the complex numbers is an $\mathfrak{sl}_2(\mathbb{C})$ -homomorphism.*

Proof. By Lemma 2.3, φ commutes with the Lie algebra action of e . By this lemma and Lemma 2.4 we have

$$\varphi f = \varphi \tau \tau f = \varphi \tau e \tau = \pm \tau' \varphi e \tau = \pm \tau' e \varphi \tau = \pm f \tau' \varphi \tau = f \varphi \tau \tau = f \varphi,$$

and so φ also commutes with the Lie algebra action of f . Since $\mathfrak{sl}_2(\mathbb{C})$ is generated by e and f the proposition follows. \square

2.3. The map φ is an $\mathrm{SL}_2(\mathbb{F})$ -isomorphism. Fix $N \in \mathbb{N}$ and $d \in \mathbb{N}_0$. The canonical basis of $\mathrm{Sym}^{N-1} E \otimes \bigwedge^{N+1} \mathrm{Sym}^{d+1} E$ is indexed by pairs (s, \mathbf{k}) with $s \in \{0, 1, \dots, N-1\}$ and $\mathbf{k} \in I^{(d+1)}(N+1)$ strictly increasing. Whenever we write a pair (s, \mathbf{k}) , it satisfies these conditions. By (1.8), the vectors

$$v_{(s, \mathbf{k})} = \sum_{\mathbf{i} \in \mathcal{B}(\mathbf{k})} F(\mathbf{i}, w - |\mathbf{i}|) \quad (2.10)$$

where $w = s + |\mathbf{k}| - N$ are the images under φ of the canonical basis of $\mathrm{Sym}^{N-1} E \otimes \bigwedge^{N+1} \mathrm{Sym}^{d+1} E$.

By Lemma 2.1, the map φ has image contained in $\Delta^{(2, 1^{N-1})} \mathrm{Sym}^d E$ and by Proposition 1.7, $\Delta^{(2, 1^{N-1})} \mathrm{Sym}^d E$ has the same dimension as the domain of φ . Therefore to complete the proof that φ is an isomorphism of $\mathrm{SL}_2(\mathbb{F})$ -representations, it suffices to show that the map is injective. To this end, we will define a total order \preceq on the basis elements $F_{\Delta}(\mathbf{i}, j)$, and we will show that the vectors $v_{(s, \mathbf{k})}$ are lower-triangular in this basis.

Preliminary results on chains. Our proof of this requires a close analysis of the neighbour map in Definition 1.2 and the chains in Lemma 1.4. Recall from Definition 1.2 that the content of a pair (\mathbf{i}, j) is the multiset $\{i_1, \dots, i_N\} \cup \{j\}$.

Lemma 2.6. *Suppose that $(\mathbf{i}, j) \in I^{(d)}(N) \times \{0, 1, \dots, d\}$ is a semistandard pair such that $F(\mathbf{i}, j)$ has a non-zero coefficient in $v_{(s, \mathbf{k})}$ and such that $j \notin \{i_1, \dots, i_N\}$. Let $(\mathbf{i}', j') = \mathcal{P}^m(\mathbf{i}, j)$ for some $m \in \mathbb{N}$, and suppose that $F(\mathbf{i}', j')$ has a non-zero coefficient in $v_{(s, \mathbf{k})}$. Then $m = 1$.*

Proof. By the definition of the neighbour map, \mathbf{i}' is strictly increasing, and since (\mathbf{i}, j) and (\mathbf{i}', j') have the same content, we may write

$$\begin{aligned} (\mathbf{i}, j) &= ((i_1, \dots, i_{\alpha-1}, i_{\alpha}, i_{\alpha+1}, \dots, i_N), j) \\ (\mathbf{i}', j') &= ((i_1, \dots, i_{\alpha-1}, j, i_{\alpha+1}, \dots, i_N), i_{\alpha}) \end{aligned}$$

for some $\alpha \in \{1, \dots, N\}$. Since $j \notin \{i_1, \dots, i_N\}$, by Lemma 1.4, the neighbour map \mathcal{P} is injective on pairs (\mathbf{i}, j) of content $\{i_1, \dots, i_N, j\}$. We claim that it suffices to prove that $(\mathbf{i}', j') = \mathcal{P}(\mathbf{i}, j)$. Indeed, this would imply

that $\mathcal{P}(\mathbf{i}, j) = \mathcal{P}(\mathcal{P}^{m-1}(\mathbf{i}, j))$, and by injectivity, $\mathcal{P}^{m-1}(\mathbf{i}, j) = (\mathbf{i}, j)$. By Lemma 1.4, this implies that $m - 1 = 0$. The claim now follows.

To see that $(\mathbf{i}', j') = \mathcal{P}(\mathbf{i}, j)$, we need to show that α is maximal such that $i_\alpha \leq j$. It is clear from Lemma 1.4 that repeated applications of \mathcal{P} replace entries in the first element of the semistandard pair (\mathbf{i}, j) by larger or equal entries. Thus, $j \geq i_\alpha$. If $\alpha = N$, we are done. Otherwise, since $\mathbf{i}' \in \mathcal{B}(\mathbf{k})$, we have $j < k_{\alpha+1} \leq i_{\alpha+1}$. Since $i_\alpha \leq i_{\alpha+1}$, the result now follows. \square

By Proposition 1.7, $\Delta^{(2, 1^{N-1})} \text{Sym}^d E$ has as a basis the $F_\Delta(\mathbf{i}, j)$ for semistandard pairs $(\mathbf{i}, j) \in I^{(d)}(N) \times \{0, 1, \dots, d\}$.

Lemma 2.7. *In the expression of $v_{(s, \mathbf{k})}$ as a linear combination of the $F_\Delta(\mathbf{i}, j)$ for semistandard pairs $(\mathbf{i}, j) \in I^{(d)}(N) \times \{0, 1, \dots, d\}$ there is at most one (\mathbf{i}, j) of any given content.*

Proof. If $j \in \{i_1, \dots, i_N\}$ then the unique basis element from the F_Δ basis whose semistandard pair has content $\{i_1, \dots, i_N\} \cup \{j\}$ is $F_\Delta(\mathbf{i}, j)$; since this element is $F(\mathbf{i}, j)$, in this case, the lemma obviously holds.

In the remaining case, the content multiset is a set, and by Lemma 1.4, there are N semistandard pairs of content $\{i_1, \dots, i_N\} \cup \{j\}$ forming a chain

$$(\mathbf{i}^{(1)}, j^{(1)}), \dots, (\mathbf{i}^{(\beta)}, j^{(\beta)}), \dots, (\mathbf{i}^{(\gamma)}, j^{(\gamma)}), \dots, (\mathbf{i}^{(N)}, j^{(N)}) \quad (2.11)$$

with $d \geq j^{(1)} > \dots > j^{(N)} \geq 1$. (To be explicit, Lemma 1.4 gives $j^{(1)} = j$ and $j^{(\alpha)} = i_{N-\alpha+2}$ for $2 \leq \alpha \leq N$.) Choose β minimal such that $F_\Delta(\mathbf{i}^{(\beta)}, j^{(\beta)})$ has a non-zero coefficient in $v_{(s, \mathbf{k})}$ and γ maximal such that $F_\Delta(\mathbf{i}^{(\gamma)}, j^{(\gamma)})$ has a non-zero coefficient in $v_{(s, \mathbf{k})}$.

Observe that $F(\mathbf{i}^{(\beta)}, j^{(\beta)})$ and $F(\mathcal{P}(\mathbf{i}^{(\gamma)}, j^{(\gamma)})) = F(\mathcal{P}^{\gamma-\beta+1}(\mathbf{i}^{(\beta)}, j^{(\beta)}))$ both have a non-zero coefficient in $v_{(s, \mathbf{k})}$ in the F basis. By Lemma 2.6 applied to $(\mathbf{i}^{(\beta)}, j^{(\beta)})$, we have $\beta = \gamma$, as required. \square

Triangularity. It will be useful to denote the content multiset of a semistandard pair $(\mathbf{i}, j) \in I^{(d)}(N) \times \{0, 1, \dots, N\}$ by $c(\mathbf{i}, j)$. Given distinct multisets $\{a_1, \dots, a_{N+1}\}$ and $\{b_1, \dots, b_{N+1}\}$ written so that $a_1 \leq \dots \leq a_{N+1}$ and $b_1 \leq \dots \leq b_{N+1}$, we order them lexicographically so that

$$\{a_1, \dots, a_{N+1}\} < \{b_1, \dots, b_{N+1}\}$$

if and only if $a_1 = b_1, \dots, a_{\gamma-1} = b_{\gamma-1}$ and $a_\gamma < b_\gamma$, where γ is minimal such that $a_\gamma \neq b_\gamma$. Using this we define a total order \preceq on semistandard pairs.

Definition 2.8. Let \prec be the total order on semistandard pairs in $I^{(d)}(N) \times \{0, 1, \dots, d\}$ defined by $(\mathbf{i}, j) \prec (\mathbf{i}', j')$ if and only if *either* $c(\mathbf{i}, j) < c(\mathbf{i}', j')$ *or* $c(\mathbf{i}, j) = c(\mathbf{i}', j')$ and $j > j'$.

For example the least element under \preceq is $((0, 1, \dots, N-1), 0)$ and the greatest is $((d-N+1, \dots, d), d)$. The final condition $j > j'$ is chosen so

that the chain in (2.11) is strictly increasing in the \preceq total order. This can be seen in the following example.

Example 2.9. Take $N = 2$ and $d = 4$. The map φ takes the canonical basis element in $E \otimes \bigwedge^3 \text{Sym}^5 E$ element associated to the pair (s, \mathbf{k}) and having weight 9 (in the sense defined before Example 1.9) to a sum $v_{(s, \mathbf{k})}$ of canonical basis elements of $\bigwedge^2 \text{Sym}^4 E \otimes \text{Sym}^4 E$ each of weight 7. The map φ , restricted to this weight space is shown in the matrix below. Each column contains the coefficients of the image of the element labelled $(s, (k_1, k_2, k_3))$ on top, corresponding to the elements labelled $((i_1, i_2), j)$ in each row.

The labels $((i_1, i_2), j)$ correspond to elements $F_\Delta((i_1, i_2), j)$. They are totally ordered

$$((0, 3), 4) \prec ((0, 4), 3) \prec ((1, 2), 4) \prec ((1, 4), 2) \prec ((1, 3), 3) \prec ((2, 3), 2)$$

by \preceq . By Lemma 2.7, at most one $F_\Delta(\mathbf{i}, j)$ of any given content appears in each column. This can be seen in the two 2×2 identity blocks in the relevant block of the matrix. We deliberately use a non-standard notation in which \cdot denotes an entry known to be zero by Lemma 2.7. Empty spaces also denote zeros, showing lower-triangularity in this example.

$$\begin{array}{c} \begin{array}{c} \begin{array}{c} (1, (0, 3, 5)) \\ (0, (0, 4, 5)) \\ (1, (1, 2, 5)) \\ (0, (1, 3, 5)) \\ (1, (1, 3, 4)) \\ (0, (2, 3, 4)) \end{array} \end{array} \\ \begin{array}{c} ((0, 3), 4) \\ ((0, 4), 3) \\ ((1, 2), 4) \\ ((1, 4), 2) \\ ((1, 3), 3) \\ ((2, 3), 2) \end{array} \begin{pmatrix} 1 & & & & & \\ \cdot & 1 & & & & \\ 0 & 0 & 1 & & & \\ 1 & 1 & \cdot & 1 & & \\ 1 & 0 & 1 & 1 & 1 & \\ 1 & 0 & 0 & 1 & 1 & 1 \end{pmatrix} \end{array}$$

The uni-triangularity seen in the previous example holds more generally, as the following proposition shows.

Proposition 2.10. *Let $(\mathbf{i}, j) \in I^{(d)}(N) \times \{0, 1, \dots, d\}$ be semistandard. Let α be maximal such that $i_\alpha \leq j$. Set $\mathbf{k} = (i_1, \dots, i_\alpha, j+1, i_{\alpha+1}+1, \dots, i_N+1)$. Then*

$$v_{(\alpha-1, \mathbf{k})} = F_\Delta(\mathbf{i}, j) + v$$

where v is a linear combination of basis elements $F_\Delta(\mathbf{i}', j')$ for semistandard (\mathbf{i}', j') such that $(\mathbf{i}', j') \succ (\mathbf{i}, j)$. Moreover, the correspondence between pairs $(\alpha-1, \mathbf{k})$ and semistandard pairs (\mathbf{i}, j) is one-to-one, so the vectors $v_{(\alpha-1, \mathbf{k})}$ are linearly independent.

Proof. We first show that we may recover the pair (\mathbf{i}, j) from α and \mathbf{k} . We have $j = k_{\alpha+1} - 1$. On the other hand,

$$i_1 = k_1, \dots, i_\alpha = k_\alpha, i_{\alpha+1} = k_{\alpha+2} - 1, \dots, i_N = k_{N+1} - 1 \quad (2.12)$$

This also shows that $\mathbf{i} \in \mathcal{B}(\mathbf{k})$. Since the quantity $s + |\mathbf{k}| - N - |\mathbf{i}|$ in the definition of φ in (1.8) is

$$(\alpha - 1) + |\mathbf{k}| - N - |\mathbf{i}| = (\alpha - 1) + (|\mathbf{i}| + N - \alpha + j + 1) - N - |\mathbf{i}| = j.$$

by (1.8), we have that $F(\mathbf{i}, j)$ has coefficient 1 in the expression of $v_{(\alpha-1, \mathbf{k})}$ in the F basis. Moreover, we have

$$\mathcal{P}(\mathbf{i}, j) = ((i_1, \dots, i_{\alpha-1}, j, i_{\alpha+1}, \dots, i_N), i_\alpha)$$

and since $k_\alpha = i_\alpha \leq j = k_{\alpha+1} - 1$, we also have $(i_1, \dots, i_{\alpha-1}, j, i_{\alpha+1}, \dots, i_N) \in \mathcal{B}(\mathbf{k})$. Therefore $F(\mathcal{P}(\mathbf{i}, j))$ has coefficient 1 in the expression of $v_{(\alpha-1, \mathbf{k})}$ in the F basis. It follows by Lemma 2.7 that $F_\Delta(\mathbf{i}, j)$ has coefficient 1 when $v_{(\alpha-1, \mathbf{k})}$ is written in the F_Δ basis, and the remaining summands in the F_Δ basis have different content to (\mathbf{i}, j) .

Let $F_\Delta(\mathbf{i}', j')$ be one of these summands with non-zero coefficient in $v_{(\alpha-1, \mathbf{k})}$. By Lemma 2.6, we see that $F(\mathbf{i}', j')$ and possibly $F(\mathcal{P}(\mathbf{i}', j'))$ are the only F -basis elements of content $c(\mathbf{i}', j')$ with non-zero coefficient in $v_{(\alpha-1, \mathbf{k})}$. It follows that $\mathbf{i}' \in \mathcal{B}(\mathbf{k})$.

Suppose for a contradiction that (\mathbf{i}', j') satisfies $(\mathbf{i}', j') \prec (\mathbf{i}, j)$, or equivalently, since $c(\mathbf{i}', j') \neq c(\mathbf{i}, j)$, that $c(\mathbf{i}', j') < c(\mathbf{i}, j)$. Let

$$\begin{aligned} \mathbf{c} &= (c_1, \dots, c_{N+1}) = (i_1, \dots, i_\alpha, j, i_{\alpha+1}, \dots, i_N) \\ \mathbf{c}' &= (c'_1, \dots, c'_{N+1}) = (i'_1, \dots, i'_\beta, j', i'_{\beta+1}, \dots, i'_N) \end{aligned}$$

where β is maximal such that $i'_\beta \leq j'$. Thus both \mathbf{c} and \mathbf{c}' are (weakly) increasing.

Suppose first of all that $\alpha < \beta$. From (2.12) we have $i_\gamma = k_\gamma$ if $1 \leq \gamma \leq \alpha$ and by definition of \mathbf{k} , we have $k_{\alpha+1} = j + 1$. Thus $\mathbf{i}' \in \mathcal{B}(\mathbf{k})$ implies that $c'_\gamma = i'_\gamma \geq k_\gamma = i_\gamma = c_\gamma$ for $1 \leq \gamma \leq \alpha$ and $c'_{\alpha+1} = i'_{\alpha+1} \geq k_{\alpha+1} = j + 1$. Since we are assuming that $c(\mathbf{i}', j') < c(\mathbf{i}, j)$, we must have $c'_\gamma = c_\gamma$ for $1 \leq \gamma \leq \alpha$. However, $c'_{\alpha+1} \geq j + 1 > j = c_{\alpha+1}$. Thus $c(\mathbf{i}', j') > c(\mathbf{i}, j)$, a contradiction.

In the remaining case we have $\alpha \geq \beta$. Similarly to the first case, we have $c'_\gamma = i'_\gamma \geq k_\gamma = i_\gamma = c_\gamma$ for $1 \leq \gamma \leq \beta$, and thus our assumption that $c(\mathbf{i}', j') < c(\mathbf{i}, j)$ implies that $c'_\gamma = c_\gamma$ for $1 \leq \gamma \leq \beta$ and $c'_{\beta+1} \leq c_{\beta+1}$. Since (\mathbf{i}, j) and (\mathbf{i}', j') lie in the same weight space, we have $|\mathbf{i}| + j = |\mathbf{i}'| + j'$. Since $c(\mathbf{i}, j) \neq c(\mathbf{i}', j')$ and $|\mathbf{i}| + j = |\mathbf{i}'| + j'$, there exists δ such that $\beta + 2 \leq \delta \leq N + 1$ and $c'_\delta > c_\delta$. Now, since $\delta \geq \beta + 2$, we have $c'_\delta = i'_{\delta-1}$. In turn, $\mathbf{i}' \in \mathcal{B}(\mathbf{k})$ implies $i'_{\delta-1} < k_\delta$. By definition of \mathbf{k} , we also have $k_\gamma \in \{c_\gamma, c_\gamma + 1\}$ for $1 \leq \gamma \leq N + 1$. In particular, $k_\delta \leq c_\delta + 1$. Putting all these together, we have

$$c'_\delta = i'_{\delta-1} < k_\delta \leq c_\delta + 1$$

and hence $c'_\delta \leq c_\delta$, a final contradiction.

Therefore, any $F_\Delta(\mathbf{i}', j')$ appearing with non-zero coefficient in the expression of $v_{(\alpha-1, \mathbf{k})}$ in the F_Δ basis satisfies $(\mathbf{i}', j') \succ (\mathbf{i}, j)$, as required. \square

The matrix representing φ in the basis $v_{(\alpha-1, \mathbf{k})}$ of the domain $\mathrm{Sym}^{N-1} E \otimes \bigwedge^{N+1} \mathrm{Sym}^{d+1} E$ and the basis $F_\Delta(\mathbf{i}, j)$ is therefore uni-triangular when the two bases are paired by the one-to-one correspondence established in Proposition 2.10. In particular, φ is injective. This concludes the proof of Theorem 1.1.

3. FINAL REMARKS

In this section we first show that the $\mathrm{SL}_2(\mathbb{F})$ isomorphism in Theorem 1.1 becomes a $\mathrm{GL}_2(\mathbb{F})$ -isomorphism provided a suitable power of the determinant is introduced. (This is typical of the general theory: see [PW21, §3.3].) We then obtain identity (1.1) by taking characters. We finish with a conjectured generalization of Theorem 1.1.

We denote the 1-dimensional determinant representation of $\mathrm{GL}_2(\mathbb{F})$ by \det and regard the domain and codomain of φ as representations of $\mathrm{GL}_2(\mathbb{F})$ in the obvious way.

Corollary 3.1. *Let $N \in \mathbb{N}$ and let $d \in \mathbb{N}_0$. The map φ defined in Definition 1.8 is an isomorphism of $\mathrm{GL}_2(\mathbb{F})$ -representations*

$$\mathrm{Sym}^{N-1} E \otimes \bigwedge^{N+1} \mathrm{Sym}^{d+1} E \cong \det^N \otimes \Delta^{(2, 1^{N-1})} \mathrm{Sym}^d E.$$

Proof. Let K be the algebraic closure of \mathbb{F} . Let $\tilde{E} = E \otimes_{\mathbb{F}} K$. It is sufficient to prove that the map

$$\tilde{\varphi} : \mathrm{Sym}^{N-1} \tilde{E} \otimes \mathrm{Sym}^{N+1} \mathrm{Sym}^{d+1} \tilde{E} \rightarrow \det^N \otimes \Delta^{(2, 1^{N-1})} \mathrm{Sym}^d \tilde{E}$$

is a $\mathrm{GL}_2(K)$ -isomorphism, since φ is defined with coefficients in the prime subfield of \mathbb{F} , and so via the inclusion $E \mapsto E \otimes 1 \subseteq E \otimes_{\mathbb{F}} K$, the map $\tilde{\varphi}$ restricts to φ . By Theorem 1.1 for the field K , the map $\tilde{\varphi}$ is an $\mathrm{SL}_2(K)$ -homomorphism. Now, because K is algebraically closed, and so every element of K has a square root in K , we have

$$\mathrm{GL}_2(K) = \left\langle \mathrm{SL}_2(K), \begin{pmatrix} \alpha & 0 \\ 0 & \alpha \end{pmatrix} : \alpha \in K \setminus \{0\} \right\rangle.$$

It therefore suffices to prove that $\tilde{\varphi}$ commutes with the action of the diagonal matrices αI for $\alpha \in K$. Using the canonical bases of the domain and codomain of $\tilde{\varphi}$, one sees that on the domain αI acts as $\alpha^{N-1+(N+1)(d+1)} = \alpha^{(N+1)d+2N}$ and on the codomain αI acts as $\alpha^{(N+1)d} \det(\alpha I)^N = \alpha^{(N+1)d} \alpha^{2N}$. Since the exponents agree, this completes the proof. \square

We now prove identity (1.1). Recall that s_λ is the Schur function canonically labelled by the partition λ . It is immediate from the combinatorial definition of Schur functions (see for instance [S99, Definition 7.10.1]) that $s_\lambda(1, q, \dots, q^d)$ is the generating function enumerating semistandard tableaux of shape λ with entries from $\{0, 1, \dots, d\}$ by their sum of entries.

This gives a combinatorial interpretation of the right-hand side in (1.1) and in Corollary 3.2 below. One of the most natural interpretations of the q -binomial coefficient $\begin{bmatrix} a \\ b \end{bmatrix}_q$ is that $q^{\frac{b(b-1)}{2}} \begin{bmatrix} a \\ b \end{bmatrix}_q$ is the generating function enumerating b -subsets of $\{0, \dots, a-1\}$ by their sum of entries. Thus, by identifying semistandard tableaux of shape (1^{N+1}) with the subset of their entries, we deduce that

$$q^{\frac{(N+1)N}{2}} \begin{bmatrix} d+2 \\ N+1 \end{bmatrix}_q = s_{(1^{N+1})}(1, q, \dots, q^{d+1}). \quad (3.1)$$

For further background on q -binomial coefficients, including the theorem that $\begin{bmatrix} a \\ b \end{bmatrix}_q$ is the generating function enumerating partitions in the $b \times (a-b)$ box by their size, we refer the reader to [S11, §1.7].

Corollary 3.2. *For any $N \in \mathbb{N}$ and $d \in \mathbb{N}_0$ we have*

$$q^{\frac{N(N-1)}{2}} [N]_q \begin{bmatrix} d+2 \\ N+1 \end{bmatrix}_q = s_{(2, 1^{N-1})}(1, q, \dots, q^d).$$

Proof. It suffices to prove the identity when q is a non-zero complex number. It is clear from the canonical basis $X^{N-1}, X^{N-1}Y, \dots, Y^{N-1}$ of $\text{Sym}^{N-1}E$ that $[N]_q = 1 + q + \dots + q^{N-1}$ is the character of $\text{Sym}^{N-1}E$ evaluated at the diagonal matrix D in $\text{GL}_2(\mathbb{C})$ with entries 1 and q . By [PW21, (10)], $s_{(1^{N+1})}(1, q, \dots, q^{d+1})$ and $s_{(2, 1^{N-1})}(1, q, \dots, q^d)$ are the characters of the $\text{GL}_2(\mathbb{C})$ -representations $\bigwedge^{N+1} \text{Sym}^{d+1}E$ and $\Delta^{(2, 1^{N-1})} \text{Sym}^dE$, also evaluated at D . Therefore, by (3.1), the character of $\text{Sym}^{N-1}E \otimes \bigwedge^{N+1} \text{Sym}^{d+1}E$ evaluated at D is $[N]_q q^{\frac{(N+1)N}{2}} \begin{bmatrix} d+2 \\ N+1 \end{bmatrix}_q$. But in Corollary 3.1 we showed that this representation is isomorphic to $\det^N \otimes \Delta^{(2, 1^{N-1})}E$. Hence, equating the character values we obtain

$$[N]_q q^{\frac{(N+1)N}{2}} \begin{bmatrix} d+2 \\ N+1 \end{bmatrix}_q = q^N s_{(2, 1^{N-1})}(1, q, \dots, q^d).$$

The result follows by cancelling q^N from each side. \square

Our main result, Theorem 1.1, is the special case when $M = 1$ of the following conjecture.

Conjecture 3.3. *Let $M, N \in \mathbb{N}$. There is an isomorphism of $\text{SL}_2(\mathbb{F})$ -representations*

$$\bigwedge^M \text{Sym}^{M+N-2}E \otimes \bigwedge^{M+N} \text{Sym}^{M+d}E \cong \Delta^{(M+1, 1^{N-1})} \text{Sym}^dE.$$

If $M = 0$, then the first factor is \mathbb{F} and since $\Delta^{(1^N)}V = \bigwedge^N V$, both sides in the claimed isomorphism are $\bigwedge^N \text{Sym}^dE$. If $N = 1$ then since $\text{Sym}^{M-1}E$ is M -dimensional, and so $\bigwedge^M \text{Sym}^{M-1}E$ is the determinant representation of $\text{SL}_2(\mathbb{F})$, which is trivial, and $\Delta^{(M+1)}V = \text{Sym}_{M+1}V$, the claimed isomorphism is $\bigwedge^{M+1} \text{Sym}^{M+d}E \cong \text{Sym}_{M+1} \text{Sym}^dE$. An explicit isomorphism

from the right-hand side to the left-hand side is given by Theorem 1.4 in [McDW22].

More broadly, it would be interesting to have field-independent results on the endomorphism rings of the two sides in Conjecture 3.3.

ACKNOWLEDGEMENTS

The first author would like to thank his advisor Mikhail Khovanov for valuable discussions. The second author gratefully acknowledges financial support from the Heilbronn Institute for Mathematical Research, Bristol, UK. The authors are very grateful to the anonymous referee for an exceptionally careful reading of the first version of this paper and its revision and many helpful comments and corrections. The authors also thank Álvaro Gutiérrez Cáceres and Michał Szwejk for helpful comments and corrections.

REFERENCES

- [dBPW21] Melanie de Boeck, Rowena Paget, and Mark Wildon. Plethysms of symmetric functions and highest weight representations. *Trans. Amer. Math. Soc.*, **374** 8013–8043, 2021.
- [EGS08] K. Erdmann, J.A. Green, and M. Schocker. *Polynomial representations of GL_n : with an appendix on Schensted Correspondence and Littelmann Paths*, volume 830 of *Lecture Notes in Mathematics*. Springer, 2008.
- [FH91] William Fulton and Joe Harris, *Representation theory: A first course*, Graduate Texts in Mathematics, vol. 129, Springer-Verlag, New York, 1991, Readings in Mathematics.
- [McDW22] Eoghan McDowell and Mark Wildon. Modular plethystic isomorphisms for two-dimensional linear groups *J. Alg.*, **602** 441–483, 2022.
- [PW21] Rowena Paget and Mark Wildon. Plethysms of symmetric functions and representations of $SL_2(\mathbb{C})$. *Algebr. Comb.*, **4** 27–68, 2021.
- [S11] Richard P. Stanley, *Enumerative combinatorics. Vol. 1*, 2nd edition, Cambridge Studies in Advanced Mathematics, vol. 49, Cambridge University Press, Cambridge, 2011.
- [S99] Richard P. Stanley, *Enumerative combinatorics. Vol. 2*, Cambridge Studies in Advanced Mathematics, vol. 62, Cambridge University Press, Cambridge, 1999, With a foreword by Gian-Carlo Rota and appendix 1 by Sergey Fomin.

Email address: `martinez@math.columbia.edu`

Email address: `mark.wildon@bristol.ac.uk`