

**SYLOW SUBGROUPS OF SYMMETRIC AND
ALTERNATING GROUPS AND THE VERTEX
OF $S^{(kp-p, 1^p)}$ IN CHARACTERISTIC p**

EUGENIO GIANNELLI, KAY JIN LIM, AND MARK WILDON

ABSTRACT. We show that the Sylow p -subgroups of a symmetric group, respectively an alternating group, are characterized as the p -subgroups containing all elementary abelian p -subgroups up to conjugacy of the symmetric group, respectively the alternating group. We apply the characterization result for symmetric groups to compute the vertices of the hook Specht modules associated to the partition $(kp - p, 1^p)$ under the assumption that $k \equiv 1 \pmod{p}$ and $k \not\equiv 1 \pmod{p^2}$.

1. INTRODUCTION

In this paper we prove two main theorems. The first gives a new characterization of the Sylow p -subgroups of the symmetric groups \mathfrak{S}_n and alternating groups \mathfrak{A}_n .

Theorem 1.1. *Suppose that G is either \mathfrak{A}_n or \mathfrak{S}_n . Let p be a prime and let $Q \leq G$ be a p -subgroup. Then Q is a Sylow p -subgroup of G if and only if Q contains a G -conjugate of every elementary abelian p -subgroup of G .*

It is easily seen that this characterization of Sylow subgroups does not hold in a general finite group. When n is large compared to p , the Sylow p -subgroups of \mathfrak{S}_n and \mathfrak{A}_n are much larger than the elementary abelian p -groups they contain. For example, if n is a power of 2 then a Sylow 2-subgroup of \mathfrak{S}_n has order 2^{n-1} , whereas the order of the largest elementary abelian 2-subgroup of \mathfrak{S}_n is $2^{n/2}$. This gives another indication that Theorem 1.1 is a non-trivial result. We state some open problems suggested by this theorem in the final section of the paper.

Date: December 7, 2015.

2010 Mathematics Subject Classification. Primary 20C30, Secondary 20B30, 20C20, 20D06, 20D20.

Key words and phrases. Specht module, vertex, symmetric group, alternating group, Sylow subgroup.

Part of this work was done while the third author was visiting the Institute for Mathematical Sciences, National University of Singapore in 2013. The visit was supported by the Institute.

The second author is supported by Singapore Ministry of Education AcRF Tier 1 grant RG13/14.

Theorem 1.1 is motivated by an application to vertices of indecomposable modules for the symmetric group. We recall the key definitions. Let F be a field of prime characteristic p , let G be a finite group and let M be an indecomposable FG -module. A subgroup Q of G is said to be a *vertex* of M if there exists a FQ -module U such that M is a direct summand of $U \uparrow^G$ and Q is minimal with this property. This definition was introduced by Green in [13], where he showed that the vertices of M form a single conjugacy class of p -subgroups of G . We note that the vertices of M do not change under field extensions of F [17, Theorem 1.21]. Vertices play a key role in the Green correspondence [14], and in Alperin's Weight Conjecture [3]. But despite the importance of vertices to local representation theory, there are relatively few cases where the vertices of particular families of modules have been computed. Our second main theorem is as follows.

Theorem 1.2. *Let p be an odd prime. Suppose that $k \equiv 1 \pmod{p}$ and that $k \not\equiv 1 \pmod{p^2}$. The Specht module $S^{(kp-p, 1^p)}$, defined over a field of prime characteristic p , has a Sylow p -subgroup of \mathfrak{S}_{kp} as a vertex.*

We prove Theorem 1.2 as a corollary of the following result, which is of independent interest.

Theorem 1.3. *Let p be an odd prime and let E be an elementary abelian p -subgroup of \mathfrak{S}_{kp} . If either*

- (i) *E has at least two orbits of size p on $\{1, \dots, kp\}$, or*
- (ii) *E has at least one orbit of size p^2 on $\{1, \dots, kp\}$,*

then E is contained in a vertex of the Specht module $S^{(kp-p, 1^p)}$, defined over an algebraically closed field of characteristic p .

The proof of Theorem 1.3 uses two key techniques: generic Jordan type for modules over elementary abelian p -groups (see [11, 26]), and the Brauer correspondence for modules, as developed by Broué in [5]. The isomorphism $S^{(kp-p, 1^p)} \cong \bigwedge^p S^{(kp-1, 1)}$ (see Section 5) is used throughout the calculations.

Background. We now survey the existing results on vertices of Specht modules, emphasising the different techniques that have been used. We begin with results that, like Theorem 1.2, apply to Specht modules labelled by partitions of a special form. Note that, by [18, Corollary 13.18], Specht modules are always indecomposable in odd characteristic.

The simplest case is a Specht module S^λ in a block of abelian defect. In this case S^λ has dimension coprime to p and by [13, Corollary 1], its vertices are the defect groups of its block; these are elementary abelian.

In [27, Theorem 2] the third author used the Brauer correspondence to show that when p does not divide n and $S^{(n-r, 1^r)}$ is indecomposable, its vertex is a Sylow p -subgroup of $\mathfrak{S}_{n-r-1} \times \mathfrak{S}_r$. When $p = 2$ and n is even

it was shown by Murphy and Peel in [23, Theorem 4.5] that $S^{(n-r, 1^r)}$ is indecomposable and has a Sylow 2-subgroup of \mathfrak{S}_n as a vertex. A key step in their proof is their Theorem 4.3, which states that if Q is an elementary abelian 2-subgroup of \mathfrak{S}_n of rank $n/2$ then $S^{(n-r, 1^r)} \downarrow_Q$ is indecomposable. The analogue of this result for odd primes is in general false: for example, when $p = 5$, calculations with the computer algebra package MAGMA [4] show that if $Q = \langle (1, 2, 3, 4, 5), (6, 7, 8, 9, 10) \rangle$ then $S^{(8, 1, 1)} \downarrow_Q$ is decomposable. It therefore seems impossible to extend the methods of [23] to the case when p is odd and p divides n .

The dimension of $S^{(kp-r, 1^r)}$ is $\binom{kp-1}{r}$, which is coprime to p whenever $r < p$. In these cases $S^{(kp-r, 1^r)}$ has a full Sylow p -subgroup of \mathfrak{S}_{kp} as a vertex again by [13, Corollary 1]. When $r = p$ the dimension is divisible by p if and only if $k \equiv 1 \pmod{p}$, and divisible by p^2 if and only if $k \equiv 1 \pmod{p^2}$. Theorem 1.2 therefore deals with the smallest case that cannot be decided solely by dimension arguments.

When $p = 2$ the Specht module $S^{(n-2, 2)}$ is always indecomposable. When n is even or $n \equiv 1 \pmod{4}$, it was shown in [6, Theorem 1] that $S^{(n-2, 2)}$ has a Sylow 2-subgroup of \mathfrak{S}_n as a vertex, except when $n = 4$, when the unique vertex of $S^{(2, 2)}$ is the normal Sylow 2-subgroup of \mathfrak{A}_4 . When $n \equiv 3 \pmod{4}$, any Sylow 2-subgroup of $\mathfrak{S}_{n-5} \times \mathfrak{S}_2 \times \mathfrak{S}_2$ is a vertex. When $p \geq 3$ the third author showed in [28, Theorem 8.1] that $S^{(n-2, 2)}$ has a Sylow p -subgroup of the defect group of its block as a vertex.

In the main result of [28] the third author used the Brauer correspondence to determine a subgroup contained in the vertex of each indecomposable Specht module. This result was later improved by the first author in [12]. In [21] the second author used the complexity theory of modules to prove a number of classification theorems on Specht modules with abelian vertices. In particular it was shown that if $p \geq 3$ and S^λ has an abelian vertex of p -rank m then λ is a p^2 -core of p -weight m , and that $S^{(p^p)}$ has a Sylow p -subgroup of \mathfrak{S}_{p^2} as a vertex.

There has been considerable work on the computation of the vertices of simple Specht modules. In [27], the third author showed that the vertices of the simple Specht modules $S^{(n-r, 1^r)}$ when n is not divisible by p are the Sylow p -subgroups of $\mathfrak{S}_{n-r-1} \times \mathfrak{S}_r$. In [16] Hemmer proved that, when $p \geq 3$, a simple Specht module is a signed Young module, as defined by Donkin [10, 2.3]. By [10, 5.1(3)], it follows that the Young vertices of simple Specht modules in odd characteristic are certain Young subgroups of symmetric groups and hence the vertices, in the sense of Green, are the Sylow p -subgroups of the corresponding Young subgroups (see [8, Theorem 3.7(b)]). In [8], Danz and the second author identified the labelings of the signed Young modules isomorphic to simple Specht modules when $p \geq 5$. As consequences, the vertices, Green correspondents, cohomological varieties and complexities of

all simple Specht modules are known except when $p = 3$ (see [8, Corollary 6.5, Remark 6.8]).

Finally we note that there has also been considerable work on the vertices of simple modules for symmetric groups. We refer the reader to [9] for a survey and to [7] for some recent advances.

Outline. In Section 2 we prove Theorem 1.1 for symmetric groups. Then in Section 3 we use this result and further arguments to prove Theorem 1.1 for alternating groups. In Section 4 we give a short proof, using Theorem 1.1, that Theorem 1.3 implies Theorem 1.2.

We then turn to the proof of Theorem 1.3. In Section 5 we collect the necessary background results on hook-Specht modules. We review the Brauer correspondence in Section 6 and give a useful general result on the Brauer homomorphism for monomial modules. The notion of generic Jordan type is presented in Section 7. Then in Sections 8 and 9 we prove Theorem 1.3, under the hypotheses (i) and (ii) of Theorem 1.3, respectively.

We end in Section 10 by stating some open problems related to our three main theorems.

2. PROOF OF THEOREM 1.1 FOR SYMMETRIC GROUPS

For background on Sylow subgroups of symmetric groups and their construction as product of iterated wreath products, we refer the reader to [19, (4.1.20)].

We need the following notation. Let \mathfrak{S}_A denote the symmetric group on a finite set A . For each $A \subset \mathbb{N}$ such that A has size p^d where $d \in \mathbb{N}$, let E_A denote a fixed elementary abelian subgroup of \mathfrak{S}_A of order p^d , acting regularly on A . All such subgroups are permutation isomorphic to the regular representation of an elementary abelian group of order p^d , and so are conjugate in \mathfrak{S}_A . We highlight that the precise choice of E_A is irrelevant.

The main work in the proof of Theorem 1.1 comes in the case when $n = mp$ for some $m \in \mathbb{N}$. Suppose that

$$(1) \quad n = m_1p + m_2p^2 + \cdots + m_r p^r$$

where $m_i \in \mathbb{N}_0$ for each i and $m_r \neq 0$. Note that this is not necessarily the p -adic expansion of n . For $i \in \{1, \dots, r\}$ and each j such that $j \in \{1, \dots, m_i\}$, let

$$z_j^i = (j-1)p^i + \sum_{s=1}^{i-1} m_s p^s$$

and let $B_j^i = \{z_j^i + 1, \dots, z_j^i + p^i\}$. Note that $|B_j^i| = p^i$ for all j such that $j \in \{1, \dots, m_i\}$ and that the B_j^i partition $\{1, \dots, mp\}$. If $m_i = 0$ then $B_j^i = \emptyset$ and we define $E_{B_j^i}$ to be the trivial group. For each expression for n

as in (1), define an elementary abelian p -subgroup of \mathfrak{S}_n by

$$E(m_1, \dots, m_r) = \prod_{i=1}^r \prod_{j=1}^{m_i} E_{B_j^i}.$$

Proposition 2.1 ([2, 1.3 Chapter VI]). *The p -subgroups $E(m_1, \dots, m_r)$ where m_1, \dots, m_r satisfy (1) form a complete irredundant set of maximal elementary abelian p -subgroups of \mathfrak{S}_{mp} up to \mathfrak{S}_{mp} -conjugacy.*

In fact each $E(m_1, \dots, m_r)$ is maximal even as an abelian subgroup of \mathfrak{S}_{mp} ; we leave this remark as an exercise to the reader.

The subgroup $E(m)$ generated by m disjoint p -cycles is critical to our argument. To emphasise the role played by this subgroup, we denote it by $C(m)$ below. Informally stated, the next lemma says that $C(m)$ is the normal base group in any Sylow p -subgroup that contains it.

Lemma 2.2. *If Q is a p -subgroup of \mathfrak{S}_{mp} containing $C(m)$ then $C(m) \trianglelefteq Q$. Moreover $Q/C(m)$ acts faithfully on the orbits of $C(m)$.*

Proof. Let P be a Sylow p -subgroup of \mathfrak{S}_{mp} containing Q . By the construction of Sylow p -subgroups in [19, (4.1.20)], P has a normal base group B generated by m disjoint p -cycles τ_1, \dots, τ_m . If $\sigma \in P$ then σ permutes the orbits of the τ_k as blocks for its action. It follows that if $\sigma \in P$ is a p -cycle then $\sigma \in \langle \tau_k \rangle$ for some k . Hence $C(m) = B$ and $C(m) \trianglelefteq P$. Moreover, if $\sigma \in P$ fixes each orbit of $C(m)$ setwise then σ is a product of disjoint p -cycles each contained in $C(m)$. Therefore $P/C(m)$ acts faithfully on the orbits of $C(m)$. \square

In particular, Lemma 2.2 determines how an elementary abelian p -subgroup that normalizes $C(m)$ can act on the orbits of $C(m)$.

Lemma 2.3. *Let $d \in \mathbb{N}$ and let $\sigma \in \mathfrak{S}_{p^d}$. Let $C(p^{d-1})$ be as defined when $n = p^d$, so $C(p^{d-1})$ is generated by p^{d-1} disjoint p -cycles, and let X be the set of orbits of $C(p^{d-1})$. Let $A = \{1, 2, \dots, p^d\}$. Suppose that a p -subgroup P of \mathfrak{S}_{p^d} contains both ${}^\sigma E_A$ and $C(p^{d-1})$. Then ${}^\sigma E_A C(p^{d-1})/C(p^{d-1})$ acts regularly on X .*

Proof. By Lemma 2.2 in the case $n = p^d$, the subgroup $C(p^{d-1})$ is normal in P and ${}^\sigma E_A C(p^{d-1})/C(p^{d-1})$ acts faithfully on X . Since E_A acts transitively on $\{1, \dots, p^d\}$, the action of ${}^\sigma E_A C(p^{d-1})/C(p^{d-1})$ on X is transitive. Since E_A is abelian, it follows that ${}^\sigma E_A C(p^{d-1})/C(p^{d-1}) \cong {}^\sigma E_A / (C(p^{d-1}) \cap {}^\sigma E_A)$ is also abelian and hence it acts regularly on X . \square

We are now ready to prove Theorem 1.1.

Proof of Theorem 1.1 when $G = \mathfrak{S}_n$. We prove the theorem by induction on n . Let $Q \leq \mathfrak{S}_n$ be a p -subgroup containing an \mathfrak{S}_n -conjugate of every elementary abelian p -subgroup of \mathfrak{S}_n . Let $n = mp + c$ where $m \in \mathbb{N}$ and

$0 \leq c < p$. If $c \geq 1$ then by conjugating Q by an element of \mathfrak{S}_n we may assume that $Q \leq \mathfrak{S}_{mp}$. By induction we get that Q is a Sylow p -subgroup of \mathfrak{S}_{mp} , and hence a Sylow p -subgroup of \mathfrak{S}_n . We may therefore assume that $n = mp$, and, by induction, that the theorem holds true for \mathfrak{S}_m .

By replacing Q with a \mathfrak{S}_{mp} -conjugate, we may assume that $C(m)$ is contained in Q . Let $X = \{B_1^1, \dots, B_m^1\}$ be the set of orbits of $C(m)$. By Lemma 2.2 we have that $C(m) \trianglelefteq Q$ and $Q/C(m)$ acts faithfully on X .

Let $\overline{H} \leq \mathfrak{S}_X$ denote the image of a subgroup H of Q under the quotient map $Q \rightarrow Q/C(m)$, regarding $Q/C(m)$ as a subgroup of \mathfrak{S}_X . We aim to prove that \overline{Q} contains a \mathfrak{S}_X -conjugate of every elementary abelian p -subgroup of \mathfrak{S}_X .

Let $m = \ell_0 + \sum_{i=1}^t \ell_i p^i$ such that $\ell_t \neq 0$. By assumption there exists $\sigma \in \mathfrak{S}_{mp}$ such that Q contains ${}^\sigma E(\ell_0, \dots, \ell_t)$. (Note that because of the index shift, this subgroup has exactly ℓ_{i-1} orbits of size p^i on $\{1, \dots, mp\}$.) We have

$${}^\sigma E(\ell_0, \dots, \ell_t) = \prod_{i=1}^{t+1} \prod_{j=1}^{\ell_{i-1}} E_{\sigma B_j^i}.$$

Fix $i \in \{1, \dots, t+1\}$ and $j \in \{1, \dots, \ell_{i-1}\}$. Let Ω_j^i be the set of all $k \in \{1, \dots, m\}$ such that the orbit B_k^1 of $C(m)$ is contained in the support of $E_{\sigma B_j^i}$, i.e. $B_k^1 \subseteq \sigma B_j^i$. Let $X_j^i = \{B_k^1 : k \in \Omega_j^i\}$. Thus X is the disjoint union of the X_j^i , the support of $E_{\sigma B_j^i}$ is $\bigcup_{k \in \Omega_j^i} B_k^1$ and $\overline{E_{\sigma B_j^i}}$ acts on X_j^i . By Lemma 2.3 this action is regular. It follows that the elementary abelian p -subgroup

$$\overline{{}^\sigma E(\ell_0, \dots, \ell_t)} \leq \mathfrak{S}_X$$

has exactly ℓ_i orbits on X of size p^i for each $i \in \{0, \dots, t\}$. Since the ℓ_i were freely chosen, we see that \overline{Q} contains a \mathfrak{S}_X -conjugate of every elementary abelian p -subgroup of \mathfrak{S}_X . It now follows by induction that \overline{Q} acts on X as a Sylow p -subgroup of \mathfrak{S}_X .

If p^a is the order of a Sylow p -subgroup of \mathfrak{S}_m then

$$|Q| = |C(m)| |\overline{Q}| = p^{m+a},$$

which is the order of a Sylow p -subgroup of \mathfrak{S}_{mp} . The theorem follows. \square

3. PROOF OF THEOREM 1.1 FOR ALTERNATING GROUPS

Case $p \geq 3$. In this case any p -subgroup of \mathfrak{S}_n is a p -subgroup of \mathfrak{A}_n . Hence if $Q \leq \mathfrak{A}_n$ contains an \mathfrak{A}_n -conjugate of every elementary abelian p -subgroup of \mathfrak{A}_n then Q contains an \mathfrak{S}_n -conjugate of every elementary abelian p -subgroup of \mathfrak{S}_n , and so by Theorem 1.1 for \mathfrak{S}_n , we get that Q is a Sylow p -subgroup of \mathfrak{S}_n .

Case $p = 2$. Let $n = 2m$. We need the classification of all maximal elementary abelian 2-subgroups of \mathfrak{A}_n up to \mathfrak{S}_n -conjugacy. Suppose that

$$(2) \quad n = 2m_1 + 2^2m_2 + \cdots + 2^r m_r$$

where $m_i \in \mathbb{N}_0$ for each i , $m_r \neq 0$ and $m_1 \neq 2$. For each such expression define

$$F(m_1, m_2, \dots, m_r) = E(m_1, m_2, \dots, m_r) \cap \mathfrak{A}_n.$$

The reason for excluding the case $m_1 = 2$ is that $E(2, m_2, \dots, m_r) \cap \mathfrak{A}_n$ is properly contained in $F(0, m_2 + 1, m_3, \dots, m_r)$. Note also that the subgroup $F(m_1, m_2, \dots, m_r)$ has exactly two fixed points on $\{1, \dots, n\}$ if $m_1 = 1$.

Proposition 3.1 ([1, Proposition 5.2]). *The subgroups $F(m_1, \dots, m_r)$ where m_1, \dots, m_r satisfy (2) form a complete irredundant set of maximal elementary abelian 2-subgroups of \mathfrak{A}_{2m} up to \mathfrak{S}_{2m} -conjugacy.*

In [1] the authors remark that the proposition follows from the analogous result for \mathfrak{S}_n . We take this opportunity to provide a complete proof. The following lemma is required.

Lemma 3.2. *Suppose that m_1, \dots, m_r are as in (2) and m'_1, \dots, m'_s are as in (1). If $F(m_1, \dots, m_r) \leq {}^\sigma E(m'_1, \dots, m'_s)$ for some $\sigma \in \mathfrak{S}_{2m}$ then $r = s$ and $m_i = m'_i$ for all $i \in \{1, \dots, r\}$.*

Proof. If $m_1 = 0$ then $F(0, m_2, \dots, m_r) = E(0, m_2, \dots, m_r)$ is maximal and by Proposition 2.1 we have $m_i = m'_i$ for all $i \in \{2, \dots, r\}$. Hence $m'_1 = 0$ and the lemma holds in this case. If $m_1 = 1$ then $F(1, m_2, \dots, m_r)$ is equal to the subgroup $E(0, m_2, \dots, m_r)$. Since $E(0, m_2, \dots, m_r)$ has exactly two fixed points and no orbit of size 2 on $\{1, \dots, 2m\}$, we have $m'_1 = 1$. It follows that σ fixes the set $\{1, 2\}$ and hence $E(1, m_2, \dots, m_r) \leq {}^\sigma E(m'_1, \dots, m'_s)$. We may again apply Proposition 2.1.

Suppose that $m_1 \geq 3$. Let $\tau_k = (2k - 1, 2k)(2k + 1, 2k + 2)$ for $k \in \{1, \dots, m_1 - 1\}$. Let B_j^2 for $j \in \{1, \dots, m'_2\}$ be as defined after (1) for the subgroup $E(m'_1, \dots, m'_s)$. If there exist j and k such that $\tau_k \in {}^\sigma E_{B_j^2}$ then $\sigma(B_j^2) = \{2k - 1, 2k, 2k + 1, 2k + 2\}$ and an element of ${}^\sigma E(m'_1, \dots, m'_s)$ moves $2k - 1$ if and only if it moves $2k + 1$. But if $k > 1$ then $\tau_{k-1} \in F(m_1, \dots, m_r)$ and otherwise $k = 1$ and $\tau_2 \in F(m_1, \dots, m_r)$, so this is a contradiction. Hence all the transpositions $(2k - 1, 2k)$ for $k \in \{1, \dots, m_1\}$ lie in $E(m'_1, \dots, m'_s)$ and so $m'_1 \geq m_1$. On the other hand, $F(m_1, \dots, m_r)$ has m_1 orbits of size 2 on $\{1, \dots, 2m\}$ and ${}^\sigma E(m'_1, \dots, m'_s)$ has m'_1 orbits of size 2 on $\{1, \dots, 2m\}$; since $F(m_1, \dots, m_r) \leq {}^\sigma E(m'_1, \dots, m'_s)$ we have $m_1 \geq m'_1$. Therefore $m_1 = m'_1$. Hence there exist $\tau, \tau' \in \mathfrak{S}_{2m}$ such that

$$\begin{aligned} F(m_1, m_2, \dots, m_r) &= F(m_1) \times {}^\tau F(0, m_2, \dots, m_r) \\ E(m'_1, m'_2, \dots, m'_s) &= {}^\sigma E(m_1) \times {}^{\tau'} E(0, m'_2, \dots, m'_s) \end{aligned}$$

and ${}^\tau F(0, m_2, \dots, m_r) \leq {}^\tau E(0, m'_2, \dots, m'_s)$. Both groups have support of size $2m - 2m_1$. Hence, by the special case $m_1 = 0$ we get $r = s$ and $m_i = m'_i$ for all $i \in \{2, \dots, r\}$. \square

Proof of Proposition 3.1. Let F be an elementary abelian p -subgroup of \mathfrak{A}_{2m} . By Proposition 2.1 there exist m'_1, \dots, m'_s and $\sigma \in \mathfrak{S}_{2m}$ such that

$$F \leq {}^\sigma E(m'_1, \dots, m'_s) \cap \mathfrak{A}_{2m} = {}^\sigma (E(m'_1, \dots, m'_s) \cap \mathfrak{A}_{2m}).$$

If $m'_1 \neq 2$ then $E(m'_1, \dots, m'_s) \cap \mathfrak{A}_{2m} = F(m'_1, \dots, m'_s)$. In the remaining case we have $E(2, m'_2, \dots, m'_s) \cap \mathfrak{A}_{2m} \leq F(0, m'_2 + 1, \dots, m'_s)$. Suppose further that $F = F(m_1, \dots, m_r)$. By Lemma 3.2, we have $r = s$ and $m_i = m'_i$ for all $i \in \{1, \dots, r\}$. Therefore the subgroups $F(m_1, \dots, m_r)$ for m_1, \dots, m_r satisfying (2) form a complete set of maximal elementary abelian 2-subgroups of \mathfrak{A}_{2m} up to \mathfrak{S}_{2m} -conjugacy. It is clear from the orbits of these subgroups on $\{1, \dots, 2m\}$ that no two of them are conjugate. \square

We are now ready to prove Theorem 1.1 in the alternating group case.

Proof of Theorem 1.1 when $G = \mathfrak{A}_n$ and $p = 2$. As in the proof of Theorem 1.1 for \mathfrak{S}_n we reduce to the case when $n = 2m$. The cases $m = 1$ and $m = 2$ are easily checked, so we may assume that $m \geq 3$.

Let $Q \leq \mathfrak{A}_{2m}$ be a 2-subgroup containing an \mathfrak{A}_{2m} -conjugate of every elementary abelian 2-subgroup of \mathfrak{A}_{2m} . Without loss of generality we may assume that $F(m)$ is contained in Q . Let P be a Sylow 2-subgroup of \mathfrak{S}_{2m} containing Q . As in the proof of Lemma 3.2, let $\tau_k = (2k - 1, 2k)(2k + 1, 2k + 2) \in Q$ for $k \in \{1, \dots, m - 1\}$. Let (x, y) be a transposition in P with $x < y$. If $y \neq x + 1$ then there exists k such that exactly one of x, y lies in $\{2k - 1, 2k, 2k + 1, 2k + 2\}$. It is easily checked that in this case $[(x, y), \tau_k]$ is a 3-cycle. So $y = x + 1$. Moreover, if $x = 2k$ and $y = 2k + 1$ for some k then either $k > 1$ and $(2k, 2k + 1)\tau_{k-1}$ contains a 3-cycle, or $k = 1$ and $(2, 3)\tau_2$ contains a 3-cycle. It follows that the transpositions in P are of the form $(2k - 1, 2k)$ for $k \in \{1, \dots, m\}$. Therefore $F(m) \leq C(m) \leq P$.

By Lemma 2.2, $C(m) \trianglelefteq P$. In particular, $Q \leq N_{\mathfrak{S}_{2m}}(C(m))$. Since $QC(m)/C(m) \cong C(m)/(C(m) \cap Q) = C(m)/F(m)$, we see that $QC(m) = \langle Q, (1, 2) \rangle$ is a 2-group having Q as a subgroup of index 2.

Let $E = E(m_1, m_2, \dots, m_r)$ be a maximal elementary abelian 2-subgroup of \mathfrak{S}_{2m} . Let $F = E(m_1, m_2, \dots, m_r) \cap \mathfrak{A}_{2m}$. By hypothesis there exists $\sigma \in \mathfrak{S}_{2m}$ such that ${}^\sigma F \leq Q$. If $m_1 = 0$ then ${}^\sigma E = {}^\sigma F \leq Q$. Suppose that $m_1 \geq 1$. Let $\Omega = \{\sigma(2m_1 + 1), \dots, \sigma(2m)\}$ and let $2i - 1 \leq \sigma(1) \leq 2i$. Since $\sigma(1) \notin \Omega$, similar arguments to those used earlier in the proof show that $\{2i - 1, 2i\} \cap \Omega = \emptyset$. Again similar arguments show that either

- (a) $(\sigma(1), \sigma(2)) = (2i - 1, 2i)$, or
- (b) $m_1 = 2$ and $(2i - 1, 2i)$ is either $(\sigma(1), \sigma(3))$ or $(\sigma(1), \sigma(4))$.

So

$$\langle (2i-1, 2i), {}^\sigma F \rangle = \begin{cases} {}^\sigma E & \text{in case (a),} \\ {}^\sigma(R \times E(0, m_2, \dots, m_r)) & \text{in case (b),} \end{cases}$$

for some Sylow 2-subgroup R of \mathfrak{S}_4 . In either case, $\langle (2i-1, 2i), {}^\sigma F \rangle$ contains a \mathfrak{S}_{2m} -conjugate of E . Clearly, $\langle (2i-1, 2i), {}^\sigma F \rangle \leq QC(m)$. It follows that $QC(m)$ contains all maximal elementary abelian 2-subgroups of \mathfrak{S}_{2m} up to \mathfrak{S}_{2m} -conjugacy. By Theorem 1.1 for \mathfrak{S}_n , we see that $QC(m)$ is a Sylow 2-subgroup of \mathfrak{S}_{2m} . Hence $QC(m) \cap \mathfrak{A}_{2m} = Q$ is a Sylow 2-subgroup of \mathfrak{A}_{2m} . \square

We note that this proof only used the weaker hypothesis that Q contains an \mathfrak{S}_{2m} -conjugate of every elementary abelian 2-subgroup of \mathfrak{A}_{2m} . As remarked in [1], the 2-subgroups $E(0, 0, m_3, \dots, m_r)$ are not normalized by any odd elements in \mathfrak{A}_{2m} , and so there are two \mathfrak{A}_{2m} -conjugacy classes of such subgroups; in all other cases the \mathfrak{A}_{2m} - and the \mathfrak{S}_{2m} -classes agree. This may be proved as follows:¹ if $A = \{1, \dots, 2^d\}$ then $N_{\mathfrak{S}_{2^d}}(E_A)$ is permutation isomorphic to the affine general linear group $\text{AGL}_d(\mathbb{F}_2)$; this group is generated by translations and transvections, both of which are even permutations of \mathbb{F}_2^d when $d \geq 3$. The normalizer of $E(0, 0, m_3, \dots, m_r)$ factors as a direct product of wreath products $N_{\mathfrak{S}_{2^d}}(E_{A_i}) \wr \mathfrak{S}_{m_i}$ where A_i has size 2^i . The permutations in a copy of the top group \mathfrak{S}_{m_i} act on blocks of size 2^i and so are even. Hence $N_{\mathfrak{S}_{2^d}}(E(0, 0, m_3, \dots, m_r)) \leq \mathfrak{A}_{2m}$. It is easily seen that $\langle (12)(34) \rangle$ and $\langle (12)(34), (13)(24) \rangle$ are normalized by odd permutations in \mathfrak{S}_4 , so these are the only exceptional cases.

4. PROOF OF THEOREM 1.2 FROM THEOREM 1.3

In this section, we deduce Theorem 1.2 from Theorem 1.3 using Theorem 1.1.

Let \overline{F} be the algebraic closure of F . By [17, Theorem 1.21], for any FG -module M and $H \leq G$, we have that $\overline{F} \otimes_F M$ is a direct summand of $\overline{F} \otimes_F (M \downarrow_H \uparrow^G) \cong (\overline{F} \otimes_F M) \downarrow_H \uparrow^G$ if and only if M is a direct summand of $M \downarrow_H \uparrow^G$. Thus the vertices of an indecomposable FG -module do not change under field extensions of F .

Without loss of generality, we may assume that F is an algebraically closed field. Let Q be a vertex of the Specht module $S^{(kp-p, 1^p)}$ where $k \equiv 1 \pmod{p}$ and $k \not\equiv 1 \pmod{p^2}$. By Theorem 1.1, it is sufficient to prove that if

$$kp = m_1p + m_2p^2 + \dots + m_r p^r$$

where $m_i \in \mathbb{N}_0$ for each $i \in \{1, \dots, r\}$ and $m_r \neq 0$ then Q contains a conjugate of $E(m_1, m_2, \dots, m_r)$. If $m_2 \neq 0$ then Theorem 1.3(ii) applies. If

¹The proof in [1] is indicated very briefly, but seems to incorrectly assume that if a subgroup of \mathfrak{A}_n is normalized by an odd element then it is normalized by a transposition.

$m_2 = 0$ then $k = m_1 + m_3p^2 + \cdots + m_r p^{r-1}$ and so $k \equiv m_1 \pmod{p^2}$. Hence $m_1 \geq 2$ and Theorem 1.3(i) applies. This completes the proof.

5. HOOK-SPECHT MODULES

For maximal generality we work in this section with modules over an arbitrary commutative ring K . The definition of Specht modules given in [18, Definition 4.3] extends easily to this setting. Fix $n \in \mathbb{N}$. Let $N = \langle e_1, \dots, e_n \rangle$ be the K -free natural permutation module for $K\mathfrak{S}_n$ of rank n . Then, by definition, $S^{(n-1,1)}$ is the K -free submodule of N with K -basis $\{e_2 - e_1, \dots, e_n - e_1\}$.

We begin by establishing the isomorphism of $K\mathfrak{S}_n$ -modules $S^{(n-r,1^r)} \cong \bigwedge^r S^{(n-1,1)}$. This isomorphism can be found, most obviously in the case when K is a finite field of characteristic 2, in Peel's papers [24, Section 6] and [25]. When K is a field of prime characteristic p and $n = p$ it was proved by Hamernik [15]; it is easily seen that Hamernik's proof also works in the general case. When K is a field it was proved by Müller and Zimmermann in [22, Proposition 2.3(a)]. The proof given here combines ideas from both [15] and [22].

Proposition 5.1. *Let $1 \leq r \leq n - 1$.*

(i) *The set*

$$\{(e_{i_1} - e_1) \wedge \cdots \wedge (e_{i_r} - e_1) : 1 < i_1 < \cdots < i_r \leq n\}$$

is a K -basis of $\bigwedge^r S^{(n-1,1)}$.

(ii) *The map sending $(e_{i_1} - e_1) \wedge \cdots \wedge (e_{i_r} - e_1) \in \bigwedge^r S^{(n-1,1)}$ to $e_{\mathfrak{t}} \in S^{(n-r,1^r)}$, where $1 < i_1 < \cdots < i_r \leq n$ and \mathfrak{t} is the unique standard tableau of shape $(n-r, 1^r)$ having i_1, \dots, i_r in its rows of length one, is an isomorphism of $K\mathfrak{S}_n$ -modules.*

Proof. Part (i) is obvious from the basis of $S^{(n-1,1)}$ above. By (i) and the Standard Basis Theorem for Specht modules (see [18, Theorem 8.4]), the map defined in (ii) is a K -linear isomorphism. So all we have to check is that it commutes with the action of \mathfrak{S}_n . For the subgroup of \mathfrak{S}_n fixing 1 this is obvious. So it suffices to check the action of the permutation (12). Let $w = (e_{i_1} - e_1) \wedge \cdots \wedge (e_{i_r} - e_1)$ where $1 < i_1 < \cdots < i_r \leq n$. If $i_1 = 2$ then it is clear that $(12)e_{\mathfrak{t}} = -e_{\mathfrak{t}}$ and $(12)w = -w$. Suppose that $i_1 \neq 2$. For each $1 \leq a \leq r$, let \mathfrak{t}_a be the standard tableau having $2, i_1, \dots, \widehat{i_a}, \dots, i_r$ in its rows of length 1, where the hat over i_a indicates this entry is omitted. By the Garnir relation (see [18, Theorem 7.2]) involving all entries in the first column of \mathfrak{t} , and the single box in the second column of \mathfrak{t} , we have

$$(12)e_{\mathfrak{t}} = e_{\mathfrak{t}} - e_{\mathfrak{t}_1} + e_{\mathfrak{t}_2} - \cdots + (-1)^r e_{\mathfrak{t}_r}.$$

On the other hand, the action on the wedge product is given by

$$(12)w = ((e_{i_1} - e_1) - (e_2 - e_1)) \wedge \cdots \wedge ((e_{i_r} - e_1) - (e_2 - e_1)) \\ = (e_{i_1} - e_1) \wedge \cdots \wedge (e_{i_r} - e_1) \\ + \sum_{a=1}^r (-1)^a (e_2 - e_1) \wedge (e_{i_1} - e_1) \wedge \cdots \wedge (\widehat{e_{i_a} - e_1}) \wedge \cdots \wedge (e_{i_r} - e_1).$$

The proof is now complete. \square

We now introduce some further ideas from [15] and basic simplicial homology. For $r \in \mathbb{N}$ define $\delta_r : \bigwedge^r N \rightarrow \bigwedge^{r-1} N$ by

$$\delta_r(e_{i_1} \wedge \cdots \wedge e_{i_r}) = \sum_{a=1}^r (-1)^{a-1} e_{i_1} \wedge \cdots \wedge \widehat{e_{i_a}} \wedge \cdots \wedge e_{i_r}.$$

The subscript r in the map δ_r will be omitted when it is clear from the context. By definition $\bigwedge^0 N = K$. We leave it to the reader to verify the relation

$$(3) \quad (e_{i_1} - e_j) \wedge \cdots \wedge (e_{i_r} - e_j) = \delta(e_j \wedge e_{i_1} \wedge \cdots \wedge e_{i_r})$$

for $1 \leq j, i_1, \dots, i_r \leq n$.

Proposition 5.2 (Long exact sequence). *The sequence*

$$0 \longrightarrow \bigwedge^n N \xrightarrow{\delta_n} \cdots \xrightarrow{\delta_{r+1}} \bigwedge^r N \xrightarrow{\delta_r} \bigwedge^{r-1} N \xrightarrow{\delta_{r-1}} \cdots \rightarrow N \xrightarrow{\delta_1} K \longrightarrow 0$$

is exact. Moreover if $1 \leq r < n$ then $\ker \delta_r = \text{im } \delta_{r+1} = \bigwedge^r S^{(n-1, 1)}$.

Proof. The K -linear map defined by $e_{i_1} \wedge \cdots \wedge e_{i_r} \mapsto e_1 \wedge e_{i_1} \wedge \cdots \wedge e_{i_r}$ defines a homotopy equivalence between the identity map on the sequence above and the zero map. This shows that the sequence is null-homotopic and hence exact. Proposition 5.1(i) and (3) imply that $\text{im } \delta_{r+1}$ is equal to $\bigwedge^r S^{(n-1, 1)}$. \square

We remark that there is an important homological interpretation of Proposition 5.2. Let e_1, \dots, e_n be the canonical basis of \mathbb{R}^n and fix an $(n-1)$ -simplex in \mathbb{R}^n with geometric vertices e_1, \dots, e_n . The wedge product $e_{i_1} \wedge \cdots \wedge e_{i_r}$ can then be identified with the oriented $(r-1)$ -simplex with vertices, in order, e_{i_1}, \dots, e_{i_r} . For each $r \geq 2$, the map $\delta_r : \bigwedge^r N \rightarrow \bigwedge^{r-1} N$ is the boundary map from simplicial homology, acting on oriented $(r-1)$ -simplices. Replace the map $\delta_1 : N \rightarrow K$ with the zero map $N \rightarrow 0$. Then Proposition 5.2 is equivalent to the fundamental result that the solid $(n-1)$ -simplex has trivial homology (with coefficients in K) in all non-zero degrees, and its zero homology group is $N / \ker \delta_1 \cong K$.

We end this section with some further notation and results that are used in Section 9. Let $r \in \mathbb{N}$. Let

$$I^{(r)} = \{(i_1, \dots, i_r) : 1 \leq i_1 < \cdots < i_r \leq n, i_1, \dots, i_r \in \mathbb{N}\}.$$

We call the elements of $I^{(r)}$ *multi-indices*. Let

$$J^{(r)} = \{\mathbf{i} \in I^{(r)} : i_1 > 1\}.$$

For $\mathbf{i} \in I^{(r)}$, let $e_{\mathbf{i}} = e_{i_1} \wedge \cdots \wedge e_{i_r}$. We say that $\{e_{\mathbf{i}} : \mathbf{i} \in I^{(r)}\}$ is the *monomial basis* of $\bigwedge^r N$. By (3), the set

$$(4) \quad \{\delta(e_1 \wedge e_{\mathbf{j}}) : \mathbf{j} \in J^{(r)}\}$$

is a K -basis for $\bigwedge^r S^{(n-1,1)}$, corresponding under the isomorphism in Proposition 5.1 to the standard basis of $S^{(n-r,1^r)}$. The following lemma gives a very useful way to express elements of $\bigwedge^r S^{(n-1,1)}$ in this basis.

Lemma 5.3 (Rewriting Lemma). *Let $u = \sum_{\mathbf{i} \in I^{(r)}} \mu_{\mathbf{i}} e_{\mathbf{i}} \in \bigwedge^r S^{(n-1,1)}$. Then*

$$u = \sum_{\mathbf{j} \in J^{(r)}} \mu_{\mathbf{j}} \delta(e_1 \wedge e_{\mathbf{j}}).$$

Proof. By (4) we may write

$$u = \sum_{\mathbf{j} \in J^{(r)}} \nu_{\mathbf{j}} \delta(e_1 \wedge e_{\mathbf{j}})$$

for some $\nu_{\mathbf{j}} \in K$. Each of the monomial summands in $\delta(e_1 \wedge e_{\mathbf{j}})$ involves e_1 , with the sole exception of the first summand $e_{\mathbf{j}}$. Therefore the coefficient of the monomial $e_{\mathbf{j}}$ in the right-hand side of the above equation is $\nu_{\mathbf{j}}$. Hence $\nu_{\mathbf{j}} = \mu_{\mathbf{j}}$, as required. \square

Finally we note that if $u \in \bigwedge^r N$ and $v \in \bigwedge^s N$, where $r, s \in \mathbb{N}$, then

$$(5) \quad \delta(u \wedge v) = \delta(u) \wedge v + (-1)^r u \wedge \delta(v).$$

6. THE BRAUER HOMOMORPHISM AND MONOMIAL MODULES

Throughout this section let F be a field of prime characteristic p and let G be a finite group.

6.1. The Brauer homomorphism. Let V be an FG -module. For $Q \leq G$, define

$$V^Q = \{v \in V : \sigma v = v \text{ for all } \sigma \in Q\}.$$

For $R \leq Q \leq G$, the *relative trace* map $\text{Tr}_R^Q : V^R \rightarrow V^Q$ is the linear map defined by

$$\text{Tr}_R^Q(v) = \sum_{\sigma} \sigma v$$

where the sum is over a set of coset representatives for Q/R . The *Brauer kernel* of V with respect to Q is the subspace $\sum_{R < Q} \text{Tr}_R^Q V^R$ of V^Q . The *Brauer quotient* of V with respect to Q is

$$V(Q) = V^Q / \sum_{R < Q} \text{Tr}_R^Q V^R.$$

It is easy to show that V^Q and $\sum_{R < Q} \text{Tr}_R^Q V^R$ are both $N_G(Q)$ -invariant, and so $V(Q)$ is a module for $F(N_G(Q)/Q)$.

The following proposition is proved in [5, (1.3)].

Proposition 6.1. *Let V be an indecomposable FG -module. Let Q be a p -subgroup of G . If $V(Q) \neq 0$ then V has a vertex containing Q .*

The converse to Proposition 6.1 is true when V has the trivial module as its source, or, equivalently, if V is a direct summand of a permutation module. In general it is false. For example, calculations using MAGMA show that if $p = 2$ and P is a Sylow 2-subgroup of \mathfrak{S}_6 then $S^{(4,1,1)}(P) = 0$. But by [23, Theorems 4.3, 4.5], $S^{(4,1,1)}$ is indecomposable with vertex P .

6.2. Monomial modules. Let H be a subgroup of G and let $\theta : H \rightarrow F^\times$ be a representation of H such that $\theta(h) = 1$ whenever $h \in H$ is a p -element. Let $\langle v \rangle$ be the corresponding FH -module, i.e. $hv = \theta(h)v$ for all $h \in H$. Let $V = FG \otimes_{FH} \langle v \rangle$ be the monomial module induced from $\langle v \rangle$. If P is a Sylow p -subgroup of H then $\langle v \rangle$ has P as a vertex, and so $\langle v \rangle$ is a direct summand of $\langle v \rangle \downarrow_P \uparrow^H = F \uparrow_P^H$. Hence V is a p -permutation module. The following proposition establishes a close connection between V and the permutation module $F \uparrow_H^G$.

Proposition 6.2. *Let $V = \langle v \rangle \uparrow_H^G$ be the FG -module defined above and let $W = \langle e_{\gamma H} : \gamma \in G \rangle$ be the permutation module of G acting on the cosets of H . Let Q be a p -subgroup of G . Let $\gamma_1 H, \dots, \gamma_L H$ be representatives for the orbits of Q on G/H . For each $\ell \in \{1, \dots, L\}$ denote by s_ℓ the element defined by*

$$s_\ell = \sum_{\tau \in T} \tau \gamma_\ell \otimes v,$$

where T is a set of representatives for the cosets of $\text{Stab}_Q(\gamma_\ell H)$ in Q . Then

- (i) the map defined by $\alpha \gamma_\ell \otimes v \mapsto \alpha e_{\gamma_\ell H}$ for $\alpha \in Q$ and $\ell \in \{1, \dots, L\}$ is an isomorphism $V \downarrow_Q \cong W \downarrow_Q$,
- (ii) $s_\ell = \text{Tr}_{\text{Stab}_Q(\gamma_\ell H)}^Q(\gamma_\ell \otimes v)$,
- (iii) $\{s_\ell : 1 \leq \ell \leq L\}$ is a basis for V^Q ,
- (iv) if $R \leq Q$ then $\{s_\ell : 1 \leq \ell \leq L, \text{Stab}_Q(\gamma_\ell H) = \text{Stab}_R(\gamma_\ell H)\}$ is a basis for $\text{Tr}_R^Q V^R$.

Proof. Let $g \in G$ and let R be a p -subgroup of ${}^g H$. By definition we have that ${}^g \langle v \rangle \downarrow_R$ is isomorphic to F_R , the trivial FR -module. Hence, part (i) follows from Mackey's induction/restriction formula since

$$V \downarrow_Q = \bigoplus_{g \in Q \backslash G/H} ({}^g \langle v \rangle \downarrow_{{}^g H \cap Q}) \uparrow^Q \cong \bigoplus_{g \in Q \backslash G/H} F \uparrow_{{}^g H \cap Q}^Q \cong (F \uparrow_H^G) \downarrow_Q \cong W \downarrow_Q.$$

The orbit sums of Q acting on the canonical permutation basis of W are a basis for W^Q . Since the orbit sum containing $e_{\gamma_\ell H}$ is $\text{Tr}_{\text{Stab}_Q(\gamma_\ell H)}^Q e_{\gamma_\ell H}$, parts (ii) and (iii) follow from (i).

In order to prove (iv) we observe that, by (iii), there exist a basis of V^R consisting of elements of the form $w_\gamma = \text{Tr}_{\text{Stab}_R(\gamma H)}^R(\gamma \otimes v)$, for $\gamma \in G$. Let $\gamma_\ell H$ and γH be in the same Q -orbit for some $1 \leq \ell \leq L$. Then $\text{Stab}_R(\gamma H) = \text{Stab}_Q(\gamma H)$ is equivalent to $\text{Stab}_R(\gamma_\ell H) = \text{Stab}_Q(\gamma_\ell H)$. In the case when $\text{Stab}_R(\gamma H) = \text{Stab}_Q(\gamma H)$, we have $\text{Tr}_R^Q w_\gamma = \text{Tr}_{\text{Stab}_Q(\gamma H)}^Q(\gamma \otimes v) = s_\ell$. Otherwise $\text{Stab}_R(\gamma H) < \text{Stab}_Q(\gamma H)$, and we have

$$\begin{aligned} \text{Tr}_R^Q w_\gamma &= \text{Tr}_{\text{Stab}_Q(\gamma H)}^Q \text{Tr}_{\text{Stab}_R(\gamma H)}^{\text{Stab}_Q(\gamma H)} \gamma \otimes v \\ &= |\text{Stab}_Q(\gamma H) : \text{Stab}_R(\gamma H)| \text{Tr}_{\text{Stab}_Q(\gamma H)}^Q \gamma \otimes v = 0. \end{aligned}$$

Therefore we deduce that (iv) holds. \square

7. GENERIC JORDAN TYPES OF MODULES

Let $E = \langle g_1, \dots, g_n \rangle$ be an elementary abelian p -group of p -rank n and let F be an algebraically closed field of characteristic p . Let M be a finite-dimensional FE -module. Let $K = F(\alpha_1, \dots, \alpha_n)$ where $\alpha_1, \dots, \alpha_n$ are indeterminates. With respect to a basis for M , the matrix representing the action of the element

$$1 + \alpha_1(g_1 - 1) + \dots + \alpha_n(g_n - 1)$$

on M has order p . If $[r]$ denotes a unipotent Jordan block of dimension r then the Jordan type of this matrix is $[1]^{s_1} \dots [p]^{s_p}$ for some $s_i \in \mathbb{N}_0$. By Wheeler [26], this Jordan type is independent of the choice of the generators of E . It is called the *generic Jordan type* of the FE -module M . The *stable generic Jordan type* of M is $[1]^{s_1} \dots [p-1]^{s_{p-1}}$. The module M is *generically free* if $s_1 = \dots = s_{p-1} = 0$. For further background on generic Jordan type, we refer the reader to [11].

We summarize below the properties which we need.

Proposition 7.1. *Suppose that E is an elementary abelian p -group of finite order.*

- (i) *The generic Jordan type of a direct sum of modules is the direct sum of the generic Jordan types of the modules.*
- (ii) *Let $0 \rightarrow M_1 \rightarrow M_2 \rightarrow M_3 \rightarrow 0$ be a short exact sequence of FE -modules.*
 - (a) *If M_2 is generically free then M_1 has stable generic Jordan type $[1]^{s_1} \dots [p-1]^{s_{p-1}}$ if and only if M_3 has stable generic Jordan type $[1]^{s_{p-1}} \dots [p-1]^{s_1}$.*
 - (b) *If M_3 is generically free then M_1 and M_2 have the same stable generic Jordan type.*
 - (c) *If M_1 is generically free then M_3 and M_2 have the same stable generic Jordan type.*

- (iii) Let D be a proper subgroup of E and let U be a FD -module. Then the induced module $U\uparrow^E$ is generically free.
- (iv) Let G be a finite group (not necessarily elementary abelian) and suppose that E is a subgroup of G . Let V be an indecomposable FG -module. If $V\downarrow_E$ is not generically free then V has a vertex containing the subgroup E .

Proof. Part (i) follows from [11, Proposition 4.7] and will be used in the proofs of (ii), (iii) and (iv). Since we do not require these ideas elsewhere in our paper, we refer the reader to [11] for details. We write $[\alpha_K](M)$ and $[\alpha_K]^*(M)$ for the generic and stable generic Jordan type of a module M respectively.

For part (ii), we note that the short exact sequence induces the short exact sequence

$$0 \rightarrow [\alpha_K](M_1) \rightarrow [\alpha_K](M_2) \rightarrow [\alpha_K](M_3) \rightarrow 0.$$

For (ii)(a) we have

$$[\alpha_K]^*(M_1) \simeq \Omega(\Omega^{-1}[\alpha_K]^*(M_1)) \simeq \Omega([\alpha_K]^*(M_3)).$$

Hence the stable generic Jordan types of M_1 and M_3 are complementary. For (ii)(b) we note that since M_3 is generically free, the short exact sequence generically splits. Hence

$$[\alpha_K]^*(M_2) \simeq [\alpha_K]^*(M_1) \oplus [\alpha_K]^*(M_3) = [\alpha_K]^*(M_1).$$

The proof of (ii)(c) is similar.

There exists a non-trivial subgroup C of E such that $E = C \times D$. We have $U\uparrow^E \cong FE \otimes_{FD} U \cong F(E/D) \otimes_F U'$ where U' has the same underlying vector space as U and the action of E is given by $(hk)v = kv$ for $h \in C$, $k \in D$ and $v \in U'$. Since $F(E/D)$ is generically free as an FE -module, it follows that $U\uparrow^E$ is generically free, as required in (iii).

For (iv), suppose that V has Q as a vertex. Then there is an FQ -module M such that V is a direct summand of $M\uparrow^G$. By the Mackey decomposition formula $(M\uparrow^G)\downarrow_E$ is a summand of a direct sum of the modules $M_g = ({}^gM\downarrow_{E \cap gQg^{-1}})\uparrow^E$ for some suitable $g \in G$. If $E \cap gQg^{-1}$ is a proper subgroup of E then M_g is generically free by (iii). Since $V\downarrow_E$ is direct summand of $(M\uparrow^G)\downarrow_E$ and, by hypothesis $V\downarrow_E$ is not generically free, there exists $g \in G$ such that $E \cap gQg^{-1} = E$. For this g we have $E \subseteq gQg^{-1}$. \square

Of independent interest, we mention a direct consequence of Proposition 7.1(iv) about hook Specht modules. Let $n \in \mathbb{N}$ and let $d = \lfloor \frac{n}{p} \rfloor$. Let

$$C(d) = \langle (1, 2, \dots, p), \dots, ((d-1)p+1, (d-1)p+2, \dots, dp) \rangle$$

be as defined in Section 2. The second author computed the generic Jordan type of $S^{(n-r, 1^r)} \downarrow_{C(d)}$ in [20, Corollary 4.2, Theorem 4.5]. In particular, the restricted module is not generically free. Hence by Proposition 7.1(iv), we obtain the following result.

Proposition 7.2. *The hook Specht module $S^{(n-r, 1^r)}$ has a vertex containing the maximal elementary abelian p -subgroup $C(d)$ where $d = \lfloor \frac{n}{p} \rfloor$.*

We end this section with a result on the generic Jordan type of a monomial module.

Proposition 7.3. *Let G be a finite group, let $H \leq G$ and let $V = \langle v \rangle \uparrow_H^G$ be a monomial FG -module. If E is an elementary abelian p -subgroup of G then $V \downarrow_E$ has generic Jordan type $[1]^s$ where s is the number of orbits of size 1 of E acting on G/H .*

Proof. We have

$$V \downarrow_E = \bigoplus_g ({}^g \langle v \rangle \downarrow_{E \cap gHg^{-1}}) \uparrow^E$$

where the sum is over a set of representatives for the orbits of E on G/H . Orbits of size 1 correspond to representatives g such that $E \subseteq gHg^{-1}$. In this case,

$$({}^g \langle v \rangle \downarrow_{E \cap gHg^{-1}}) \uparrow^E = {}^g \langle v \rangle \downarrow_E.$$

Since ${}^g \langle v \rangle \downarrow_E$ is 1-dimensional, its generic Jordan type is clearly $[1]$. The result now follows from parts (i) and (iii) of Proposition 7.1. \square

8. PROOF OF THEOREM 1.3 UNDER HYPOTHESIS (I)

Let F be an algebraically closed field of prime characteristic p . As in Section 5 we let $N = \langle e_1, \dots, e_{kp} \rangle$ be the kp -dimensional natural permutation module for \mathfrak{S}_{kp} . Recall from the paragraph preceding Lemma 5.3 that $\bigwedge^r N$ has as a basis the elements $e_{\mathbf{i}} = e_{i_1} \wedge \dots \wedge e_{i_r}$ for $\mathbf{i} \in I^{(r)}$. Considering the action of $H = \mathfrak{S}_{\{1, \dots, r\}} \times \mathfrak{S}_{\{r+1, \dots, kp\}}$ on the generator $e_1 \wedge \dots \wedge e_r$ of $\bigwedge^r N$, we see that

$$(6) \quad \bigwedge^r N \cong \langle v \rangle \uparrow^{\mathfrak{S}_{kp}}$$

where $\langle v \rangle$ affords the representation $\text{sgn} \boxtimes F$ of H . Hence $\bigwedge^r N$ is a monomial module.

Corollary 8.1. *Let E be an elementary abelian p -subgroup of \mathfrak{S}_{kp} . Then the stable generic Jordan type of $(\bigwedge^r N) \downarrow_E$ is $[1]^s$ where s is the number of r -subsets of $\{1, \dots, kp\}$ fixed by E in its action on all r -subsets of $\{1, \dots, kp\}$. In particular, $(\bigwedge^r N) \downarrow_E$ is generically free if $r \not\equiv 0 \pmod{p}$ and E has no fixed point on $\{1, \dots, kp\}$.*

Proof. Since $\bigwedge^r N$ is a monomial module by (6), we may apply Proposition 7.3. Identifying \mathfrak{S}_{kp}/H with the set of r -subsets of $\{1, \dots, kp\}$ we see that orbits of E on \mathfrak{S}_{kp}/H of size 1 correspond to r -subsets of $\{1, \dots, kp\}$ fixed by E . Hence the generic Jordan type of $(\bigwedge^r N) \downarrow_E$ is $[1]^s$ where s is the number of r -subsets of $\{1, \dots, kp\}$ which are fixed under the action of E . Suppose further now that E has no fixed point on the set $\{1, \dots, kp\}$. Since all the orbits of E on $\{1, \dots, kp\}$ have sizes p^j for some $j \geq 1$, we have $s = 0$ if r is not divisible by p . \square

We are now ready to prove Theorem 1.3 under the hypothesis (i).

Proof of Theorem 1.3 under the hypothesis (i). Since E has at least two orbits of size p on $\{1, \dots, kp\}$, E is a subgroup of a maximal elementary abelian p -subgroup $E(m_1, \dots, m_t)$ where $m_1 \geq 2$. To show that E is contained in a vertex of the Specht module $S^{(kp-p, 1^p)}$, it suffices to show this inclusion for $E(m_1, \dots, m_t)$. Hence, we may assume that $E = E(m_1, m_2, \dots, m_t)$ where $m_1 \geq 2$. Notice that E has no fixed point on the set $\{1, \dots, kp\}$. By Proposition 7.1(iv), it suffices to show that $S^{(kp-p, 1^p)} \downarrow_E \cong \bigwedge^p S^{(kp-1, 1)} \downarrow_E$ is not generically free. We truncate the long exact sequence in Proposition 5.2 to obtain

$$0 \longrightarrow \bigwedge^p S^{(kp-1, 1)} \longrightarrow \bigwedge^p N \xrightarrow{\delta_p} \bigwedge^{p-1} N \xrightarrow{\delta_{p-1}} \dots \longrightarrow N \xrightarrow{\delta_1} F \longrightarrow 0.$$

Restrict each term in the long exact sequence to E . For each $1 \leq i \leq p$, we have a short exact sequence

$$0 \longrightarrow \bigwedge^i S^{(kp-1, 1)} \downarrow_E \longrightarrow \bigwedge^i N \downarrow_E \xrightarrow{\delta_i} \bigwedge^{i-1} S^{(kp-1, 1)} \downarrow_E \longrightarrow 0.$$

By Corollary 8.1, $\bigwedge^i N \downarrow_E$ is generically free if $1 \leq i \leq p-1$. Since $F \downarrow_E$ has generic Jordan type $[1]$, it follows from Proposition 7.1(ii)(a) that $S^{(kp-1, 1)} \downarrow_E$ has stable generic Jordan type $[p-1]$. Repeating this argument, with $\bigwedge^2 S^{(kp-1, 1)} \downarrow_E$ up to $\bigwedge^{p-1} S^{(kp-1, 1)} \downarrow_E$, we find that $\bigwedge^{p-1} S^{(kp-1, 1)}$ has stable generic Jordan type $[1]$.

Suppose for a contradiction that $\bigwedge^p S^{(kp-1, 1)} \downarrow_E$ is generically free. Then, by Proposition 7.1(ii)(c), $\bigwedge^p N \downarrow_E$ has stable generic Jordan type $[1]$. On the other hand, since $E = E(m_1, m_2, \dots, m_t)$ where $m_1 \geq 2$, there are exactly m_1 orbits of E on $\{1, \dots, kp\}$ of size p . Each such orbit corresponds to a p -subset of $\{1, \dots, kp\}$ fixed by E so, by Corollary 8.1, $\bigwedge^p N \downarrow_E$ has stable generic Jordan type $[1]^{m_1}$. The contradiction shows that $S^{(kp-p, 1^p)} \downarrow_E$ is not generically free. This completes the proof. \square

9. PROOF OF THEOREM 1.3 UNDER HYPOTHESIS (II)

Let F be a field of prime characteristic p . Under the hypothesis (ii) of Theorem 1.3, we have that $k \geq p$. As in Section 5, we let $N = \langle e_1, \dots, e_{kp} \rangle$ be the

kp -dimensional natural permutation module for \mathfrak{S}_{kp} and define $S^{(kp-1,1)}$ to be the submodule with F -basis $\{e_2 - e_1, \dots, e_{kp} - e_1\}$. Let $W = \bigwedge^p S^{(kp-1,1)}$. By Proposition 5.1 we have $W \cong S^{(kp-p,1^p)}$.

Let T be a Sylow p -subgroup of the symmetric group on $\{p^2 + 1, \dots, kp\}$. Let Q be the subgroup of \mathfrak{S}_{kp} generated by T together with

$$\begin{aligned}\alpha &= (1, 2, \dots, p) \cdots (p^2 - p + 1, p^2 - p + 2, \dots, p^2), \\ \beta &= (1, p + 1, \dots, p^2 - p + 1) \cdots (p, 2p, \dots, p^2),\end{aligned}$$

and let

$$w = (e_1 + e_{p+1} + \cdots + e_{(p-1)p+1}) \wedge \cdots \wedge (e_p + e_{2p} + \cdots + e_{p^2}).$$

Since each factor in the wedge product lies in $S^{(kp-1,1)}$, and w is fixed by α, β and T , we have $w \in W^Q$. Observe that the coefficient of $e_1 \wedge e_2 \wedge \cdots \wedge e_p$ in w is 1.

The main result of this section is Proposition 9.1 below. It gives us the statement of Theorem 1.3(ii) almost immediately.

Proposition 9.1. *If $R < Q$ then no element of $\mathrm{Tr}_R^Q W^R$ has a non-zero coefficient of $e_1 \wedge e_2 \wedge \cdots \wedge e_p$ when expressed in the monomial basis of $\bigwedge^p N$.*

Proof of Theorem 1.3(ii) assuming Proposition 9.1. Let E be an elementary abelian p -subgroup of \mathfrak{S}_{kp} having at least one orbit of size p^2 . There exists $\sigma \in \mathfrak{S}_{kp}$ such that

$$\sigma E \sigma^{-1} \leq \langle \alpha, \beta \rangle \times T = Q.$$

Since $w \in W^Q$ and w has a summand $e_1 \wedge e_2 \wedge \cdots \wedge e_p$ with non-zero coefficient, by Proposition 9.1, we have $W(Q) \neq 0$. Theorem 1.3(ii) now follows from Proposition 6.1. \square

The proof of Proposition 9.1 occupies the remainder of this section. It is divided into two steps: in the first we reduce the proposition to a question about a specific relative trace map on the smaller module $\bigwedge^p S^{(p^2-1,1)}$. In the second step we answer this question (see Proposition 9.5) using a carefully chosen filtration of $\bigwedge^p S^{(p^2-1,1)}$.

Step 1: Reduction. We need the following lemma.

Lemma 9.2. *Let $R \leq Q$, let $B^{(1)}, \dots, B^{(m)}$ be representatives for the orbits of R on the set of p -subsets of $\{1, \dots, kp\}$ and let $\mathbf{i}^{(1)}, \dots, \mathbf{i}^{(m)}$ be the corresponding multi-indices. Then a basis for $(\bigwedge^p N)^R$ is*

$$\mathcal{B} = \{\mathrm{Tr}_{\mathrm{Stab}_R(B^{(k)})}^R e_{\mathbf{i}^{(k)}} : 1 \leq k \leq m\}.$$

Proof. By (6) at the beginning of Section 8, the module $\bigwedge^p N$ is induced from the representation $\mathrm{sgn} \boxtimes F$ of $H = \mathfrak{S}_{\{1, \dots, p\}} \times \mathfrak{S}_{\{p+1, \dots, kp\}}$. Since $\mathrm{sgn}(g) = 1$ for all p -elements $g \in \mathfrak{S}_p$, the hypotheses of Proposition 6.2 are satisfied. The lemma now follows from parts (ii) and (iii) of this proposition, noting

as in the proof of Corollary 8.1 that the coset space \mathfrak{S}_{kp}/H is isomorphic as an \mathfrak{S}_{kp} -set to the set of all p -subsets of $\{1, \dots, kp\}$. \square

We now show that only one subgroup R needs to be considered in Proposition 9.1.

Proposition 9.3. *If R is a proper subgroup of Q such that $e_1 \wedge e_2 \wedge \dots \wedge e_p$ appears with a non-zero coefficient in an element of $\text{Tr}_R^Q W^R$, then $R = \langle \alpha, T \rangle$.*

Proof. Since W^R is a submodule of $(\wedge^p N)^R$, it is sufficient to prove the proposition with W replaced by $\wedge^p N$. Let \mathcal{B} be the basis of $(\wedge^p N)^Q$ given by Lemma 9.2. The unique element of \mathcal{B} containing $e_1 \wedge e_2 \wedge \dots \wedge e_p$ with a non-zero coefficient is

$$\text{Tr}_{\langle \alpha, T \rangle}^Q(e_1 \wedge e_2 \wedge \dots \wedge e_p) = e_1 \wedge e_2 \wedge \dots \wedge e_p + \dots + e_{(p-1)p+1} \wedge e_{(p-1)p+2} \wedge \dots \wedge e_{p^2}.$$

By Proposition 6.2(iv), $\text{Tr}_R^Q(\wedge^p N)^R$ has, as a basis, a subset \mathcal{B}' of \mathcal{B} such that \mathcal{B}' contains $\text{Tr}_{\langle \alpha, T \rangle}^Q(e_1 \wedge e_2 \wedge \dots \wedge e_p)$ if and only if

$$\text{Stab}_Q(\{1, 2, \dots, p\}) = \text{Stab}_R(\{1, 2, \dots, p\}).$$

Since $\text{Stab}_Q(\{1, 2, \dots, p\}) = \langle \alpha, T \rangle$ is a maximal subgroup of Q , this condition holds if and only if $R = \langle \alpha, T \rangle$. \square

The next proposition completes the reduction step.

Proposition 9.4. *If $e_1 \wedge e_2 \wedge \dots \wedge e_p$ appears with a non-zero coefficient in $\text{Tr}_{\langle \alpha, T \rangle}^Q v$ for some $v \in (\wedge^p S^{(kp-1, 1)})^{\langle \alpha, T \rangle}$ then $e_1 \wedge e_2 \wedge \dots \wedge e_p$ appears with a non-zero coefficient in $\text{Tr}_{\langle \alpha \rangle}^{\langle \alpha, \beta \rangle} v'$ for some $v' \in (\wedge^p S^{(p^2-1, 1)})^{\langle \alpha \rangle}$.*

Proof. Let $B^{(1)}, \dots, B^{(L)}$ be representatives for the orbits of $\langle \alpha \rangle$ on the set of p -subsets of $\{1, \dots, p^2\}$ and let $D^{(1)}, \dots, D^{(M)}$ be representatives for the orbits of $\langle \alpha, T \rangle$ on the set of p -subsets of $\{1, \dots, kp\}$ that have non-empty intersection with $\{p^2+1, \dots, kp\}$. Let $\mathbf{i}^{(1)}, \dots, \mathbf{i}^{(L)}$ and $\mathbf{k}^{(1)}, \dots, \mathbf{k}^{(M)}$ be the corresponding multi-indices, and let

$$s_\ell = \text{Tr}_{\text{Stab}_{\langle \alpha, T \rangle}(B^{(\ell)})}^{\langle \alpha, T \rangle} e_{\mathbf{i}^{(\ell)}} \quad \text{for } 1 \leq \ell \leq L,$$

$$t_m = \text{Tr}_{\text{Stab}_{\langle \alpha, T \rangle}(D^{(m)})}^{\langle \alpha, T \rangle} e_{\mathbf{k}^{(m)}} \quad \text{for } 1 \leq m \leq M.$$

By Lemma 9.2, $\{s_\ell : 1 \leq \ell \leq L\}$ is a basis for $(\wedge^p \langle e_1, \dots, e_{p^2} \rangle)^{\langle \alpha \rangle}$ and $\{s_\ell : 1 \leq \ell \leq L\} \cup \{t_m : 1 \leq m \leq M\}$ is a basis for $(\wedge^p N)^{\langle \alpha, T \rangle}$.

Let $v = \sum_{\ell=1}^L \lambda_\ell s_\ell + \sum_{m=1}^M \mu_m t_m$ where the coefficients λ_ℓ and μ_m are in F . By Proposition 5.2, $\wedge^p S^{(kp-1, 1)}$ is the kernel of the map $\delta : \wedge^p N \rightarrow \wedge^{p-1} N$. There is a vector space decomposition $\wedge^{p-1} N = U \oplus U'$ where

$$U = \langle e_{\mathbf{i}} : \mathbf{i} \in I^{(p-1)}, 1 \leq i_1 < \dots < i_{p-1} \leq p^2 \rangle,$$

$$U' = \langle e_{\mathbf{i}} : \mathbf{i} \in I^{(p-1)}, i_{p-1} > p^2 \rangle.$$

For each s_ℓ we have $\delta(s_\ell) \in U$. Let $1 \leq m \leq M$. If $|D^{(m)} \cap \{p^2+1, \dots, kp\}| \geq 2$ then every monomial summand of t_m involves two or more of e_{p^2+1}, \dots, e_{kp} , and so $\delta(t_m)$ lies in U' . Suppose that $D^{(m)} = \{i_1, \dots, i_{p-1}, i_p\}$ where $1 \leq i_1 \leq \dots \leq i_{p-1} \leq p^2$ and $i_p > p^2$. Since $\text{Stab}_{\langle \alpha, T \rangle}(\{i_1, \dots, i_{p-1}\}) = T$, we have

$$\begin{aligned} t_m &= \text{Tr}_{\text{Stab}_T(\{i_p\})}^{\langle \alpha, T \rangle}((e_{i_1} \wedge \dots \wedge e_{i_{p-1}}) \wedge e_{i_p}) \\ &= \left(\text{Tr}_1^{\langle \alpha \rangle}(e_{i_1} \wedge \dots \wedge e_{i_{p-1}}) \right) \wedge \text{Tr}_{\text{Stab}_T(\{i_p\})}^T e_{i_p}. \end{aligned}$$

Since the orbit of T containing i_p has size a multiple of p , it now follows from (5) at the end of Section 5 that

$$\delta(t_m) = \delta\left(\text{Tr}_1^{\langle \alpha \rangle}(e_{i_1} \wedge \dots \wedge e_{i_{p-1}})\right) \wedge \text{Tr}_{\text{Stab}_T(\{i_p\})}^T e_{i_p}.$$

Hence $\delta(t_m) \in U'$. Thus $\delta(\sum_{\ell=1}^L \lambda_\ell s_\ell) \in U$, whereas $\delta(\sum_{m=1}^M \mu_m t_m) \in U'$. Since $\delta(v) = 0$ and $U \cap U' = \{0\}$, it follows that $\delta(\sum_{\ell=1}^L \lambda_\ell s_\ell) = 0$. By Proposition 5.2, $\sum_{\ell=1}^L \lambda_\ell s_\ell \in \bigwedge^p(S^{(p^2-1,1)})$. Since s_ℓ is fixed by α , we have $\sum_{\ell=1}^L \lambda_\ell s_\ell \in \bigwedge^p(S^{(p^2-1,1)})^{\langle \alpha \rangle}$. It is clear that no element of $\text{Tr}_{\langle \alpha, T \rangle}^Q \langle t_1, \dots, t_M \rangle$ contains the monomial $e_1 \wedge e_2 \wedge \dots \wedge e_p$ with a non-zero coefficient. Therefore $v' = \sum_{\ell=1}^L \lambda_\ell s_\ell$ has the properties required in the proposition. \square

Step 2: Proof of the reduced version of Proposition 9.1. Fix $u \in (\bigwedge^p S^{(p^2-1,1)})^{\langle \alpha \rangle}$. By Propositions 9.3 and 9.4, to prove Proposition 9.1, it suffices to show that the coefficient of $e_1 \wedge e_2 \wedge \dots \wedge e_p$ in $\text{Tr}_{\langle \alpha \rangle}^{\langle \alpha, \beta \rangle} u$ is zero. We do this by analysing the behaviour of this trace map on a filtration of $(\bigwedge^p S^{(p^2-1,1)})^{\langle \alpha \rangle}$. From now on we work inside $\bigwedge^p \langle e_1, \dots, e_{p^2} \rangle$. To simplify the notation, let $W = \bigwedge^p S^{(p^2-1,1)}$, and let $M = \bigwedge^p \langle e_1, \dots, e_{p^2} \rangle$. Let $I = I^{(p)}$ and let $J = J^{(p)}$, where these sets of multi-indices are as defined in Section 5 when $n = p^2$. Thus $I = \{(i_1, \dots, i_p) : 1 \leq i_1 < \dots < i_p \leq p^2, i_1, \dots, i_r \in \mathbb{N}\}$ and $J = \{\mathbf{i} \in I : i_1 > 1\}$.

Let $\Gamma = \{1, 2, \dots, p\}$. For $0 \leq c \leq p$, let

$$I_c = \{\mathbf{i} \in I : i_1, \dots, i_c \in \Gamma, i_{c+1}, \dots, i_p \notin \Gamma\}$$

and let $M_c = \langle e_{\mathbf{i}} : \mathbf{i} \in I_c \rangle$. Since Γ is an orbit of α , the subspaces M_c are invariant under α , so as an $F\langle \alpha \rangle$ -module we have $M = M_0 \oplus M_1 \oplus \dots \oplus M_p$. For $0 \leq c \leq p$, let

$$W_c = W \cap (M_c \oplus \dots \oplus M_p).$$

Since M_p is spanned by $e_1 \wedge e_2 \wedge \dots \wedge e_p$ and $\delta(e_1 \wedge e_2 \wedge \dots \wedge e_p) \neq 0$, we have $W_p = \{0\}$. The required result that the coefficient of $e_1 \wedge e_2 \wedge \dots \wedge e_p$ in $\text{Tr}_{\langle \alpha \rangle}^{\langle \alpha, \beta \rangle} u$ is zero therefore follows immediately from the case $c = p - 1$ of the next proposition.

Proposition 9.5. *Let $0 \leq c \leq p-1$ and $u \in W^{(\alpha)}$. There exists $v_0 \in W_0^{(\alpha)}$, \dots , $v_c \in W_c^{(\alpha)}$ such that*

- (i) *the coefficient of $e_1 \wedge \dots \wedge e_p$ in $\text{Tr}_{(\alpha)}^{(\alpha, \beta)}(v_0 + \dots + v_c)$ is zero,*
- (ii) *$u \in v_0 + \dots + v_c + W_{c+1}$.*

We prove Proposition 9.5 by induction on c , using the following two lemmas. First we need some notation. Fix $m \in \{2, \dots, p\}$ and let $\wp(\Gamma)$ denote the power set of Γ . We define a function $\iota_m : \wp(\Gamma) \rightarrow I$ given by

$$\iota_m(X)_a = \begin{cases} a & \text{if } a \in X, \\ (m-1)p + a & \text{if } a \notin X, \end{cases}$$

for $X \subseteq \Gamma$.

Lemma 9.6. *Let $2 \leq m \leq p$, and let*

$$z_m = \sum_{X \subseteq \Gamma} (-1)^{|X|} e_{\iota_m(X)}.$$

Then

- (i) $z_m \in W^{(\alpha)}$,
- (ii) $z_m \in e_{(m-1)p+1} \wedge \dots \wedge e_{(m-1)p+p} + M_1 \oplus \dots \oplus M_p$,
- (iii) *the coefficient of $e_1 \wedge e_2 \wedge \dots \wedge e_p$ in $\text{Tr}_{(\alpha)}^{(\alpha, \beta)} z_m$ is zero.*

Proof. The monomial summands in

$$\delta(z_m) = \sum_{X \subseteq \Gamma} (-1)^{|X|} \sum_{a=1}^p (-1)^{a-1} e_{\iota_m(X)_1} \wedge \dots \wedge \widehat{e_{\iota_m(X)_a}} \wedge \dots \wedge e_{\iota_m(X)_p}$$

are indexed by pairs (X, a) where $X \subseteq \Gamma$ and $1 \leq a \leq p$. We define an involution \star on the set of such pairs by

$$(X, a)^\star = (X \Delta \{a\}, a).$$

where Δ denotes symmetric difference. The summands for (X, a) and $(X, a)^\star$ cancel. Hence $\delta(z_m) = 0$ and $z_m \in W$. Since $\alpha e_{\iota_m(X)} = e_{\iota_m(\alpha X)}$ and obviously $|\alpha X| = |X|$, we have $z_m \in W^{(\alpha)}$. The unique monomial summand of z_m in M_0 is $e_{\iota_m(\emptyset)} = e_{(m-1)p+1} \wedge \dots \wedge e_{(m-1)p+p}$, so we have (ii). The two summands of z_m that contribute to the coefficient of $e_1 \wedge e_2 \wedge \dots \wedge e_p$ in $\text{Tr}_{(\alpha)}^{(\alpha, \beta)} z_m$ are $e_{\iota_m(\emptyset)}$, which appears with coefficient $+1$, and $e_{\iota_m(\Gamma)} = e_1 \wedge e_2 \wedge \dots \wedge e_p$, which appears with coefficient -1 . Their contributions cancel, hence (iii). \square

Lemma 9.7. *Let $c \in \{0, \dots, p-1\}$ and let $\mathbf{j} = (j_1, \dots, j_p) \in J \cap I_c$. Define*

$$w(\mathbf{j}) = \sum_{\ell=0}^{p-1} \alpha^\ell \delta(e_1 \wedge e_{\mathbf{j}}).$$

Then

- (i) $w(\mathbf{j}) \in W_c^{(\alpha)}$,

- (ii) $w(\mathbf{j}) \in \sum_{\ell=0}^{p-1} \alpha^\ell (\delta(e_1 \wedge e_{\mathbf{d}}) \wedge e_{\mathbf{k}}) + M_{c+1}$, where $\mathbf{d} = (j_1, \dots, j_c)$ and $\mathbf{k} = (j_{c+1}, \dots, j_p)$.
- (iii) the coefficient of $e_1 \wedge e_2 \wedge \dots \wedge e_p$ in $\text{Tr}_{\langle \alpha \rangle}^{(\alpha, \beta)} w(\mathbf{j})$ is zero.

Proof. We will use the following facts. The orbits of $\langle \alpha \rangle$ on $\{1, \dots, p^2\}$ are $\Gamma, \beta(\Gamma), \dots, \beta^{p-1}(\Gamma)$, and so are permuted by β . Let $b \in \{1, \dots, p-1\}$ and let $\mathbf{i} = e_{i_1} \wedge \dots \wedge e_{i_p} \in I_b$. Since $\{i_1, \dots, i_p\}$ meets at least two orbits of $\langle \alpha \rangle$, the same is true for $\{\beta^\ell i_1, \dots, \beta^\ell i_p\}$ for any $\ell \in \{0, 1, \dots, p-1\}$. Hence the coefficient of $e_1 \wedge e_2 \wedge \dots \wedge e_p$ in $\sum_{\ell=0}^{p-1} \beta^\ell e_{\mathbf{i}}$ is zero. Therefore the coefficient of $e_1 \wedge e_2 \wedge \dots \wedge e_p$ in $\text{Tr}_{\langle \alpha \rangle}^{(\alpha, \beta)}(y)$ is zero, for any $y \in M_b^{(\alpha)}$.

Clearly $w(\mathbf{j}) \in W^{(\alpha)}$. The monomial summands of

$$\alpha^\ell \delta(e_1 \wedge e_{\mathbf{j}}) = \delta(e_{1+\ell} \wedge e_{\alpha^\ell(j_1)} \wedge \dots \wedge e_{\alpha^\ell(j_c)} \wedge e_{\alpha^\ell(j_{c+1})} \wedge \dots \wedge e_{\alpha^\ell(j_p)})$$

are $e_{\alpha^\ell(j_1)} \wedge \dots \wedge e_{\alpha^\ell(j_p)}$ and $e_{1+\ell} \wedge e_{\alpha^\ell(j_1)} \wedge \dots \wedge \widehat{e_{\alpha^\ell(j_a)}} \wedge \dots \wedge e_{\alpha^\ell(j_p)}$ for $1 \leq a \leq p$. The first summand and the summands for a such that $1 \leq a \leq c$ are in M_c , and the rest are in M_{c+1} , hence (i) and (ii) follow. Also, since $w(\mathbf{j}) \in M_c + M_{c+1}$, using the previous paragraph, we obtain (iii) provided that $1 \leq c \leq p-2$. It remains to prove (iii) when $c=0$ or $c=p-1$.

Suppose that $c=0$, so $j_1, \dots, j_p \geq p+1$. (We give an example of this case following the proof.) By (5) at the end of Section 5 we have $\delta(e_1 \wedge e_{\mathbf{j}}) = e_{\mathbf{j}} - e_1 \wedge \delta(e_{\mathbf{j}})$. Hence

$$w(\mathbf{j}) = \sum_{\ell=0}^{p-1} \alpha^\ell e_{\mathbf{j}} - \sum_{\ell=0}^{p-1} \alpha^\ell (e_1 \wedge \delta(e_{\mathbf{j}})).$$

We consider the two summands separately. Let $k, \ell \in \{0, 1, \dots, p-1\}$ and suppose that $\beta^k \alpha^\ell e_{\mathbf{j}} = e_1 \wedge \dots \wedge e_p$. Then $\beta^k \alpha^\ell (\{j_1, \dots, j_p\}) = \{1, \dots, p\}$, so by the observation on the orbits of $\langle \alpha \rangle$ in the first paragraph, $\{j_1, \dots, j_p\} = \beta^{-k}(\Gamma)$ and $\{j_1, \dots, j_p\}$ is an orbit of $\langle \alpha \rangle$. Hence $\sum_{\ell=0}^{p-1} \alpha^\ell e_{\mathbf{j}} = p e_{\mathbf{j}} = 0$. Now let $y = \sum_{\ell=0}^{p-1} \alpha^\ell (e_1 \wedge \delta(e_{\mathbf{j}}))$. Each monomial summand of each $\alpha^\ell (e_1 \wedge \delta(e_{\mathbf{j}}))$ is equal to $e_{\mathbf{i}}$ for some $\mathbf{i} \in I_1$. Hence $y \in M_1^{(\alpha)}$ and by the first paragraph the coefficient of $e_1 \wedge \dots \wedge e_p$ in $\text{Tr}_{\langle \alpha \rangle}^{(\alpha, \beta)} y$ is zero.

Suppose that $c=p-1$. Then $\mathbf{j} = (2, 3, \dots, p, j_p)$ where $j_p \in \{(m-1)p+1, \dots, (m-1)p+p\}$ for some m such that $2 \leq m \leq p$ and

$$w(\mathbf{j}) = \delta(e_1 \wedge e_2 \wedge \dots \wedge e_p \wedge (e_{(m-1)p+1} + \dots + e_{(m-1)p+p})).$$

It now follows from (5) at the end of Section 5 that

$$\begin{aligned} w(\mathbf{j}) &= (-1)^{p-1} \cdot p \cdot e_1 \wedge \dots \wedge e_p + \sum_{i=1}^p \delta(e_1 \wedge \dots \wedge e_p) \wedge e_{(m-1)p+i} \\ &= \sum_{i=1}^p \delta(e_1 \wedge \dots \wedge e_p) \wedge e_{(m-1)p+i} \in M_{p-1}^{(\alpha)}. \end{aligned}$$

By the first paragraph, the proof of (iii) for $c=p-1$ is complete. \square

In the example below we illustrate the case $c = 0$ in the proof of Lemma 9.7.

Example 9.8. Let $p = 3$. Then $\alpha = (1, 2, 3)(4, 5, 6)(7, 8, 9)$ and $\beta = (1, 4, 7)(2, 5, 8)(3, 6, 9)$. Let $\mathbf{j} = (4, 6, 8)$, $\mathbf{j}' = (4, 5, 6) \in J \cap I_0$. By definition

$$\begin{aligned} w(\mathbf{j}) &= \delta(e_1 \wedge e_4 \wedge e_6 \wedge e_8) + \delta(e_2 \wedge e_5 \wedge e_4 \wedge e_9) + \delta(e_3 \wedge e_6 \wedge e_5 \wedge e_7) \\ &= x + y, \end{aligned}$$

where $x = e_4 \wedge e_6 \wedge e_8 + e_5 \wedge e_4 \wedge e_9 + e_6 \wedge e_5 \wedge e_7$ and $y \in M_1^{(\alpha)}$. By the first paragraph of the proof, we deduce that the coefficient of $e_1 \wedge e_2 \wedge e_3$ in $\text{Tr}_{\langle \alpha \rangle}^{(\alpha, \beta)}(y)$ is zero. Moreover, since $\{4, 6, 8\} \neq \beta^\ell(\{1, 2, 3\})$ for any $\ell \in \{0, 1, 2\}$, the coefficient of $e_1 \wedge e_2 \wedge e_3$ in $\text{Tr}_{\langle \alpha \rangle}^{(\alpha, \beta)}(x)$ is also zero. Hence the coefficient of $e_1 \wedge e_2 \wedge e_3$ in $\text{Tr}_{\langle \alpha \rangle}^{(\alpha, \beta)}(w(\mathbf{j}))$ is zero.

We now consider $w(\mathbf{j}')$. Notice that in this case we have

$$\{j_1, j_2, j_3\} = \{4, 5, 6\} = \beta(\{1, 2, 3\}).$$

By definition $w(\mathbf{j}') = x' + y'$ where $x' = e_4 \wedge e_5 \wedge e_6 + e_5 \wedge e_6 \wedge e_4 + e_6 \wedge e_4 \wedge e_5$ and $y' \in M_1^{(\alpha)}$. Since $y' \in M_1$ the only contributions to the coefficient of $e_1 \wedge e_2 \wedge e_3$ in $\text{Tr}_{\langle \alpha \rangle}^{(\alpha, \beta)}(w(\mathbf{j}'))$ come from $\text{Tr}_{\langle \alpha \rangle}^{(\alpha, \beta)}(x')$. But this is equal to zero since $x' = 3(e_4 \wedge e_5 \wedge e_6) = 0$. Hence also in this case we conclude that the coefficient of $e_1 \wedge e_2 \wedge e_3$ in $\text{Tr}_{\langle \alpha \rangle}^{(\alpha, \beta)}(w(\mathbf{j}'))$ is equal to zero.

In the proof of Proposition 9.5 below we use the bilinear form on M defined on the monomial basis of M by

$$(e_{\mathbf{i}}, e_{\mathbf{i}'}) = \begin{cases} 1 & \text{if } \mathbf{i} = \mathbf{i}', \\ 0 & \text{if } \mathbf{i} \neq \mathbf{i}', \end{cases}$$

where $\mathbf{i}, \mathbf{i}' \in I$.

Proof of Proposition 9.5. We work by induction on c . By induction we may assume that $u \in W_c^{(\alpha)}$. For $\mathbf{i} \in I$ let $\mu_{\mathbf{i}} = (u, e_{\mathbf{i}})$, so we have

$$u = \sum_{\mathbf{i} \in I_c \cup \dots \cup I_p} \mu_{\mathbf{i}} e_{\mathbf{i}}.$$

It suffices to find $v \in W_c^{(\alpha)}$ such that the coefficient of $e_1 \wedge e_2 \wedge \dots \wedge e_p$ in $\text{Tr}_{\langle \alpha \rangle}^{(\alpha, \beta)} v$ is zero and $u - v \in W_{c+1}$.

Case $c = 0$. Let

$$v' = \sum_{m=2}^p \mu_{((m-1)p+1, \dots, (m-1)p+p)} z_m$$

where z_m is as defined in Lemma 9.6. Note that the sets $\{(m-1)p+1, \dots, (m-1)p+p\}$ where $2 \leq m \leq p$ are the singleton orbits of $\langle \alpha \rangle$ on the set of all p -subsets of $\{1, \dots, p^2\} \setminus \Gamma$. Let $B^{(1)}, \dots, B^{(S)}$ be representatives for

the remaining orbits of size p , and let $\mathbf{j}^{(1)}, \dots, \mathbf{j}^{(S)} \in J$ be the corresponding multi-indices. Let

$$v'' = \sum_{s=1}^S \mu_{\mathbf{j}^{(s)}} w(\mathbf{j}^{(s)})$$

where $w(\mathbf{j})$ is as defined in Lemma 9.7 and let $v = v' + v''$. It follows from Lemmas 9.6 and 9.7 that $v \in W^{(\alpha)}$ and that the coefficient of $e_1 \wedge e_2 \wedge \dots \wedge e_p$ in $\text{Tr}_{(\alpha)}^{(\alpha, \beta)} v$ is zero.

Let $\mathbf{j} \in J \cap I_0$. Suppose that $\{j_1, \dots, j_p\}$ is not fixed by α . There exists a unique set $B^{(t)}$ and a unique q such that $0 \leq q < p$ and $\alpha^q(\{j_1, \dots, j_p\}) = B^{(t)}$. Note that $\alpha^q e_{\mathbf{j}} = \pm e_{\mathbf{j}^{(t)}}$ where the sign is determined by the permutation that puts $(\alpha^q(j_1), \dots, \alpha^q(j_p))$ in increasing order. By Lemma 9.2 there is a basis of $M^{(\alpha)}$ in which $\text{Tr}_1^{(\alpha)} e_{\mathbf{j}}$ is the unique basis element involving $e_{\mathbf{j}}$. Since $\alpha u = u$, it follows that $(u, e_{\mathbf{j}}) = (u, \alpha^q e_{\mathbf{j}}) = (u, \pm e_{\mathbf{j}^{(t)}})$. We also have $(v, e_{\mathbf{j}^{(t)}}) = (v, \alpha^{-q} e_{\mathbf{j}}) = \pm (v, e_{\mathbf{j}})$ where the sign is as before. By Lemmas 9.6(ii) and 9.7(ii), we have $(u, e_{\mathbf{j}^{(t)}}) = \mu_{\mathbf{j}^{(t)}} = (v, e_{\mathbf{j}^{(t)}})$. It follows that $(u, e_{\mathbf{j}}) = (v, e_{\mathbf{j}})$. Therefore $u - v \in W_1$.

Case $1 \leq c \leq p - 1$. Let X be the set of all $(p - c)$ -subsets of $\{1, \dots, p^2\} \setminus \Gamma$. Each orbit of $\langle \alpha \rangle$ on X has size p . Let $B^{(1)}, \dots, B^{(S)}$ be representatives for these orbits and let $\mathbf{k}^{(1)}, \dots, \mathbf{k}^{(S)} \in I^{(p-c)}$ be the corresponding multi-indices. Define

$$D = \{(i_1, \dots, i_c) \in I^{(c)} : i_1, \dots, i_c \in \Gamma \setminus \{1\}\}.$$

For each $\mathbf{k}^{(s)}$ define

$$v_{\mathbf{k}^{(s)}} = \sum_{\mathbf{d} \in D} \mu_{\mathbf{d} : \mathbf{k}^{(s)}} w(\mathbf{d} : \mathbf{k}^{(s)})$$

where $\mathbf{d} : \mathbf{k}^{(s)} \in I$ is the multi-index obtained by concatenating \mathbf{d} and $\mathbf{k}^{(s)}$. Let

$$v = \sum_{s=1}^S v_{\mathbf{k}^{(s)}}.$$

It follows from parts (i) and (iii) of Lemma 9.7 that $v \in W_c^{(\alpha)}$ and that the coefficient of $e_1 \wedge e_2 \wedge \dots \wedge e_p$ in $\text{Tr}_{(\alpha)}^{(\alpha, \beta)} v$ is zero. It remains to show that $u - v \in W_{c+1}$.

Let $\mathbf{i} \in I_c$. There exists a unique set $B^{(t)}$ and a unique q such that $0 \leq q \leq p - 1$ and $\alpha^q(\{i_{c+1}, \dots, i_p\}) = B^{(t)}$. Let $\mathbf{g} = (g_1, \dots, g_c)$ be the multi-index corresponding to the set $\alpha^q(\{i_1, \dots, i_c\}) \subseteq \Gamma$. We have $\alpha^q e_{\mathbf{i}} = \pm e_{\mathbf{g} : \mathbf{k}^{(t)}}$, where the sign is determined by the permutation putting $(\alpha^q(i_1), \dots, \alpha^q(i_p))$ into increasing order. As in the previous case we have $(u, e_{\mathbf{i}}) = (u, \alpha^q e_{\mathbf{i}}) = \pm (u, e_{\mathbf{g} : \mathbf{k}^{(t)}})$ and $(v, e_{\mathbf{g} : \mathbf{k}^{(t)}}) = (v, \alpha^{-q} e_{\mathbf{i}}) = \pm (v, e_{\mathbf{i}})$

where the signs agree. Therefore $(u, e_i) = (v, e_i)$, if and only if $(u, e_{\mathbf{g}:\mathbf{k}^{(t)}}) = (v, e_{\mathbf{g}:\mathbf{k}^{(t)}})$. By Lemma 9.7(ii) we have

$$\begin{aligned}
(v, e_{\mathbf{g}:\mathbf{k}^{(t)}}) &= \sum_{\mathbf{d} \in D} \mu_{\mathbf{d}:\mathbf{k}^{(t)}} \left(\sum_{\ell=0}^{p-1} \alpha^\ell (\delta(e_1 \wedge e_{\mathbf{d}}) \wedge e_{\mathbf{k}^{(t)}}), e_{\mathbf{g}:\mathbf{k}^{(t)}} \right) \\
&= \sum_{\mathbf{d} \in D} \mu_{\mathbf{d}:\mathbf{k}^{(t)}} (\delta(e_1 \wedge e_{\mathbf{d}}) \wedge e_{\mathbf{k}^{(t)}}), e_{\mathbf{g}:\mathbf{k}^{(t)}}) \\
(7) \qquad &= \sum_{\mathbf{d} \in D} \mu_{\mathbf{d}:\mathbf{k}^{(t)}} (\delta(e_1 \wedge e_{\mathbf{d}}), e_{\mathbf{g}}).
\end{aligned}$$

On the other hand, by (4) in Section 5 and Lemma 5.3 we have

$$u = \sum_{\mathbf{d} \in D} \mu_{\mathbf{d}:\mathbf{k}^{(t)}} \delta(e_1 \wedge e_{\mathbf{d}} \wedge e_{\mathbf{k}^{(t)}}) + u'$$

where u' is an F -linear combination of elements $\delta(e_1 \wedge e_j)$ of the standard basis for $\mathbf{j} \in J'$, where

$$J' = \{\mathbf{j} \in (I_c \cup \dots \cup I_p) \cap J : \mathbf{j} \neq \mathbf{d} : \mathbf{k}^{(t)} \text{ for any } \mathbf{d} \in D\}.$$

Note that $(e_{\mathbf{g}:\mathbf{k}^{(t)}}, \delta(e_1 \wedge e_j))$ is zero for all $\mathbf{j} \in J'$. (This is clear if $g_1 \neq 1$, since then $(e_{\mathbf{g}:\mathbf{k}^{(t)}}, \delta(e_1 \wedge e_j)) = (e_{\mathbf{g}:\mathbf{k}^{(t)}}, e_j) = 0$. If $g_1 = 1$ then $\mathbf{k}^{(t)}$ must be a subsequence of \mathbf{j} ; since $\mathbf{j} \in I_c \cup \dots \cup I_p$, it follows that $\mathbf{j} \in I_c$, and all the entries of \mathbf{j} not in $\mathbf{k}^{(t)}$ lie in Γ , and so $\mathbf{j} = \mathbf{d} : \mathbf{k}^{(t)}$ where $\mathbf{d} \in D$. But then $\mathbf{j} \notin J'$.) Hence $(e_{\mathbf{g}:\mathbf{k}^{(t)}}, u') = 0$. It follows from (5) at the end of Section 5 that

$$u \in \sum_{\mathbf{d} \in D} \mu_{\mathbf{d}:\mathbf{k}^{(t)}} \delta(e_1 \wedge e_{\mathbf{d}}) \wedge e_{\mathbf{k}^{(t)}} + u' + M_{c+1}$$

and so

$$(u, e_{\mathbf{g}:\mathbf{k}^{(t)}}) = \sum_{\mathbf{d} \in D} \mu_{\mathbf{d}:\mathbf{k}^{(t)}} (\delta(e_1 \wedge e_{\mathbf{d}}), e_{\mathbf{g}}).$$

Comparing with (7) we get $(v, e_{\mathbf{g}:\mathbf{k}^{(t)}}) = (u, e_{\mathbf{g}:\mathbf{k}^{(t)}})$, as required. Hence $u - v \in W_{c+1}$. \square

The proof of Proposition 9.1 is now complete.

10. OPEN PROBLEMS

We end with some open problems suggested by our three main theorems.

Sylow subgroups. Say that a group G is *p-elementarily large* if it has the property in Theorem 1.1 that if Q is a p -subgroup of G containing a G -conjugate of every elementary abelian p -subgroup of G then Q is a Sylow p -subgroup of G . As remarked in the introduction, not every group is p -elementarily large. For example, if G has a quaternionic Sylow 2-subgroup then any non-trivial 2-subgroup of G contains the unique elementary abelian 2-subgroup of G up to G -conjugacy. An abelian group is p -elementarily large if and only if its p -Sylow subgroup is elementary abelian.

Problem 10.1. *Let p be a prime. Find sufficient conditions for a finite group G to be p -elementarily large. In particular, which finite simple groups are p -elementarily large?*

Vertices. By Theorem 1.2, the following conjecture holds when $r = p$, $k \equiv 1 \pmod{p}$ and $k \not\equiv 1 \pmod{p^2}$.

Conjecture 10.2. *Let p be an odd prime. The hook Specht module $S^{(kp-r, 1^r)}$ has a Sylow p -subgroup of \mathfrak{S}_{kp} as a vertex.*

We hope that this conjecture will motivate new methods for computing vertices. The following two examples show some of the limitations of the main methods in this paper. Take $p = 3$. Let $E = E(0, 0, 1)$; thus E is an elementary abelian 3-subgroup of \mathfrak{S}_{27} acting regularly on the set $\{1, \dots, 27\}$. Since the dimension of $\bigwedge^3 S^{(26,1)}$ is $\binom{26}{3}$, which is coprime to 3, $\bigwedge^3 S^{(26,1)}$ has a Sylow 3-subgroup of \mathfrak{S}_{27} as a vertex. However calculations using MAGMA show that $(\bigwedge^3 S^{(26,1)})(E) = 0$, so the Brauer correspondence fails to detect that E is contained in a vertex of $S^{(24,1^3)}$. Similarly if $E = E(1, 1) \leq \mathfrak{S}_{12}$ then the $FE(1,1)$ -module $S^{(9,1^3)} \downarrow_{E(1,1)}$ is generically free, but by Theorem 1.2, $S^{(9,1^3)}$ has a Sylow 3-subgroup of \mathfrak{S}_{12} as a vertex.

Sources. If V is an FG -module with vertex Q and V is a direct summand of $U \uparrow^G$ where U is an indecomposable FQ -module, then U is said to be a *source* of V ; the module U is well-defined up to conjugacy by $N_G(Q)$. If n is not divisible by p then $S^{(n-1,1)}$ is a direct summand of the permutation module $\langle e_1, \dots, e_n \rangle$ and it easily follows that $\bigwedge^r S^{(n-1,1)}$ has trivial source whenever it is indecomposable. When $p = 2$ and n is even it follows from the result of Murphy and Peel mentioned in the introduction that the source of $S^{(n-r, 1^r)}$ is $S^{(n-r, 1^r)} \downarrow_P$ where P is a Sylow 2-subgroup of \mathfrak{S}_n . The indirect methods used in this paper give no information about the source of $S^{(kp-p, 1^p)}$ when p is odd.

Problem 10.3. *Determine the source of $S^{(kp-p, 1^p)}$ when p is odd.*

Problem 10.4. *Is there an indecomposable Specht module with a vertex properly contained in the defect group of its p -block that does not have trivial source?*

REFERENCES

- [1] A. Adem, D. Karagueuzian, R. J. Milgram and K. Umland, *The cohomology of the Lyons group and double covers of alternating groups*, J. Algebra **208** (1998), no. 2, 452–479.
- [2] A. Adem and R. J. Milgram, *Cohomology of Finite Groups*, Volume 309, Springer-Verlag 2004.
- [3] J. L. Alperin, *Weights for finite groups*, Proc. Sympos. Pure Math. **47** (1987) 369–379.

- [4] W. Bosma, J. Cannon, and C. Playoust, *The Magma algebra system. I. The user language*, J. Symbolic Comput. **24** (1997), no. 3-4, 235–265, Computational algebra and number theory (London, 1993).
- [5] M. Broué, *On Scott modules and p -permutation modules: an approach through the Brauer homomorphism*, Proc. Amer. Math. Soc. **93**, no. 3 (1985), 401–408.
- [6] S. Danz and K. Erdmann, *The vertices of a class of Specht modules and simple modules for symmetric groups in characteristic 2*, Algebra Colloq. **19** (2012), no. 1 987–1016.
- [7] S. Danz and E. Giannelli, *Vertices of simple modules of symmetric groups labelled by hook partitions*, J. Group Theory **18**, no. 2, (2014), 313–334.
- [8] S. Danz and K. J. Lim, *Signed Young modules and simple Specht modules*, arXiv:1504.02823v2.
- [9] S. Danz and B. Külshammer, *Vertices of simple modules for symmetric groups: a survey*, Proceedings of the International Conference on Modules and Representation Theory, Presa Univ. Clujeană, Cluj-Napoca, 2009, pp. 61–77.
- [10] S. Donkin, *Symmetric and exterior powers, linear source modules and representations of Schur superalgebras*, Proc. London Math. Soc. **83** (2001), 647–680.
- [11] E. Friedlander, J. Pevtsova and A. Suslin, *Generic and maximal Jordan types*, Invent. Math. **168** (2007), no. 3, 485–522.
- [12] E. Giannelli, *A lower bound on the vertices of Specht modules for symmetric groups*, Arch. Math. (Basel) **103** (2014), no. 1, 1–9.
- [13] J. A. Green, *On the indecomposable representations of a finite group*, Math. Zeitschrift **70** (1958/59), 430–445.
- [14] J. A. Green, *A transfer theorem for modular representations*, J. Algebra **1** (1964), 73–84.
- [15] W. Hamernik, *Specht modules and the radical of the group ring over the symmetric group γ_p* , Comm. Algebra **4**, (1976), 435–475.
- [16] D. J. Hemmer, *Irreducible Specht modules are signed Young modules*, J. Algebra **305** (2006), no. 1, 433–441.
- [17] B. Huppert and N. Blackburn, *Finite Groups II*, Springer, Berlin, 1982.
- [18] G. D. James. *The representation theory of the symmetric groups*, volume 682 of *Lecture Notes in Mathematics*. Springer, Berlin, 1978.
- [19] G. James and A. Kerber, *The representation theory of the symmetric group*, Encyclopedia of Mathematics and its Applications, vol. 16, Addison-Wesley Publishing Co., Reading, Mass., 1981.
- [20] K. J. Lim, *The complexity of the Specht modules corresponding to hook partitions*, Arch. Math. (Basel) **93** (2009), no. 1, 11–22.
- [21] K. J. Lim, *Specht modules with abelian vertices*, J. Algebraic Combin. **35** (2012), no. 1, 157–171.
- [22] J. Müller and R. Zimmermann, *Green vertices and sources of simple modules of the symmetric group labelled by hook partitions*, Arch. Math. (Basel) **89** (2007), no. 2, 97–108.
- [23] G. M. Murphy and M. H. Peel, *Vertices of Specht modules*, J. Algebra **86** (1984), 85–97.
- [24] M. H. Peel, *Hook representations of the symmetric groups*, Glasgow Math. J. **12** (1971), 136–149.
- [25] M. H. Peel, *Specht modules and symmetric groups*, J. Algebra **36** (1975), 88–97.
- [26] W. W. Wheeler, *Generic module theory*, J. Algebra **185** (1996), 205–228.

- [27] M. Wildon, *Two theorems on the vertices of Specht modules*, Arch. Math. (Basel) **81** (2003), no. 5, 505–511.
- [28] M. Wildon, *Vertices of Specht modules and blocks of the symmetric group*, J. Algebra **323** (2010), no. 8, 2243–2256.

(E. Giannelli) DEPARTMENT OF MATHEMATICS, ROYAL HOLLOWAY, UNIVERSITY OF LONDON, UNITED KINGDOM.

E-mail address: `Eugenio.Giannelli.2011@live.rhul.ac.uk`

(K. J. Lim) DIVISION OF MATHEMATICAL SCIENCES, NANYANG TECHNOLOGICAL UNIVERSITY, SPMS-MAS-03-01, 21 NANYANG LINK, SINGAPORE 637371.

E-mail address: `limkj@ntu.edu.sg`

(M. Wildon) DEPARTMENT OF MATHEMATICS, ROYAL HOLLOWAY, UNIVERSITY OF LONDON, UNITED KINGDOM.

E-mail address: `mark.wildon@rhul.ac.uk`