

ON SIGNED YOUNG PERMUTATION MODULES AND SIGNED p -KOSTKA NUMBERS

EUGENIO GIANNELLI, KAY JIN LIM, WILLIAM O'DONOVAN, AND MARK WILDON

ABSTRACT. We prove the existence and main properties of signed Young modules for the symmetric group, using only basic facts about symmetric group representations and the Broué correspondence. We then prove new reduction theorems for the signed p -Kostka numbers, defined to be the multiplicities of signed Young modules as direct summands of signed Young permutation modules. We end by classifying the indecomposable signed Young permutation modules and determining their endomorphism algebras.

1. INTRODUCTION

Let F be a field of odd prime characteristic p and let \mathfrak{S}_n denote the symmetric group of degree n . In this article we investigate the modular structure of the p -permutation $F\mathfrak{S}_n$ -modules defined by inducing a linear representation of a Young subgroup of \mathfrak{S}_n to \mathfrak{S}_n .

Let $\mathcal{P}^2(n)$ be the set of all pairs of partitions $(\alpha|\beta)$ such that $|\alpha| + |\beta| = n$. For $(\alpha|\beta) \in \mathcal{P}^2(n)$, the *signed Young permutation module* $M(\alpha|\beta)$ is the $F\mathfrak{S}_n$ -module defined by

$$M(\alpha|\beta) = \text{Ind}_{\mathfrak{S}_\alpha \times \mathfrak{S}_\beta}^{\mathfrak{S}_n} (F(\mathfrak{S}_\alpha) \boxtimes \text{sgn}(\mathfrak{S}_\beta)). \quad (1.1)$$

In [7, page 651], Donkin defines a *signed Young module* to be an indecomposable direct summand of a signed Young permutation module and proves the following theorem.

Theorem 1.1 (Donkin [7]). *There exist indecomposable $F\mathfrak{S}_n$ -modules $Y(\lambda|p\mu)$ for $(\lambda|p\mu) \in \mathcal{P}^2(n)$ with the following properties:*

- (i) *if $(\alpha|\beta) \in \mathcal{P}^2(n)$ then $M(\alpha|\beta)$ is isomorphic to a direct sum of modules $Y(\lambda|p\mu)$ for $(\lambda|p\mu) \in \mathcal{P}^2(n)$ such that $(\lambda|p\mu) \triangleright (\alpha|\beta)$,*
- (ii) $[M(\lambda|p\mu) : Y(\lambda|p\mu)] = 1$,
- (iii) *if $\lambda = \sum_{i=0}^r p^i \lambda(i)$ and $\mu = \sum_{i=0}^{r-1} p^i \mu(i)$ are the p -adic expansions of λ and μ , as defined in (2.1), then $Y(\lambda|p\mu)$ has as a vertex a Sylow p -subgroup of the Young subgroup \mathfrak{S}_ρ , where ρ is the partition of n having exactly $|\lambda(i)| + |\mu(i-1)|$ parts of size p^i for each $i \in \{0, \dots, r\}$.*

The first author is supported by the ERC advanced grant 291512 and the London Mathematical Society Postdoctoral Mobility Grant PMG14-15 02. The second author is supported by Singapore Ministry of Education AcRF Tier 1 grant RG13/14. Part of this work was done while the first and fourth authors visited National University of Singapore in December 2014; this visit was supported by London Mathematical Society grant 41406 and funding from National University of Singapore and Royal Holloway, University of London.

Here $(\lambda|p\mu) \supseteq (\alpha|\beta)$ refers to the dominance order on $\mathcal{P}^2(n)$, as defined in Section 2.3 below and, in (iii), $\mu(-1)$ should be interpreted as the partition of 0.

Donkin's definition of signed Young modules and his proof of his theorem use the Schur superalgebra. In Section 4 we give an independent proof using only basic facts about symmetric group representations and the Broué correspondence for p -permutation modules; our proof shows that the $Y(\lambda|p\mu)$ may be defined by Definition 4.11. (Theorem 1.1 characterizes the signed Young module $Y(\lambda|p\mu)$ as the unique summand of $M(\lambda|p\mu)$ appearing in $M(\alpha|\beta)$ only if $(\lambda|p\mu) \supseteq (\alpha|\beta)$, so the two definitions are equivalent). As a special case we obtain the existence and main properties of the *Young modules*, which we define by $Y^\lambda = Y(\lambda|\emptyset)$. These are precisely the indecomposable summands of the *Young permutation modules* $M^\alpha = M(\alpha|\emptyset)$. We state this result, and discuss the connection with [10], and with the original definition of Young modules via the Schur algebra [19], in Subsection 5.1.

In [15], Hemmer conjectured, motivated by known results on tilting modules for Schur algebras, that the signed Young modules are exactly the self-dual modules for symmetric groups with Specht filtrations. This was shown to be false in [24]; the fourth author later proved in [27] that if $n \geq 66$ and G is a subgroup of \mathfrak{S}_n such that the ordinary character of $M = \text{Ind}_G^{\mathfrak{S}_n} F$ is multiplicity free, then every indecomposable summand of M is a self-dual module with a Specht filtration. Despite the failure of Hemmer's conjecture, it is clear that signed Young modules are of considerable interest. In particular, a strong connection between simple Specht modules and signed Young modules has been established by Hemmer [15] and by Danz and the second author [6]. More precisely, Hemmer showed that every simple Specht module is isomorphic to a signed Young module, and Danz and the second author established their labels.

In Section 6 we study *signed p -Kostka numbers*, defined to be the multiplicities of signed Young modules as direct summands of signed Young permutation modules. These generalize the p -Kostka numbers considered in [11], [12], [16] and [17]. Given the p -Kostka numbers for \mathfrak{S}_n it is routine to calculate the decomposition matrix of \mathfrak{S}_n in characteristic p (see [12, §3]). It is therefore no surprise that a complete understanding of the p -Kostka numbers seems to be out of reach. However, as the references above demonstrate, many partial results and significant advances have been obtained. Our first main theorem is a relation between signed p -Kostka numbers. We refer the reader to Notation 3.8 for the definitions of the composition $\delta^{(0)}$ and the set $\Lambda((\alpha|\beta), \rho)$.

Theorem 1.2. *Let $(\alpha|\beta), (\lambda|p\mu) \in \mathcal{P}^2(n)$. Then*

$$[M(p\alpha|p\beta) : Y(p\lambda|p^2\mu)] \leq [M(\alpha|\beta) : Y(\lambda|p\mu)].$$

Furthermore, if $\delta^{(0)} = \emptyset$ for all $(\gamma|\delta) \in \Lambda((\alpha|\beta), \rho)$ then equality holds.

Example 6.4 shows that strict inequality may hold in Theorem 1.2. This is an important fact, since it appears to rule out a routine proof of Theorem 1.2 using the theory of weights for the Schur superalgebra: we explain this obstacle later in the introduction. However, in

Corollary 6.3, we obtain the following asymptotic stability of the signed p -Kostka numbers

$$[M(\alpha|\beta) : Y(\lambda|p\mu)] \geq [M(p\alpha|p\beta) : Y(p\lambda|p^2\mu)] = [M(p^2\alpha|p^2\beta) : Y(p^2\lambda|p^3\mu)] = \cdots$$

If $\beta = \emptyset$ then the condition on $\delta^{(0)}$ holds for all $(\gamma|\delta) \in \Lambda((\alpha|\beta), \rho)$ and Theorem 1.2 specializes to Gill's result [12, Theorem 1] that $[M^{p\alpha} : Y^{p\lambda}] = [M^\alpha : Y^\lambda]$ for all partitions α and λ of n .

Our second main theorem describes the relation between signed p -Kostka numbers for partitions differing by a p -power of a partition. Let $\mathcal{C}^2(m)$ be the set consisting of all pairs of compositions $(\alpha|\beta)$ such that $|\alpha| + |\beta| = m$. We refer the reader to Equation 5.1 in Subsection 5.2 for the definition of $\ell_p(\lambda|p\mu)$.

Theorem 1.3. *Let m, n and k be natural numbers. Let $(\pi|\tilde{\pi}) \in \mathcal{C}^2(m)$, $(\lambda|p\mu) \in \mathcal{P}^2(m)$, $(\phi|\tilde{\phi}) \in \mathcal{C}^2(n)$ and $(\alpha|p\beta) \in \mathcal{P}^2(n)$. If $k > \ell_p(\lambda|p\mu)$, then*

$$\begin{aligned} [M(\pi + p^k\phi|\tilde{\pi} + p^k\tilde{\phi}) : Y(\lambda + p^k\alpha|p(\mu + p^k\beta))] \\ \geq [M(\pi|\tilde{\pi}) : Y(\lambda|p\mu)] [M(p\phi|p\tilde{\phi}) : Y(p\alpha|p^2\beta)]. \end{aligned}$$

Moreover, if $p^k > \max\{\pi_1, \tilde{\pi}_1\}$, then equality holds.

In particular, taking $\phi = \alpha = (r)$ and $\tilde{\phi} = \beta = \emptyset$, we see that $[M(\pi + p^k(r)|\tilde{\pi}) : Y(\lambda + p^k(r)|p\mu)] \geq M[(\pi|\tilde{\pi}) : Y(\lambda|p\mu)]$ with equality whenever $p^k > \max\{\pi_1, \tilde{\pi}_1\}$.

Our third main theorem classifies the indecomposable signed Young permutation modules.

Theorem 1.4. *Let $(\alpha|\beta) \in \mathcal{P}^2(n)$. The signed Young permutation module $M(\alpha|\beta)$ is indecomposable if and only if one of the following conditions holds.*

- (i) $(\alpha|\beta) = ((m)|(n))$ for some non-negative integers m, n such that either
 - (a) $m = 0$,
 - (b) $n = 0$, or
 - (c) $m + n$ is divisible by p .
- (ii) $(\alpha|\beta)$ is either $((kp - 1, 1)|\emptyset)$ or $(\emptyset|(kp - 1, 1))$ for some $k \in \mathbb{N}$.

In cases (i)(a) and (i)(b), we have $\text{End}_{F\mathfrak{S}_n} M(\alpha|\beta) \cong F$. In the remaining cases we have $\text{End}_{F\mathfrak{S}_n} M(\alpha|\beta) \cong F[x]/\langle x^2 \rangle$.

In particular, Theorem 1.4 classifies all indecomposable Young permutation modules up to isomorphism, recovering [12, Theorem 2] for fields of odd characteristic. Note that the Young permutation module $M^{(n-1,1)} = M((n-1, 1)|\emptyset) = M((n-1)|(1))$ appears in both parts (i) and (ii). If $M(\alpha|\beta)$ is indecomposable then there exist unique partitions λ and μ such that $M(\alpha|\beta) \cong Y(\lambda|p\mu)$. These partitions are determined in Proposition 7.1.

Schur algebras. Our results may be applied to obtain corollaries on modules for the Schur algebra. Fix $n, d \in \mathbb{N}$ with $d \geq n$ and let $\text{GL}_d(F)$ be the general linear group of $d \times d$ matrices over F . Let $\rho : \text{GL}_d(F) \rightarrow \text{GL}_m(F)$ be a representation of $\text{GL}_d(F)$ of dimension m . We say that ρ is a *polynomial representation of degree n* if the matrix coefficients $\rho(X)_{ij}$ for each $i, j \in \{1, \dots, m\}$ are polynomials of degree n in the coefficients

of the matrix X . Given a polynomial representation $\rho : \mathrm{GL}_d(F) \rightarrow \mathrm{GL}(V)$ of degree n , the image of V under the *Schur functor* f is the subspace of V on which the diagonal matrices $\mathrm{diag}(a_1, \dots, a_d) \in \mathrm{GL}_d(F)$ act as $a_1 \dots a_n$. It is easily seen that $f(V)$ is preserved by the permutation matrices in $\mathrm{GL}_d(F)$ that fix the final $d - n$ vectors in the standard basis of F^d . Thus $f(V)$ is a module for $F\mathfrak{S}_n$.

The category of polynomial representations of $\mathrm{GL}_d(F)$ of degree n is equivalent to the category of modules for the Schur algebra $S_F(d, n)$. We refer the reader to [14] for the definition of $S_F(d, n)$ and further background. In this setting, the Schur functor may be defined by $V \mapsto eV$ where $e \in S_F(d, n)$ is an idempotent such that $eS_F(d, n)e \cong F\mathfrak{S}_n$. It follows that f is an exact functor from the category of polynomial representations of $\mathrm{GL}_d(F)$ of degree n to the category of $F\mathfrak{S}_n$ -modules.

Let E denote the natural $\mathrm{GL}_d(F)$ -module. Given $\alpha \in \mathcal{P}(n)$, let $\mathrm{Sym}^\alpha(E)$ and $\bigwedge^\beta(E)$ denote the corresponding divided symmetric and exterior powers of E , defined as quotient modules of $E^{\otimes n}$. The *mixed powers* $\mathrm{Sym}^\alpha E \otimes \bigwedge^\beta E$ for $(\alpha|\beta) \in \mathcal{P}^2(n)$ generate the category of $\mathrm{GL}_d(F)$ -modules of degree n . In [7], Donkin defines a *listing module* to be an indecomposable direct summand of a mixed power. (As the nautical parlance suggests, listing modules generalize tilting modules). By [7, Proposition 3.1c], for each $(\lambda|\mu) \in \mathcal{P}^2(n)$ there exists a unique listing module $\mathrm{List}(\lambda|p\mu)$ such that $f(\mathrm{List}(\lambda|p\mu)) \cong Y(\lambda|p\mu)$. By [7, Proposition 3.1a], we have $f(\mathrm{Sym}^\alpha E \otimes \bigwedge^\beta E) \cong M(\alpha|\beta)$. Moreover, by [7, Proposition 3.1b], the Schur functor induces an isomorphism

$$\mathrm{End}_{\mathrm{GL}_d(F)}(\mathrm{Sym}^\alpha E \otimes \bigwedge^\beta E) \cong \mathrm{End}_{\mathfrak{S}_n}(M(\alpha|\beta)).$$

Thus each of our three main theorems has an immediate translation to a result on multiplicities of listing modules in certain mixed powers. For example Theorem 1.4 classifies the indecomposable $\mathrm{GL}_d(F)$ -mixed powers and shows that each has an endomorphism algebra, as a $\mathrm{GL}_d(F)$ -module, of dimension at most 2. It is also worth noting that many of Gill's results from [12] are reproved in greater generality in the Schur algebra setting in a recent paper of Donkin [8].

Steinberg Tensor Product Formula. As Gill remarks in [12], some of his results can be obtained using weight spaces and the Steinberg Tensor Product Theorem for irreducible representations of $\mathrm{GL}_d(F)$. We explain the connection here, since this remark is also relevant to this work. Let α be a composition of n where $d \geq n$ and let $\xi_\alpha \in S_F(d, n)$ be the idempotent defined in [14, Section 3.2] such that $\xi_\alpha V$ is the α -weight space, denoted V_α , of the $S_F(d, n)$ -module V ; the idempotent e defining the Schur functor is $\xi_{(1^n)}$. For λ a partition of n , let $L(\lambda)$ denote the irreducible representation of $\mathrm{GL}_d(F)$ with highest weight λ , thought of as a module for $S_F(d, n)$. Let $\mathrm{Proj}(\lambda)$ be the projective cover of $L(\lambda)$. By James' original definition of Young modules (this is shown to be equivalent to ours in Subsection 5.1), we have $Y^\lambda = f(\mathrm{Proj}(\lambda))$; moreover $[M^\alpha : Y^\lambda] = [\mathrm{Sym}_\alpha(E) : \mathrm{Proj}(\lambda)] = \dim_F \mathrm{Hom}(S(d, n)\xi_\alpha, L(\lambda)) = \dim_F \xi_\alpha L(\lambda)$. (Here $\mathrm{Sym}_\alpha(E) \subseteq E^{\otimes n}$ is the contravariant dual, as defined in [14, 2.7a], of the quotient module $\mathrm{Sym}^\alpha(E)$ of $E^{\otimes n}$). Thus

$$[M^\alpha : Y^\lambda] = \dim_F L(\lambda)_\alpha. \tag{1.2}$$

As an example of this relationship between p -Kostka numbers and dimensions of weight spaces, we use (1.2) to deduce Theorem 1 in [12]. By the Steinberg Tensor Product Theorem, $L(p\lambda) = L(\lambda)^{\mathcal{F}}$, where \mathcal{F} is the Frobenius map, acting on representing matrices by sending each entry to its p th power. Clearly there is a canonical vector space isomorphism $(L(\lambda)^{\mathcal{F}})_{p\alpha} \cong L(\lambda)_{\alpha}$. Therefore

$$[M^{p\alpha} : Y^{p\lambda}] = \dim_F L(p\lambda)_{p\alpha} = \dim_F L(\lambda)_{\alpha} = [M^{\alpha} : Y^{\lambda}]$$

as required.

Schur superalgebras. Our Theorem 1.2 generalizes the result just proved, so it is natural to ask if it can be proved in a similar way, replacing the Schur algebra with the Schur superalgebra defined in [7]. Let $a, b \in \mathbb{N}$. Given $(\lambda|p\mu) \in \mathcal{P}^2(n)$ where λ has at most a parts and μ has at most b parts, let $L(\lambda|p\mu)$ denote the irreducible module of highest weight $(\lambda|p\mu)$ for the Schur superalgebra $S(a|b, n)$, defined in [7, page 661]. By [7, §2.3], we have

$$[M(\alpha|\beta) : Y(\lambda|p\mu)] = \dim_F L(\lambda|p\mu)_{(\alpha|\beta)}. \quad (1.3)$$

generalizing (1.2).

Let $\mathrm{GL}(a|b)$ denote the super general linear group defined in [4, Section 2]. Since $E^{\otimes n}$ is a generator for the category of polynomial representations of $\mathrm{GL}(a|b)$ of degree n , it follows from [7, page 660, (1)] that the category of such modules is equivalent to the module category of $S(a|b, n)$. Taking the even degree part of $\mathrm{GL}(a|b)$ recovers $\mathrm{GL}_a(F) \times \mathrm{GL}_b(F)$. (More precisely, the even degree part is isomorphic to the product of the affine group schemes corresponding to these two general linear groups). The Frobenius map is identically zero on the odd degree part of $\mathrm{GL}(a|b)$, so induces a map $\mathcal{F} : \mathrm{GL}(a|b) \rightarrow \mathrm{GL}_a(F) \times \mathrm{GL}_b(F)$. Let \mathcal{F}^* be the corresponding inflation functor, sending modules for $\mathrm{GL}_a(F) \times \mathrm{GL}_b(F)$ to modules for $\mathrm{GL}(a|b)$. By [4, Remark 4.6(iii)] we have

$$L(p\lambda|p\mu) = \mathcal{F}^*(L(\lambda) \boxtimes L(\mu))$$

where \boxtimes denotes an outer tensor product. Taking weight spaces we get $L(p\lambda|p\mu)_{(p\alpha|p\beta)} \cong L(\lambda)_{\alpha} \boxtimes L(\mu)_{\beta}$. By (1.3) we have $[M(p\alpha|p\beta) : Y(p\lambda|p\mu)] = \dim_F L(\lambda)_{\alpha} \dim_F L(\mu)_{\beta}$. Replacing μ with $p\mu$ and applying the Steinberg Tensor Product Formula, this implies the asymptotic stability of signed p -Kostka numbers mentioned after Theorem 1.2.

Stated for $\mathrm{GL}(a|b)$ -modules, the remaining part of Theorem 1.2 becomes

$$\dim_F L(p\lambda|p\mu^2)_{(p\alpha|p\beta)} \leq \dim_F L(\lambda|p\mu)_{(\alpha|\beta)}.$$

This does not follow from the results mentioned so far, or from the version of the Steinberg Tensor Product Theorem for $\mathrm{GL}(a|b)$ -modules proved in [23], because the module on the right-hand side is not an inflation. Moreover, translated into this setting, a special case of Example 6.4 shows that $\dim_F L((1)|\emptyset)_{(\emptyset|(1))} = 1$ whereas $\dim_F L((p)|\emptyset)_{(\emptyset|(p))} = \dim_F L((p))_{\emptyset} \dim_F L(\emptyset)_{(p)} = 0$, so it is certainly not the case that equality always holds. (Further examples of this type are given by the general case of Example 6.4). Whether or

not a proof using supergroups is possible, the authors believe that since Theorem 1.2 can be stated within the context of symmetric groups, it deserves a proof in this setting.

Klyachko's Multiplicity Formula. Klyachko's Multiplicity Formula [22, Corollary 9.2] expresses the p -Kostka number $[M^\alpha : Y^\lambda]$ in terms of p -Kostka numbers for p -restricted partitions. Our Corollary 5.2 gives a generalization to signed Young modules. Specializing this result we obtain a symmetric group proof of Klyachko's formula in the form

$$[M^\alpha : Y^\lambda] = \sum_{(\gamma|\emptyset) \in \Lambda((\alpha|\emptyset), \rho)} \prod_{i=0}^r [M^{\gamma^{(i)}} : Y^{\lambda^{(i)}}], \quad (1.4)$$

where $\lambda = \sum_{i=0}^r p^i \lambda^{(i)}$ is the p -adic expansion of λ , ρ is the partition defined in Theorem 1.1(iii) and the set $\Lambda((\alpha|\emptyset), \rho)$ is as defined in Notation 3.8.

Outline. In Section 2 we recall the main ideas concerning the Brauer construction for p -permutation modules and set up our notation for symmetric group modules and modules for wreath products. In Section 3 we find the Broué quotients of signed Young permutation modules. In Section 4 we use these results, together with James' Submodule Theorem, to define Young modules and signed Young modules in the symmetric group setting. We then prove Donkin's Theorem 1.1. We give some immediate corollaries of this theorem in Section 5.2. In Sections 6 and 7, we prove Theorems 1.2, 1.3 and 1.4.

2. PRELIMINARIES

We work with left modules throughout. For background on vertices and sources and other results from modular representation theory we refer the reader to [1]. For an account of the representation theory of the symmetric group we refer the reader to [18] or [20], or for more recent developments, to [21].

2.1. Indecomposable summands. Let G be a finite group. Let M and N be FG -modules. We write $N \mid M$ if N is isomorphic to a direct summand of M . We have already used the notation $[M : N]$ for the number of summands in a direct sum decomposition of M that are isomorphic to the indecomposable module N . This multiplicity is well defined by the Krull–Schmidt Theorem (see [1, Section 4, Lemma 3]). The proof of the following lemma is easy.

Lemma 2.1. *Let M and N be FG -modules, and let N be indecomposable. Suppose that H is a normal subgroup of G acting trivially on both the modules M and N . Let \overline{M} and \overline{N} be the corresponding $F[G/H]$ -modules. Then $[M : N] = [\overline{M} : \overline{N}]$.*

2.2. Broué correspondence. Let G be a finite group. An FG -module V is said to be a p -permutation module if for every Sylow p -subgroup P of G there exists a linear basis of V that is permuted by P . A useful characterization of p -permutation modules is given by the following theorem (see [3, (0.4)]).

Theorem 2.2. *An indecomposable FG -module V is a p -permutation module if and only if there exists a p -subgroup P of G such that $V \mid \text{Ind}_P^G F$; equivalently, V has trivial source.*

It easily follows that the class of p -permutation modules is closed under restriction and induction and under taking direct sums, direct summands and tensor products.

We now recall the definition and the basic properties of Brauer quotients. Given an FG -module V and P a p -subgroup of G , the set of fixed points of P on V is denoted by

$$V^P = \{v \in V : gv = v \text{ for all } g \in P\}.$$

It is easy to see that V^P is an $FN_G(P)$ -module on which P acts trivially. For Q a proper subgroup of P , the relative trace map $\text{Tr}_Q^P : V^Q \rightarrow V^P$ is the linear map defined by

$$\text{Tr}_Q^P(v) = \sum_{g \in P/Q} gv,$$

where the sum is over a complete set of left coset representatives for Q in P . The definition of this map does not depend on the choice of the set of representatives. We observe that

$$\text{Tr}^P(V) = \sum_{Q < P} \text{Tr}_Q^P(V^Q)$$

is an $FN_G(P)$ -module on which P acts trivially. We define the *Brauer quotient* of V with respect to P to be the $F[N_G(P)/P]$ -module

$$V(P) = V^P / \text{Tr}^P(V).$$

If V is an indecomposable FG -module and P is a p -subgroup of G such that $V(P) \neq 0$, then P is contained in a vertex of V . Broué proved in [3] that the converse holds for p -permutation modules.

Theorem 2.3 ([3, Theorem 3.2]). *Let V be an indecomposable p -permutation module and P be a vertex of V . Let Q be a p -subgroup of G . Then $V(Q) \neq 0$ if and only if $Q \leq {}^gP$ for some $g \in G$.*

Here gP denotes the conjugate gPg^{-1} of P . If V is an FG -module with p -permutation basis \mathcal{B} with respect to a Sylow p -subgroup \tilde{P} of G and $P \leq \tilde{P}$, then, taking for each orbit of P on \mathcal{B} the sum of the basis elements in that orbit, we obtain a basis for V^P . Each sum over an orbit of size p or more is a relative trace from a proper subgroup of P . Hence $V(P)$ is isomorphic to the F -span of

$$\mathcal{B}^P = \{v \in \mathcal{B} : gv = v \text{ for all } g \in P\}.$$

Thus Theorem 2.3 has the following corollary.

Corollary 2.4. *Let V be a p -permutation FG -module with p -permutation basis \mathcal{B} with respect to a Sylow p -subgroup of G containing a subgroup P . The Brauer quotient $V(P)$ has \mathcal{B}^P as a basis. Moreover, V has an indecomposable summand with a vertex containing P if and only if $\mathcal{B}^P \neq \emptyset$.*

The next result states what is now known as the Broué correspondence.

Theorem 2.5 ([3, Theorems 3.2 and 3.4]). *An indecomposable p -permutation FG -module V has vertex P if and only if $V(P)$ is a projective $F[N_G(P)/P]$ -module. Furthermore, we have the following statements.*

- (i) *The Brauer map sending V to $V(P)$ is a bijection between the isomorphism classes of indecomposable p -permutation FG -modules with vertex P and the isomorphism classes of indecomposable projective modules for $F[N_G(P)/P]$. Regarded as an $FN_G(P)$ -module, $V(P)$ is the Green correspondent of V .*
- (ii) *Suppose that V has vertex P . If M is a p -permutation FG -module then V is a direct summand of M if and only if $V(P)$ is a direct summand of $M(P)$. Moreover,*

$$[M : V] = [M(P) : V(P)].$$

The following lemma allows the Broué correspondence to be applied to monomial modules such as signed Young permutation modules.

Lemma 2.6. *Let A be a subset of F^\times . Let M be an FG -module with an F -basis $\mathcal{B} = \{m_1, \dots, m_r\}$ such that, if $g \in G$ and $m_i \in \mathcal{B}$ then $gm_i = am_j$ for some $a \in A$ and some $m_j \in \mathcal{B}$. Then, for any p -subgroup P of G , there exist coefficients $a_1, \dots, a_r \in A$ such that $\{a_1m_1, \dots, a_rm_r\}$ is a p -permutation basis of M with respect to P .*

Proof. Let $\{i_1, \dots, i_s\}$ be a subset of $\{1, \dots, r\}$ such that \mathcal{B} is the disjoint union of $\mathcal{B}_1, \dots, \mathcal{B}_s$ where, for each $1 \leq j \leq s$,

$$\mathcal{B}_j = \{m_k : gm_{i_j} = a_g m_k \text{ for some } g \in P \text{ and } a_g \in A\}.$$

Suppose that $gm_{i_j} = am_k$ and $g'm_{i_j} = a'm_k$ for some $g, g' \in P$ and $a, a' \in A$. Then $g^{-1}g'm_{i_j} = a'a^{-1}m_{i_j}$ and, consequently, Fm_{i_j} is a one dimensional $F\langle g^{-1}g' \rangle$ -module. Since P is a p -subgroup, Fm_{i_j} is the trivial $F\langle g^{-1}g' \rangle$ -module. Hence $a = a'$. Thus the coefficient a_g is independent of the choice of g , and depends only on m_{i_j} and m_k .

For each $1 \leq j \leq s$, let

$$\Lambda_j = \{a_k m_k : gm_{i_j} = a_k m_k \text{ for some } g \in P \text{ and } a_k \in A\}.$$

By the previous paragraph, $\bigcup_{j=1}^s \Lambda_j$ is a basis of M . It is sufficient to prove that each Λ_j is permuted by P . Let $x \in P$, and let $a_k m_k, a_{k'} m_{k'} \in \Lambda_j$. Suppose that $x(a_{k'} m_{k'}) = b(a_k m_k)$ for some $b \in F$. We have $gm_{i_j} = a_k m_k$ and $g'm_{i_j} = a_{k'} m_{k'}$ for some $g, g' \in P$. Thus $g^{-1}xg'm_{i_j} = bm_{i_j}$. Repeating the argument in the first paragraph, we see that Fm_{i_j} is the trivial $F\langle g^{-1}xg' \rangle$ -module and so $b = 1$. \square

The Brauer quotient of an outer tensor product of p -permutation modules is easily described.

Lemma 2.7. *Let G_1 and G_2 be finite groups, let M_1, M_2 be p -permutation FG_1 - and FG_2 -modules, and let P_1 and P_2 be p -subgroups of G_1, G_2 , respectively. Then*

$$(M_1 \boxtimes M_2)(P_1 \times P_2) \cong M_1(P_1) \boxtimes M_2(P_2)$$

as a representation of $N_{G_1 \times G_2}(P_1 \times P_2)/(P_1 \times P_2) \cong (N_{G_1}(P_1)/P_1) \times (N_{G_2}(P_2)/P_2)$.

Proof. The statement follows from an easy application of Theorem 2.5. \square

2.3. Partitions and compositions. Let $n \in \mathbb{N}_0$. A *composition* of n is a sequence of non-negative integers $\alpha = (\alpha_1, \dots, \alpha_r)$ such that $\alpha_r \neq 0$ and $\alpha_1 + \dots + \alpha_r = n$. In this case, we write $\ell(\alpha) = r$ and $|\alpha| = n$. The unique composition of 0 is denoted by \emptyset ; we have $\ell(\emptyset) = 0$. The *Young subgroup* \mathfrak{S}_α is the subgroup $\mathfrak{S}_{\alpha_1} \times \dots \times \mathfrak{S}_{\alpha_r}$ of \mathfrak{S}_n , where the i th factor \mathfrak{S}_{α_i} acts on the set $\{\alpha_1 + \dots + \alpha_{i-1} + 1, \dots, \alpha_1 + \dots + \alpha_{i-1} + \alpha_i\}$. Let $\alpha = (\alpha_1, \dots, \alpha_r)$ and $\beta = (\beta_1, \dots, \beta_s)$ be compositions and let $q \in \mathbb{N}$. We denote by $q\alpha$ and $\alpha \bullet \beta$ the compositions of $q|\alpha|$ and $|\alpha| + |\beta|$ defined by

$$\begin{aligned} q\alpha &= (q\alpha_1, \dots, q\alpha_r), \\ \alpha \bullet \beta &= (\alpha_1, \dots, \alpha_r, \beta_1, \dots, \beta_s), \end{aligned}$$

respectively. We set $0\alpha = \emptyset$. We denote by $\alpha + \beta$ the composition of $|\alpha| + |\beta|$ defined by

$$\alpha + \beta = (\alpha_1 + \beta_1, \dots, \alpha_s + \beta_s, \alpha_{s+1}, \dots, \alpha_r)$$

where we have assumed, without loss of generality, that $s \leq r$. We define $\alpha - \beta$ similarly, in the case when $\beta_i \leq \alpha_i$ for each $i \leq s$.

A composition α is a *partition* if it is non-increasing. A partition α is *p -restricted* if $\alpha_i - \alpha_{i+1} < p$ for all $i \geq 1$. We denote the set of compositions, partitions and p -restricted partitions of n by $\mathcal{C}(n)$, $\mathcal{P}(n)$ and $\mathcal{RP}(n)$, respectively. A partition α is *p -regular* if its conjugate α' , defined by $\alpha'_j = |\{i : \alpha_i \geq j\}|$, is p -restricted. It is well known that if λ is a partition then there exist unique p -restricted partitions $\lambda(i)$ for $i \in \mathbb{N}_0$ such that

$$\lambda = \sum_{i \geq 0} p^i \lambda(i). \quad (2.1)$$

We call this expression the *p -adic expansion* of λ .

Let $\mathcal{P}^2(n)$, $\mathcal{C}^2(n)$ and $\mathcal{RP}^2(n)$ be the sets consisting of all pairs $(\lambda|\nu)$ of partitions, compositions and p -restricted partitions, respectively, such that $|\lambda| + |\nu| = n$. Here λ or ν may be the empty composition \emptyset . For $(\lambda|\nu), (\alpha|\beta) \in \mathcal{P}^2(n)$, we say that $(\lambda|\nu)$ *dominates* $(\alpha|\beta)$, and write $(\lambda|\nu) \supseteq (\alpha|\beta)$, if, for all $k \geq 1$, we have

- (a) $\sum_{i=1}^k \lambda_i \geq \sum_{i=1}^k \alpha_i$, and
- (b) $|\lambda| + \sum_{i=1}^k \nu_i \geq |\alpha| + \sum_{i=1}^k \beta_i$.

(As a standing convention we declare that $\lambda_i = 0$ whenever λ is a partition and $i > \ell(\lambda)$). This defines a partial order on the set $\mathcal{P}^2(n)$ called the *dominance order*. This order becomes the usual dominance order on partitions when restricted to the subsets $\{(\lambda|\emptyset) \in \mathcal{P}^2(n)\}$ or $\{(\emptyset|\nu) \in \mathcal{P}^2(n)\}$ of $\mathcal{P}^2(n)$.

2.4. Modules for symmetric groups. Let $n \in \mathbb{N}_0$, let \mathfrak{S}_n be the symmetric group on the set $\{1, \dots, n\}$ and let \mathfrak{A}_n be its alternating subgroup. Given a subgroup H of \mathfrak{S}_n , we denote the trivial representation of H by $F(H)$, and the restriction of the sign representation of \mathfrak{S}_n to H by $\text{sgn}(H)$. In the case when $H = \mathfrak{S}_\gamma$ for some composition γ of n we write $F(\gamma)$ and $\text{sgn}(\gamma)$ for $F(H)$ and $\text{sgn}(H)$, respectively. If $\gamma = (n)$ we reduce the number of parentheses by writing $F(n)$ and $\text{sgn}(n)$, respectively.

For λ a p -regular partition of n , let D^λ be the $F\mathfrak{S}_n$ -module defined by

$$D^\lambda = S^\lambda / \text{rad}(S^\lambda),$$

where S^λ is the Specht module labelled by λ (see [18, Chapter 4]). By [18, Theorem 11.5] each D^λ is simple, and each simple $F\mathfrak{S}_n$ -module is isomorphic to a unique D^λ . The simple $F\mathfrak{S}_n$ -modules can also be labelled by p -restricted partitions. For $\lambda \in \mathcal{RP}(n)$ we set $D_\lambda = \text{soc}(S^\lambda)$. The connection between the two labelings is given by

$$D_\lambda \cong D^\lambda \otimes \text{sgn}(n).$$

For $\lambda \in \mathcal{RP}(n)$, let P^λ denote the projective cover of the simple $F\mathfrak{S}_n$ -module D_λ .

Finally, for $\gamma \in \mathcal{P}(n)$, let χ^γ denote the ordinary irreducible character of S^γ , defined over the rational field.

2.5. Modules for wreath products. Let $m \in \mathbb{N}$ and let G be a finite group. Recall that the multiplication in the group $G \wr \mathfrak{S}_m$ is given by

$$(g_1, \dots, g_m; \sigma)(g'_1, \dots, g'_m; \sigma') = (g_1 g'_{\sigma^{-1}(1)}, \dots, g_m g'_{\sigma^{-1}(m)}; \sigma \sigma'),$$

for $(g_1, \dots, g_m; \sigma), (g'_1, \dots, g'_m; \sigma') \in G \wr \mathfrak{S}_m$. (Our notation for wreath products is taken from [20, Section 4.1]). Let M be an FG -module. The m -fold tensor product of M becomes an $F[G \wr \mathfrak{S}_m]$ -module with the action given by

$$(g_1, \dots, g_m; \sigma) \cdot (v_1 \otimes \dots \otimes v_m) = \text{sgn}(\sigma) g_1 v_{\sigma^{-1}(1)} \otimes \dots \otimes g_m v_{\sigma^{-1}(m)}$$

for $(g_1, \dots, g_m; \sigma) \in G \wr \mathfrak{S}_m$ and $v_1, \dots, v_m \in M$. We denote this module by $\widehat{M}^{\otimes m}$. Note that we have twisted the action of the top group \mathfrak{S}_m by the sign representation. Thus, in the notation of [20, 4.3.14], we have

$$\widehat{M}^{\otimes m} = (\# M)^{\sim} \otimes \text{Inf}_{\mathfrak{S}_m}^{G \wr \mathfrak{S}_m}(\text{sgn}(m)).$$

The 1-dimensional module $\widehat{\text{sgn}(k)}^{\otimes n}$ will be important to us. In our applications k will be a p -power, and so odd. Since a transposition in the top group \mathfrak{S}_n acts on $\{1, \dots, kn\}$ as a product of k disjoint transpositions, and so has odd sign, there is a simpler definition of this module, as $\text{Res}_{\mathfrak{S}_k \wr \mathfrak{S}_n}^{\mathfrak{S}_{kn}} \text{sgn}(kn)$. More generally, given $\alpha \in \mathcal{C}(n)$ and an odd number k , we define

$$\widehat{\text{sgn}(k)}^{\otimes \alpha} = \text{Res}_{(\mathfrak{S}_k \wr \mathfrak{S}_{\alpha_1}) \times \dots \times (\mathfrak{S}_k \wr \mathfrak{S}_{\alpha_r})}^{\mathfrak{S}_{kn}} \text{sgn}(kn). \quad (2.2)$$

For use in the proof of Proposition 4.5 we briefly recall the character theory of the group $C_2 \wr \mathfrak{S}_n$. Let χ^λ denote the irreducible character of \mathfrak{S}_n labelled by $\lambda \in \mathfrak{S}_n$. For $(\lambda|\mu) \in \mathcal{P}^2(n)$, with $|\lambda| = m_1$ and $|\mu| = m_2$, we define $\chi^{(\lambda|\mu)}$ to be the ordinary character of the following module for $C_2 \wr \mathfrak{S}_n$

$$\text{Ind}_{C_2 \wr (\mathfrak{S}_{m_1} \times \mathfrak{S}_{m_2})}^{C_2 \wr \mathfrak{S}_n} \left(\text{Inf}_{\mathfrak{S}_{m_1}}^{C_2 \wr \mathfrak{S}_{m_1}}(\chi^\lambda) \boxtimes \left(\text{Inf}_{\mathfrak{S}_{m_2}}^{C_2 \wr \mathfrak{S}_{m_2}}(\chi^\mu) \otimes \widehat{\text{sgn}(2)}^{\otimes m_2} \right) \right).$$

A standard Clifford theory argument (see for instance [20, Theorem 4.34]) shows that the characters $\chi^{(\lambda|\mu)}$ for $(\lambda|\mu) \in \mathcal{P}^2(n)$ are precisely the irreducible characters of $C_2 \wr \mathfrak{S}_n$.

2.6. Sylow p -subgroups of \mathfrak{S}_n . Let P_p be the cyclic group $\langle (1, 2, \dots, p) \rangle \leq \mathfrak{S}_p$ of order p . Let $P_1 = \{1\}$ and, for $d \geq 1$, set

$$P_{p^{d+1}} = P_{p^d} \wr P_p = \{(\sigma_1, \dots, \sigma_p; \pi) : \sigma_1, \dots, \sigma_p \in P_{p^d}, \pi \in P_p\}.$$

By [20, 4.1.22, 4.1.24], P_{p^d} is a Sylow p -subgroup of \mathfrak{S}_{p^d} .

Let $n \in \mathbb{N}$. Let $n = \sum_{i=0}^r n_i p^i$, where $0 \leq n_i < p$ for $i \in \{0, \dots, r\}$, and $n_r \neq 0$ be the p -adic expansion of n . By [20, 4.1.22, 4.1.24], the Sylow p -subgroups of \mathfrak{S}_n are each conjugate to the direct product $\prod_{i=0}^r (P_{p^i})^{n_i}$. Hence if we define P_n to be a Sylow p -subgroup of the Young subgroup $\prod_{i=0}^r (\mathfrak{S}_{p^i})^{n_i}$ then P_n is a Sylow p -subgroup of \mathfrak{S}_n . The normalizer $N_{\mathfrak{S}_n}(P_n)$ of P_n in \mathfrak{S}_n is denoted by N_n .

Whenever $\rho = (\rho_1, \dots, \rho_r) \in \mathcal{C}(n)$, we denote by P_ρ a Sylow p -subgroup of \mathfrak{S}_ρ , defined so that $P_\rho = \prod_{i=1}^r P_{\rho_i}$. In the special case when

$$\rho = (1^{m_0}, p^{m_1}, \dots, (p^s)^{m_s}) = (\underbrace{1, \dots, 1}_{m_0 \text{ copies}}, \underbrace{p, \dots, p}_{m_1 \text{ copies}}, \dots, \underbrace{p^s, \dots, p^s}_{m_s \text{ copies}}),$$

where $m_i \in \mathbb{N}_0$ for each i , we have $P_\rho = \prod_{i=0}^s (P_{p^i})^{m_i}$; in particular, the group P_ρ has precisely m_i orbits of size p^i on the set $\{1, 2, \dots, n\}$ for each i . We write $N_\rho = N_{\mathfrak{S}_n}(P_\rho)$.

3. THE BRAUER QUOTIENTS OF SIGNED YOUNG PERMUTATION MODULES

In this section, we determine the Brauer quotients of signed Young permutation modules with respect to Sylow subgroups of Young subgroups. Our main result is Proposition 3.12; this generalizes [9, Proposition 1]. The description of the Brauer quotients is combinatorial, using the $(\alpha|\beta)$ -tableaux defined below.

Fix $n \in \mathbb{N}$ and $(\alpha|\beta) \in \mathcal{C}^2(n)$. Suppose that $\alpha = (\alpha_1, \dots, \alpha_r)$ and $\beta = (\beta_1, \dots, \beta_s)$. The *diagram* $[\alpha] \bullet [\beta]$ is the set consisting of the *boxes* $(i, j) \in \mathbb{N}^2$ for i and j such that either $1 \leq i \leq r$ and $1 \leq j \leq \alpha_i$ or $r+1 \leq i \leq r+s$ and $1 \leq j \leq \beta_{i-r}$. A box (i, j) is said to be in *row* i . The subset of $[\alpha] \bullet [\beta]$ consisting of the boxes belonging to the first r rows (respectively, the last s rows) is denoted by $[\alpha] \bullet \emptyset$ (respectively, $\emptyset \bullet [\beta]$).

Definition 3.1. An $(\alpha|\beta)$ -*tableau* T is a bijective function $T : [\alpha] \bullet [\beta] \rightarrow \{1, \dots, n\}$. For $(i, j) \in [\alpha] \bullet [\beta]$, the (i, j) -*entry* of T is $T(i, j)$.

We represent an $(\alpha|\beta)$ -*tableau* T by putting the (i, j) -entry of T in the box (i, j) of the diagram $[\alpha] \bullet [\beta]$. Considering $[\alpha] \bullet \emptyset$ as the Young diagram $[\alpha]$, we denote the α -tableau $T([\alpha] \bullet \emptyset)$ by T_+ . Similarly, we denote the β -tableau $T(\emptyset \bullet [\beta])$ by T_- . It will sometimes be useful to write

$$T = (T_+ | T_-).$$

The $(\alpha|\beta)$ -tableau T is *row standard* if the entries in each row of T are increasing from left to right, i.e. both T_+ and T_- are row standard in the usual sense. We denote by $T^{\alpha|\beta}$ the unique row standard $(\alpha|\beta)$ -tableau such that for all $i, j \in \{1, \dots, n\}$, if i is in row a

of $T^{\alpha|\beta}$ and j is in row b of $T^{\alpha|\beta}$ and $i \leq j$ then $a \leq b$. For example,

$$T^{(2,1)|(3)} = \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & \\ \hline 4 & 5 & 6 \\ \hline \end{array}$$

where the thicker line separates the two parts of the tableau.

Let $\mathcal{T}(\alpha|\beta)$ be the set of all $(\alpha|\beta)$ -tableaux. If $T \in \mathcal{T}(\alpha|\beta)$ and $g \in \mathfrak{S}_n$ then we define $g \cdot T$ to be the $(\alpha|\beta)$ -tableau obtained by applying g to each entry of T , i.e. $(g \cdot T)(i, j) = g(T(i, j))$. This defines an action of \mathfrak{S}_n on the set $\mathcal{T}(\alpha|\beta)$. The vector space $F\mathcal{T}(\alpha|\beta)$ over F with basis $\mathcal{T}(\alpha|\beta)$ is therefore a permutation $F\mathfrak{S}_n$ -module.

For each $T \in \mathcal{T}(\alpha|\beta)$, let $R(T) \leq \mathfrak{S}_n$ be the *row stabilizer* of T in \mathfrak{S}_n , consisting of those $g \in \mathfrak{S}_n$ such that the rows of T and $g \cdot T$ coincide as sets. Then $R(T) = R(T_+) \times R(T_-)$, where $R(T_+)$ and $R(T_-)$ are the row stabilizers of T_+ and T_- respectively, in the usual sense. Denote by $U(\alpha|\beta)$ the subspace of $F\mathcal{T}(\alpha|\beta)$ spanned by

$$\{T - \text{sgn}(g_2)g_1g_2 \cdot T : T \in \mathcal{T}(\alpha|\beta), (g_1, g_2) \in R(T_+) \times R(T_-)\}.$$

In fact $U(\alpha|\beta)$ is an $F\mathfrak{S}_n$ -submodule of $F\mathcal{T}(\alpha|\beta)$, since for all $h \in \mathfrak{S}_n$ and for any $(g_1, g_2) \in R(T_+) \times R(T_-)$ and $T \in \mathcal{T}(\alpha|\beta)$ we have

$$h \cdot (T - \text{sgn}(g_2)g \cdot T) = h \cdot T - \text{sgn}({}^h g_2){}^h g \cdot (h \cdot T) \in U(\alpha|\beta),$$

where $g = g_1g_2$, since ${}^h g \in {}^h R(T) = R(h \cdot T)$ and ${}^h g_2 \in R((h \cdot T)_-)$.

Definition 3.2. For each $T \in \mathcal{T}(\alpha|\beta)$, we write

$$\{T\} = \{(T_+|T_-)\}$$

for the element $T + U(\alpha|\beta) \in F\mathcal{T}(\alpha|\beta)/U(\alpha|\beta)$ and call it an $(\alpha|\beta)$ -*tabloid*.

Note that $g\{T\} = \{g \cdot T\}$ for all $g \in \mathfrak{S}_n$ and $T \in \mathcal{T}(\alpha|\beta)$. If $T, T' \in \mathcal{T}(\alpha|\beta)$ are such that $T_- = T'_-$ and T'_+ is obtained by swapping two entries in the same row of T_+ then $\{T\} = \{T'\}$. On the other hand, if $T_+ = T'_+$ and T'_- is obtained by swapping two entries in the same row of T_- then $\{T'\} = -\{T\}$. The graphical representation of $(\alpha|\beta)$ -tableaux is shown in Example 3.5 below.

Let

$$\Omega(\alpha|\beta) = \{\{T\} : T \text{ is a row standard } (\alpha|\beta)\text{-tableau}\} \subseteq F\mathcal{T}(\alpha|\beta)/U(\alpha|\beta).$$

It is clear that $\Omega(\alpha|\beta)$ is an F -basis of $F\mathcal{T}(\alpha|\beta)/U(\alpha|\beta)$. We write $F\Omega(\alpha|\beta)$ for the $F\mathfrak{S}_n$ -module $F\mathcal{T}(\alpha|\beta)/U(\alpha|\beta)$.

Lemma 3.3. *Let $(\alpha|\beta) \in \mathcal{C}^2(n)$.*

- (i) *The $F\mathfrak{S}_n$ -module $F\Omega(\alpha|\beta)$ is isomorphic to the signed Young permutation $M(\alpha|\beta)$.*
- (ii) *For any p -subgroup P of \mathfrak{S}_n , there exist coefficients $a_{\{T\}} \in \{\pm 1\}$ for each $\{T\} \in \Omega(\alpha|\beta)$, such that*

$$\{a_{\{T\}}\{T\} : \{T\} \in \Omega(\alpha|\beta)\}$$

is a p -permutation basis for $F\Omega(\alpha|\beta) \cong M(\alpha|\beta)$ with respect to P .

Proof. By the remarks after Definition 3.2 there is an isomorphism $F(\alpha) \boxtimes \text{sgn}(\beta) \cong F\{\mathbf{T}^{\alpha|\beta}\}$ of $F[\mathfrak{S}_\alpha \times \mathfrak{S}_\beta]$ -modules. Since $|\Omega(\alpha|\beta)| = \dim_F M(\alpha|\beta)$, part (i) follows from the characterization of induced modules in [1, Section 8, Corollary 3]. Part (ii) follows from Lemma 2.6, since, for all $\{\mathbf{T}\} \in \Omega(\alpha|\beta)$ and $\sigma \in \mathfrak{S}_n$, we have $\sigma\{\mathbf{T}\} = \pm\{\mathbf{T}'\}$ for some $\{\mathbf{T}'\} \in \Omega(\alpha|\beta)$. \square

In view of Lemma 3.3(i), we shall identify $M(\alpha|\beta)$ with $F\Omega(\alpha|\beta)$, so that $M(\alpha|\beta)$ has the set of $(\alpha|\beta)$ -tabloids as a basis.

The next corollary follows from Lemma 3.3 and Corollary 2.4.

Corollary 3.4. *Let $(\alpha|\beta) \in \mathcal{C}^2(n)$.*

- (i) *Let P be a p -subgroup of \mathfrak{S}_n . The $F[N_{\mathfrak{S}_n}(P)/P]$ -module $M(\alpha|\beta)(P)$ has a linear basis consisting of all the $(\alpha|\beta)$ -tabloids $\{\mathbf{T}\}$ that are fixed by P .*
- (ii) *Let $\rho = (1^{n_0}, p^{n_1}, \dots, (p^r)^{n_r})$ be a partition of n . The group*

$$N_\rho/P_\rho \cong \mathfrak{S}_{n_0} \times ((N_p/P_p) \wr \mathfrak{S}_{n_1}) \times \cdots \times ((N_{p^r}/P_{p^r}) \wr \mathfrak{S}_{n_r})$$

acts on the set of P_ρ -fixed $(\alpha|\beta)$ -tabloids by transitively permuting the entries in P_ρ -orbits of size p^i according to \mathfrak{S}_{n_i} and, within each P_ρ -orbit of size p^i , permuting its entries according to N_{p^i}/P_{p^i} , for all $i \in \{0, 1, \dots, r\}$.

More explicitly, the basis in Corollary 3.4(i) consists of all $(\alpha|\beta)$ -tabloids $\{\mathbf{T}\}$ such that \mathbf{T} is row standard and each row of \mathbf{T} is a union of orbits of P on $\{1, \dots, n\}$. This can be seen in the following example.

Example 3.5. Let $p = 3$. Consider the 3-subgroups $Q_1 = \langle (1, 2, 3), (4, 5, 6), (7, 8, 9) \rangle$ and $Q_2 = \langle (4, 5, 6), (7, 8, 9) \rangle$ of \mathfrak{S}_9 . By Corollary 3.4(i), since there are no $((2, 1)|(6))$ -tabloids fixed by Q_1 , we have $M((2, 1)|(6))(Q_1) = 0$. On the other hand, $M((2, 1)|(6))(Q_2)$ has a basis consisting of the $((2, 1)|(6))$ -tabloids

$$\left\{ \begin{array}{c} \overline{1 \ 2} \\ \mathbf{3} \\ \hline 4 \ 5 \ 6 \ 7 \ 8 \ 9 \end{array} \right\}, \quad \left\{ \begin{array}{c} \overline{1 \ 3} \\ \mathbf{2} \\ \hline 4 \ 5 \ 6 \ 7 \ 8 \ 9 \end{array} \right\}, \quad \left\{ \begin{array}{c} \overline{2 \ 3} \\ \mathbf{1} \\ \hline 4 \ 5 \ 6 \ 7 \ 8 \ 9 \end{array} \right\}$$

where the bold line separates each \mathbf{T}_+ from \mathbf{T}_- . Taking $\rho = (1, 1, 1, 3, 3)$ we have $P_\rho = Q_2$ and

$$N_{\mathfrak{S}_9}(Q_2) = \mathfrak{S}_3 \times (N_{\mathfrak{S}_3}(P_3) \wr \mathfrak{S}_2) = \mathfrak{S}_3 \times (\mathfrak{S}_3 \wr \mathfrak{S}_2).$$

The first factor \mathfrak{S}_3 permutes the entries 1, 2, 3 of each tabloid without sign, and the second factor $\mathfrak{S}_3 \wr \mathfrak{S}_2$ permutes the entries 4, 5, 6, 7, 8, 9 with sign. The subgroup Q_2 acts trivially on the tabloids. Thus if $\{\mathbf{U}\}$ and $\{\mathbf{V}\}$ are the first two $((2, 1)|(6))$ -tabloids above then

$$\text{Res}_{\mathfrak{S}_3 \wr \mathfrak{S}_2}^{N_{\mathfrak{S}_9}(Q_2)}(F\{\mathbf{U}\}) \cong \widehat{\text{sgn}(3)}^{\otimes 2}$$

and $(23)(45)\widehat{\{\mathbf{V}\}} = -\{\mathbf{U}\}$. Note that the isomorphism above requires the sign twist in the definition of \widehat{M} for M an $F\mathfrak{S}_m$ -module that we commented on in Section 2.5.

Given $m \in \mathbb{N}$, we define a 1-dimensional $F[N_k \wr \mathfrak{S}_m]$ -module by

$$\widehat{\text{sgn}(N_k)}^{\otimes m} = \text{Res}_{N_k \wr \mathfrak{S}_m}^{\mathfrak{S}_k \wr \mathfrak{S}_m} \widehat{\text{sgn}(k)}^{\otimes m}.$$

Using this we may now define three key families of modules. Let \mathcal{F} denote the bifunctor sending a pair $(U|V)$ where U is an $F\mathfrak{S}_{m_1}$ -module and V is an $F\mathfrak{S}_{m_2}$ -module to the $F[C_2 \wr \mathfrak{S}_m]$ -module

$$\text{Ind}_{C_2 \wr (\mathfrak{S}_{m_1} \times \mathfrak{S}_{m_2})}^{C_2 \wr \mathfrak{S}_m} \left(\text{Inf}_{\mathfrak{S}_{m_1}}^{C_2 \wr \mathfrak{S}_{m_1}}(U) \boxtimes \left(\text{Inf}_{\mathfrak{S}_{m_2}}^{C_2 \wr \mathfrak{S}_{m_2}}(V) \otimes \widehat{\text{sgn}(2)}^{\otimes m_2} \right) \right).$$

Definition 3.6. Let $k \in \mathbb{N}$, let $m \in \mathbb{N}_0$ and let $(\gamma|\delta) \in \mathcal{C}^2(m)$. Let $|\gamma| = m_1$ and $|\delta| = m_2$.

(i) We define $V_k(\gamma|\delta)$ to be the $F[\mathfrak{S}_k \wr \mathfrak{S}_m]$ -module

$$\text{Ind}_{(\mathfrak{S}_k \wr \mathfrak{S}_{m_1}) \times (\mathfrak{S}_k \wr \mathfrak{S}_{m_2})}^{\mathfrak{S}_k \wr \mathfrak{S}_m} \left(\text{Inf}_{\mathfrak{S}_{m_1}}^{\mathfrak{S}_k \wr \mathfrak{S}_{m_1}}(M^\gamma) \boxtimes \left(\text{Inf}_{\mathfrak{S}_{m_2}}^{\mathfrak{S}_k \wr \mathfrak{S}_{m_2}}(M^\delta) \otimes \widehat{\text{sgn}(k)}^{\otimes m_2} \right) \right)$$

(ii) We define $W_k(\gamma|\delta)$ to be the $F[(N_k/P_k) \wr \mathfrak{S}_m]$ -module obtained from

$$\text{Res}_{N_k \wr \mathfrak{S}_m}^{\mathfrak{S}_k \wr \mathfrak{S}_m} V_k(\gamma|\delta)$$

$$\cong \text{Ind}_{N_k \wr (\mathfrak{S}_{m_1} \times \mathfrak{S}_{m_2})}^{N_k \wr \mathfrak{S}_m} \left(\text{Inf}_{\mathfrak{S}_{m_1}}^{N_k \wr \mathfrak{S}_{m_1}}(M^\gamma) \boxtimes \left(\text{Inf}_{\mathfrak{S}_{m_2}}^{N_k \wr \mathfrak{S}_{m_2}}(M^\delta) \otimes \widehat{\text{sgn}(N_k)}^{\otimes m_2} \right) \right)$$

via the canonical surjection $N_k \wr \mathfrak{S}_m \rightarrow (N_k \wr \mathfrak{S}_m)/(P_k)^m \cong (N_k/P_k) \wr \mathfrak{S}_m$.

(iii) For $k \geq 2$, we define $\overline{W}_k(\gamma|\delta)$ to be the $F[C_2 \wr \mathfrak{S}_m]$ -module $\mathcal{F}(M^\gamma|M^\delta)$. We define

$$\overline{W}_1(\gamma|\delta) = \text{Inf}_{\mathfrak{S}_m}^{C_2 \wr \mathfrak{S}_m} W_1(\gamma|\delta).$$

We note that $W_k(\gamma|\delta)$ may equivalently be defined to be the $F[(N_k/P_k) \wr \mathfrak{S}_m]$ -module obtained from

$$\text{Ind}_{N_k \wr (\mathfrak{S}_\gamma \times \mathfrak{S}_\delta)}^{N_k \wr \mathfrak{S}_m} \left(F(N_k \wr \mathfrak{S}_\gamma) \boxtimes \widehat{\text{sgn}(N_k)}^{\otimes \delta} \right) \quad (3.1)$$

via the canonical surjection as in Definition 3.6(ii). We have $W_1(\gamma|\delta) = V_1(\gamma|\delta) \cong M(\gamma|\delta)$ as $F\mathfrak{S}_m$ -modules. When $k \geq 2$, the $F[C_2 \wr \mathfrak{S}_m]$ -module $\overline{W}_k(\alpha|\beta)$ is isomorphic to $V_k(\alpha|\beta)$, considered as a $F[C_2 \wr \mathfrak{S}_m]$ -module via the canonical surjection $\mathfrak{S}_k \wr \mathfrak{S}_m \rightarrow (\mathfrak{S}_k \wr \mathfrak{S}_m)/\mathfrak{A}_k^m \cong (\mathfrak{S}_k/\mathfrak{A}_k) \wr \mathfrak{S}_m \cong C_2 \wr \mathfrak{S}_m$. Similarly, $\overline{W}_k(\alpha|\beta)$ is isomorphic to $\text{Res}_{N_k \wr \mathfrak{S}_m}^{\mathfrak{S}_k \wr \mathfrak{S}_m} V_k(\alpha|\beta)$, considered as a $F[C_2 \wr \mathfrak{S}_m]$ -module via the canonical surjection $N_k \wr \mathfrak{S}_m \rightarrow (N_k \wr \mathfrak{S}_m)/(N_{\mathfrak{A}_k}(P_k))^m \cong C_2 \wr \mathfrak{S}_m$.

Lemma 3.7. For all $k \geq 2$ and all $(\gamma|\delta) \in \mathcal{C}^2(m)$, we have

$$\overline{W}_k(\gamma|\delta) \cong V_2(\gamma|\delta),$$

as $F[C_2 \wr \mathfrak{S}_m]$ -modules.

Proof. It suffices to show that $\widehat{\text{sgn}(k)}^{\otimes m_2} \cong \widehat{\text{sgn}(2)}^{\otimes m_2}$ as $F[C_2 \wr \mathfrak{S}_{m_2}]$ -modules, where $\text{sgn}(k)$ is regarded as an FC_2 -module via the canonical surjection $\mathfrak{S}_k \rightarrow \mathfrak{S}_k/\mathfrak{A}_k \cong C_2$. This is clear since $\text{sgn}(k) \cong \text{sgn}(2)$ as FC_2 -modules in this regard. \square

The following notation will be used to describe the direct summands of the Brauer quotients of the signed Young permutation modules $M(\alpha|\beta)$.

Notation 3.8. Let $(\alpha|\beta) \in \mathcal{C}^2(n)$ and let $\rho = (1^{n_0}, p^{n_1}, (p^2)^{n_2}, \dots, (p^r)^{n_r}) \in \mathcal{C}(n)$. We write $\Lambda((\alpha|\beta), \rho)$ for the set consisting of all pairs of tuples of compositions $(\gamma|\delta) = (\gamma^{(0)}, \gamma^{(1)}, \dots, \gamma^{(r)}|\delta^{(0)}, \delta^{(1)}, \dots, \delta^{(r)})$ such that:

- (1) $\alpha = \sum_{i=0}^r p^i \gamma^{(i)}$, $\beta = \sum_{i=0}^r p^i \delta^{(i)}$, and
- (2) $|\gamma^{(i)}| + |\delta^{(i)}| = n_i$ for each $i \in \{0, \dots, r\}$.

Let $(\alpha|\beta) \in \mathcal{C}^2(n)$. Recall that $\Omega(\alpha|\beta)$ is the basis of $M(\alpha|\beta)$ consisting of all $(\alpha|\beta)$ -tabloids. As remarked after Corollary 3.4, an F -basis of $M(\alpha|\beta)(P_\rho)$ is obtained by taking those $(\alpha|\beta)$ -tabloids $\{(T_+|T_-)\} \in \Omega(\alpha|\beta)$ such that the rows of T_+ and T_- are unions of the orbits of P_ρ . Given such a basis element $\{(T_+|T_-)\}$ and $i \in \{0, \dots, r\}$, let $\gamma_j^{(i)}$ and $\delta_k^{(i)}$ be the numbers of P_ρ -orbits of length p^i in rows j and k of T_+ and T_- , respectively. For each $i \in \{0, \dots, r\}$, let

$$\begin{aligned}\gamma^{(i)} &= (\gamma_1^{(i)}, \gamma_2^{(i)}, \dots), \\ \delta^{(i)} &= (\delta_1^{(i)}, \delta_2^{(i)}, \dots).\end{aligned}$$

Note that $|\gamma^{(i)}| + |\delta^{(i)}| = n_i$ for each i , and so $(\gamma^{(0)}, \gamma^{(1)}, \dots, \gamma^{(r)}|\delta^{(0)}, \delta^{(1)}, \dots, \delta^{(r)}) \in \Lambda((\alpha|\beta), \rho)$. We say that the $(\alpha|\beta)$ -tabloid $\{(T_+|T_-)\}$ is of ρ -type $(\gamma|\delta)$. For example, if $p = 3$, $n = 9$ and $\rho = (1^3, 3^2)$, so $P_\rho = \langle (4, 5, 6), (7, 8, 9) \rangle$, then the $((3, 3)|(3))$ -tabloid

$$\left\{ \begin{array}{|c|c|c|} \hline 1 & 2 & 3 \\ \hline 7 & 8 & 9 \\ \hline 4 & 5 & 6 \\ \hline \end{array} \right\}$$

has ρ -type $((3), (0, 1)|\emptyset, (1))$.

We denote the set of all $(\alpha|\beta)$ -tabloids of ρ -type $(\gamma|\delta)$ by $\Omega((\alpha|\beta), \rho)_{(\gamma|\delta)}$. Then the disjoint union

$$\Omega((\alpha|\beta), \rho) = \bigcup_{(\gamma|\delta) \in \Lambda((\alpha|\beta), \rho)} \Omega((\alpha|\beta), \rho)_{(\gamma|\delta)} \quad (3.2)$$

is an F -basis of $M(\alpha|\beta)(P_\rho)$. Thus, as F -vector spaces, we have

$$M(\alpha|\beta)(P_\rho) = F\Omega((\alpha|\beta), \rho) = \bigoplus_{(\gamma|\delta) \in \Lambda((\alpha|\beta), \rho)} F\Omega((\alpha|\beta), \rho)_{(\gamma|\delta)}. \quad (3.3)$$

It is clear that (3.3) is in fact a decomposition of FN_ρ -modules, since N_ρ permutes orbits of P_ρ of the same size as blocks for its action, and therefore preserves the ρ -type in its action on $(\alpha|\beta)$ -tabloids. Furthermore, P_ρ fixes all $(\alpha|\beta)$ -tabloids having a specified ρ -type. Therefore we obtain the following lemma.

Lemma 3.9. *Let $(\alpha|\beta) \in \mathcal{C}^2(n)$ and $\rho = (1^{n_0}, p^{n_1}, \dots, (p^r)^{n_r})$ be a partition of n . The Brauer quotient of $M(\alpha|\beta)$ with respect to the subgroup P_ρ has the following direct sum decomposition into $F[N_\rho/P_\rho]$ -modules:*

$$M(\alpha|\beta)(P_\rho) = \bigoplus_{(\gamma|\delta) \in \Lambda((\alpha|\beta), \rho)} F\Omega((\alpha|\beta), \rho)_{(\gamma|\delta)}.$$

By Lemma 3.9, to understand the Brauer quotient $M(\alpha|\beta)(P_\rho)$ of the signed Young permutation module $M(\alpha|\beta)$, it suffices to understand each of the $F[N_\rho/P_\rho]$ -modules $F\Omega((\alpha|\beta), \rho)_{(\gamma|\delta)}$.

Definition 3.10. Suppose that $(\alpha|\beta) \in \mathcal{C}^2(n)$ and that $\rho = (1^{n_0}, p^{n_1}, \dots, (p^r)^{n_r})$ is a partition of n . Let the orbits of P_ρ of size p^i be $\mathcal{O}_{i,1}, \dots, \mathcal{O}_{i,n_i}$. Let

$$\Theta : \Omega((\alpha|\beta), \rho) \rightarrow \bigcup_{(\gamma|\delta) \in \Lambda((\alpha|\beta), \rho)} \prod_{i=0}^r \Omega(\gamma^{(i)}|\delta^{(i)})$$

be the bijective function defined as follows. Suppose that $\{T\} \in \Omega(\alpha|\beta)$ is of ρ -type $(\gamma|\delta)$. For each $0 \leq i \leq r$, let $\{T_i\}$ be the $(\gamma^{(i)}|\delta^{(i)})$ -tabloid such that T_i is row standard, and row k of $(T_i)_+$ (respectively, $(T_i)_-$) contains j if and only if row k of T_+ (respectively, T_-) contains the orbit $\mathcal{O}_{i,j}$. Define $\Theta(\{T\}) = (\{T_i\})_{i=0,1,\dots,r}$.

We note that, by definition of P_ρ ,

$$\mathcal{O}_{i,j} = \left\{ (j-1)p^i + 1 + \sum_{\ell=0}^{i-1} n_\ell p^\ell, \dots, jp^i + \sum_{\ell=0}^{i-1} n_\ell p^\ell \right\},$$

for $i \in \{0, \dots, r\}$ and $j \in \{1, \dots, n_i\}$. Clearly, the bijection Θ in Definition 3.10 restricts to a bijection, also denoted Θ ,

$$\Theta : \Omega((\alpha|\beta), \rho)_{(\gamma|\delta)} \rightarrow \prod_{i=0}^r \Omega(\gamma^{(i)}|\delta^{(i)})$$

Since $|\Omega(\gamma^{(i)}|\delta^{(i)})| = \dim_F M(\gamma^{(i)}|\delta^{(i)}) = [\mathfrak{S}_{n_i} : (\mathfrak{S}_{\gamma^{(i)}} \times \mathfrak{S}_{\delta^{(i)}})]$, we obtain the following lemma.

Lemma 3.11. *Let $(\alpha|\beta) \in \mathcal{C}^2(n)$, $\rho = (1^{n_0}, p^{n_1}, \dots, (p^r)^{n_r})$ such that $|\rho| = n$, and let $(\gamma|\delta) \in \Lambda((\alpha|\beta), \rho)$. Set*

$$H = \prod_{i=0}^r N_{p^i} \wr (\mathfrak{S}_{\gamma^{(i)}} \times \mathfrak{S}_{\delta^{(i)}}) = \prod_{i=0}^r (N_{p^i} \wr \mathfrak{S}_{\gamma^{(i)}}) \times (N_{p^i} \wr \mathfrak{S}_{\delta^{(i)}}) \leq N_\rho.$$

Then $|\Omega((\alpha|\beta), \rho)_{(\gamma|\delta)}| = [N_\rho : H]$.

We have reached the main result of this section.

Proposition 3.12. *Suppose that $(\alpha|\beta) \in \mathcal{C}^2(n)$ and that $\rho = (1^{n_0}, p^{n_1}, \dots, (p^r)^{n_r}) \in \mathcal{C}(n)$. Regarded as an $F[N_\rho/P_\rho]$ -module, the Brauer quotient $M(\alpha|\beta)(P_\rho)$ of the signed Young permutation module $M(\alpha|\beta)$ with respect to P_ρ satisfies*

$$M(\alpha|\beta)(P_\rho) \cong \bigoplus_{(\gamma|\delta) \in \Lambda((\alpha|\beta), \rho)} W_1(\gamma^{(0)}|\delta^{(0)}) \boxtimes W_p(\gamma^{(1)}|\delta^{(1)}) \boxtimes \dots \boxtimes W_{p^r}(\gamma^{(r)}|\delta^{(r)}).$$

Proof. Recall that for each $(\lambda|\mu) \in \mathcal{C}^2(n)$, we have defined a row-standard $(\lambda|\mu)$ -tableau $T^{\lambda|\mu}$ immediately after Definition 3.1. Fix $(\gamma|\delta) \in \Lambda((\alpha|\beta), \rho)$ and let $Z = F\Omega((\alpha|\beta), \rho)_{(\gamma|\delta)}$.

By Lemma 3.9, it suffices to show that $Z \cong \boxtimes_{i=0}^r W_{p^i}(\gamma^{(i)}|\delta^{(i)})$ as FN_ρ -modules with P_ρ acting trivially, or equivalently, by (3.1), that

$$Z \cong \boxtimes_{i=0}^r \text{Ind}_{N_{p^i} \wr (\mathfrak{S}_{\gamma^{(i)}} \times \mathfrak{S}_{\delta^{(i)}})}^{N_{p^i} \wr \mathfrak{S}_{n_i}} (F(N_{p^i} \wr \mathfrak{S}_{\gamma^{(i)}}) \boxtimes \widehat{\text{sgn}}(N_{p^i})^{\otimes \delta^{(i)}}). \quad (3.4)$$

Let $\{S\} \in \Omega((\alpha|\beta), \rho)_{(\gamma|\delta)}$ be the unique $(\alpha|\beta)$ -tabloid such that

$$\Theta(\{S\}) = (T^{\gamma^{(0)}|\delta^{(0)}}, T^{\gamma^{(1)}|\delta^{(1)}}, \dots, T^{\gamma^{(r)}|\delta^{(r)}}) \in \prod_{i=0}^r \Omega(\gamma^{(i)}|\delta^{(i)}).$$

Using the N_ρ -action on Z , we observe that Z is a cyclic FN_ρ -module generated by $\{S\}$. Let X be the subspace of Z linearly spanned by $\{S\}$. By the definition of $\{S\}$, the subspace X is an FH -module where

$$H = \prod_{i=0}^r N_{p^i} \wr (\mathfrak{S}_{\gamma^{(i)}} \times \mathfrak{S}_{\delta^{(i)}}) = \prod_{i=0}^r ((N_{p^i} \wr \mathfrak{S}_{\gamma^{(i)}}) \times (N_{p^i} \wr \mathfrak{S}_{\delta^{(i)}})) \leq N_\rho,$$

and there is an isomorphism

$$X \cong (F(N_{p^0} \wr \mathfrak{S}_{\gamma^{(0)}}) \boxtimes \widehat{\text{sgn}}(N_{p^0})^{\otimes \delta^{(0)}}) \boxtimes \cdots \boxtimes (F(N_{p^r} \wr \mathfrak{S}_{\gamma^{(r)}}) \boxtimes \widehat{\text{sgn}}(N_{p^r})^{\otimes \delta^{(r)}})$$

of FH -modules. Since $\dim_F Z = [N_\rho : H] \dim_F X$ by Lemma 3.11, we have $Z \cong \text{Ind}_H^{N_\rho} X$ by the characterization of induced modules in [1, Section 8, Corollary 3]. Hence we obtain the isomorphism (3.4) as desired. \square

4. YOUNG MODULES AND SIGNED YOUNG MODULES

In this section we define Young modules and signed Young modules in the setting of the symmetric group and prove Theorem 1.1.

4.1. Vertices. As a first step we identify the possible vertices of summands of signed Young modules. Recall from Subsection 2.6 that P_k denotes a Sylow subgroup of \mathfrak{S}_k and, if ρ is a partition of n , then P_ρ denotes a Sylow subgroup of the Young subgroup \mathfrak{S}_ρ of \mathfrak{S}_n . We require the following lemma from [9]; a proof, slightly shorter than the one in [9], is included to make the article self-contained.

Lemma 4.1 (Erdmann [9, Lemma 1]). *Let G be a finite group and let M be a p -permutation FG -module. If P and \tilde{P} are p -subgroups of G such that $P < \tilde{P}$ and $\dim_F M(P) = \dim_F M(\tilde{P})$ then no indecomposable summand of M has vertex P .*

Proof. Suppose, for a contradiction, that U is such a summand. Let $M = U \oplus V$ where V is a complementary FG -module. By Corollary 2.4, we have $U(P) \neq 0$ and $U(\tilde{P}) = 0$. Thus

$$M(\tilde{P}) = U(\tilde{P}) \oplus V(\tilde{P}) = V(\tilde{P}), \quad \text{and} \quad M(P) = U(P) \oplus V(P).$$

This is a contradiction, since taking a p -permutation basis for V and applying Corollary 2.4 shows that $\dim_F V(P) \geq \dim_F V(\tilde{P})$. \square

Proposition 4.2. *Let $(\alpha|\beta) \in \mathcal{C}^2(n)$. If P is a vertex of an indecomposable summand of $M(\alpha|\beta)$ then there exists $\rho = (1^{n_0}, p^{n_1}, \dots, (p^r)^{n_r}) \in \mathcal{C}(n)$ such that P is conjugate in \mathfrak{S}_n to P_ρ .*

Proof. Let H be the Young subgroup of \mathfrak{S}_n having the same orbits as P on $\{1, \dots, n\}$ and let \tilde{P} be a Sylow p -subgroup of H . Note that \tilde{P} has the same orbits on $\{1, \dots, n\}$ as P : suppose that each subgroup has exactly n_i orbits of size p^i for each $i \in \{0, \dots, r\}$, so \tilde{P} is conjugate in \mathfrak{S}_n to P_ρ . It suffices to prove that $P = \tilde{P}$.

Let $\{T\}$ be an $(\alpha|\beta)$ -tabloid fixed by P . As remarked following Corollary 3.4, each row of T is a union of orbits of P . Therefore each row is a union of orbits of \tilde{P} , and so if $g \in \tilde{P}$ then $g\{T\} = \pm\{T\}$. Since g has p -power order, we see that $g\{T\} = \{T\}$. It now follows from Corollary 3.4 that $\dim_F M(\alpha|\beta)(P) = \dim_F M(\alpha|\beta)(\tilde{P})$. By Lemma 4.1 we have $P = \tilde{P}$, as required. \square

Combining Proposition 3.12 and Proposition 4.2, we see that the Broué correspondents of the non-projective indecomposable summands of $M(\alpha|\beta)$ are certain outer tensor products of the projective indecomposable summands of the $F[(N_{p^i}/P_{p^i}) \wr \mathfrak{S}_m]$ -modules $W_{p^i}(\gamma|\delta)$ in Definition 3.6. In fact, it is most convenient to factor out a further subgroup that acts trivially, and consider projective summands of the $F[C_2 \wr \mathfrak{S}_m]$ -modules $\overline{W}_{p^i}(\gamma|\delta)$.

4.2. Projective summands of $\overline{W}_k(\gamma|\delta)$. Fix $k \in \mathbb{N}$ and $m_1, m_2 \in \mathbb{N}_0$. Let $m = m_1 + m_2$. Recall from Subsection 2.5 that if $\alpha \in \mathcal{RP}(n)$, that is, α is a p -restricted partition of n , then P^α denotes the projective cover of the simple $F\mathfrak{S}_n$ -module D_α .

We remind the reader that the bifunctor \mathcal{F} was defined just before Definition 3.6.

Definition 4.3. Let $(\alpha|\beta) \in \mathcal{RP}^2(m)$. We define $\overline{Q}(\alpha|\beta) = \mathcal{F}(P^\alpha|P^\beta)$.

Example 4.10 gives an example of these modules. Note that each tensor factor is projective, so each $\overline{Q}(\alpha|\beta)$ is projective.

Lemma 4.4. *The $F[C_2 \wr \mathfrak{S}_m]$ -modules $\mathcal{F}(D_\alpha|D_\beta)$ for $(\alpha|\beta) \in \mathcal{RP}^2(m)$ form a complete set of non-isomorphic simple $F[C_2 \wr \mathfrak{S}_m]$ -modules. Moreover, the $F[C_2 \wr \mathfrak{S}_m]$ -module $\overline{Q}(\alpha|\beta)$ is the projective cover of $\mathcal{F}(D_\alpha|D_\beta)$ and the modules $\overline{Q}(\alpha|\beta)$ for $(\alpha|\beta) \in \mathcal{RP}^2(m)$ form a complete set of non-isomorphic indecomposable projective modules for $F[C_2 \wr \mathfrak{S}_m]$.*

Proof. The first claim follows from the construction of simple modules for wreath products stated in Theorem 4.34 of [20]. For the second, note that by functoriality, there is a surjection $\overline{Q}(\alpha|\beta) = \mathcal{F}(P^\alpha|P^\beta) \rightarrow \mathcal{F}(D_\alpha|D_\beta)$. Hence the projective $F[C_2 \wr \mathfrak{S}_m]$ -module $\overline{Q}(\alpha|\beta)$ has the projective cover of $\mathcal{F}(D_\alpha|D_\beta)$ as a summand. Since the inertial group of

$$\mathcal{F}(P^\alpha|P^\beta) = \text{Inf}_{\mathfrak{S}_{m_1}}^{C_2 \wr \mathfrak{S}_{m_1}}(P^\alpha) \boxtimes (\text{Inf}_{\mathfrak{S}_{m_2}}^{C_2 \wr \mathfrak{S}_{m_2}}(P^\beta) \otimes \widehat{\text{sgn}(2)}^{\otimes m_2})$$

is $(C_2 \wr \mathfrak{S}_{m_1}) \times (C_2 \wr \mathfrak{S}_{m_2})$, it follows from [2, Proposition 3.13.2] that $\overline{Q}(\alpha|\beta)$ is indecomposable. Therefore $\overline{Q}(\alpha|\beta)$ is the projective cover of $\mathcal{F}(D_\alpha|D_\beta)$. \square

Let G be a finite group. By Section 3.11 in [2], we may associate a character to a p -permutation FG -module M by taking a p -modular system (K, \mathcal{O}, F) compatible with F

and an $\mathcal{O}G$ -module $M_{\mathcal{O}}$ whose p -modular reduction is M . The *ordinary character* of M is then the character of the KG -module $K \otimes_{\mathcal{O}} M_{\mathcal{O}}$. If M is projective and indecomposable, the ordinary character of M may equivalently be defined by Brauer reciprocity (see for instance [26, Section 15.4]).

Proposition 4.5. *Let $(\gamma|\delta) \in \mathcal{P}^2(m)$ where $|\gamma| = m_1$ and $|\delta| = m_2$. Each indecomposable projective summand of $\overline{W}_k(\gamma|\delta)$ is isomorphic to some $\overline{Q}(\alpha|\beta)$, where $(\alpha|\beta) \in \mathcal{R}\mathcal{P}^2(m)$ satisfies*

- (i) $|\alpha| = m_1$ and $|\beta| = m_2$;
- (ii) $\alpha \succeq \gamma$ and $\beta \succeq \delta$.

Proof. By Lemma 4.4, each indecomposable projective summand of $\overline{W}_k(\gamma|\delta)$ is isomorphic to some $\overline{Q}(\alpha|\beta)$. By the ‘wedge’ shape of the decomposition matrix of \mathfrak{S}_n with columns labelled by p -restricted partitions (see for instance [21, Theorem 5.2]) and Brauer reciprocity, the ordinary character of P^α contains the irreducible character χ^α exactly once. Hence the ordinary character of $\overline{Q}(\alpha|\beta)$ contains the character

$$\chi^{(\alpha|\beta)} = \text{Ind}_{C_2 \wr (\mathfrak{S}_{m_1} \times \mathfrak{S}_{m_2})}^{C_2 \wr \mathfrak{S}_m} \left(\text{Inf}_{\mathfrak{S}_{m_1}}^{C_2 \wr \mathfrak{S}_{m_1}}(\chi^\alpha) \times (\text{Inf}_{\mathfrak{S}_{m_2}}^{C_2 \wr \mathfrak{S}_{m_2}}(\chi^\beta) \otimes \widehat{\text{sgn}(2)}^{\otimes m_2}) \right)$$

defined in Subsection 2.5 exactly once.

We now consider when the ordinary character of $\overline{W}_k(\gamma|\delta)$ contains $\chi^{(\alpha|\beta)}$. The restriction of $\overline{W}_k(\gamma|\delta)$ to the base group in the wreath product $C_2 \wr \mathfrak{S}_m$ is a direct sum of 1-dimensional submodules. In each such submodule, m_1 of the factors in the product C_2^m act trivially and m_2 of the factors act as $\text{sgn}(2)$. It follows by basic Clifford theory that the ordinary character of $\overline{W}_k(\gamma|\delta)$ contains the character $\chi^{(\alpha|\beta)}$ only if $|\alpha| = m_1$ and $|\beta| = m_2$. By Young’s rule (see for instance [18, Theorem 13.13]), the ordinary character of M^γ contains χ^α only if $\alpha \succeq \gamma$, and similarly the ordinary character of M^δ contains χ^β only if $\beta \succeq \delta$. It follows that if $\overline{Q}(\alpha|\beta)$ is a summand of $\overline{W}_k(\gamma|\delta)$ then $\alpha \in \mathcal{P}(m_1)$, $\beta \in \mathcal{P}(m_2)$, $\alpha \succeq \gamma$ and $\beta \succeq \delta$. \square

4.3. Definition of signed Young modules. We define signed Young modules as the Broué correspondents of tensor products of suitable inflations of the modules $\overline{Q}(\alpha|\beta)$. To make this precise, we need the three further families of modules defined below: their definition follows the same pattern as the p -permutation modules $V_k(\gamma|\delta)$, $W_k(\gamma|\delta)$ and $\overline{W}_k(\gamma|\delta)$ in Definition 3.6.

Definition 4.6. Let $k \in \mathbb{N}$, let $m \in \mathbb{N}_0$, and let $(\alpha|\beta) \in \mathcal{R}\mathcal{P}^2(m)$. Let $m_1 = |\alpha|$ and $m_2 = |\beta|$. The $F[\mathfrak{S}_k \wr \mathfrak{S}_m]$ -module $R_k(\alpha|\beta)$ is defined by

$$R_k(\alpha|\beta) = \mathcal{F}(P^\alpha | P^\beta).$$

By convention, $R_k(\emptyset|\beta) = \text{Inf}_{\mathfrak{S}_{m_2}}^{\mathfrak{S}_k \wr \mathfrak{S}_{m_2}}(P^\beta) \otimes \widehat{\text{sgn}(k)}^{\otimes m_2}$, and similarly for $R_k(\alpha|\emptyset)$. Furthermore if $m = 0$, then $R_k(\emptyset|\emptyset)$ is the trivial $F\mathfrak{S}_0$ -module. If $k = 1$ then we identify $\mathfrak{S}_k \wr \mathfrak{S}_m$ with \mathfrak{S}_m and get

$$R_1(\alpha|\beta) = \text{Ind}_{\mathfrak{S}_{m_1} \times \mathfrak{S}_{m_2}}^{\mathfrak{S}_m} (P^\alpha \boxtimes (P^\beta \otimes \text{sgn}(m_2))).$$

Recall from Section 2.6 that P_k is a fixed Sylow p -subgroup of \mathfrak{S}_k and that $N_k = N_{\mathfrak{S}_k}(P_k)$.

Definition 4.7. Let $k \in \mathbb{N}$, let $m \in \mathbb{N}_0$, and let $(\alpha|\beta) \in \mathcal{RP}^2(m)$. Let $Q_k(\alpha|\beta)$ be the $F[(N_k/P_k) \wr \mathfrak{S}_m]$ -module defined by

$$Q_k(\alpha|\beta) = \text{Res}_{N_k \wr \mathfrak{S}_m}^{\mathfrak{S}_k \wr \mathfrak{S}_m} R_k(\alpha|\beta)$$

considered as an $F[(N_k/P_k) \wr \mathfrak{S}_m]$ -module via the canonical surjection $N_k \wr \mathfrak{S}_m \rightarrow (N_k \wr \mathfrak{S}_m)/(P_k)^m \cong (N_k/P_k) \wr \mathfrak{S}_m$.

Again if $k = 1$ we identify $N_1 \wr \mathfrak{S}_m$ with \mathfrak{S}_m and we have $Q_1(\alpha|\beta) = R_1(\alpha|\beta)$. Since $(\mathfrak{A}_k)^m$ acts trivially on $R_k(\alpha|\beta)$ we see that $(N_{\mathfrak{A}_k}(P_k)/P_k)^m$ acts trivially on $Q_k(\alpha|\beta)$. It is clear that

$$Q_k(\alpha|\beta) \cong \text{Ind}_{N_k \wr (\mathfrak{S}_{m_1} \times \mathfrak{S}_{m_2})}^{N_k \wr \mathfrak{S}_m} \left(\text{Inf}_{\mathfrak{S}_{m_1}}^{N_k \wr \mathfrak{S}_{m_1}}(P^\alpha) \boxtimes \left(\text{Inf}_{\mathfrak{S}_{m_2}}^{N_k \wr \mathfrak{S}_{m_2}}(P^\beta) \otimes \widehat{\text{sgn}(N_k)}^{\otimes m_2} \right) \right), \quad (4.1)$$

again regarded as an $F[(N_k/P_k) \wr \mathfrak{S}_m]$ -module by this canonical surjection.

Definition 4.8. Let $k \in \mathbb{N}$, let $m \in \mathbb{N}_0$, and let $(\alpha|\beta) \in \mathcal{RP}^2(m)$. For $k \geq 2$, let $\overline{Q}_k(\alpha|\beta)$ be the $F[C_2 \wr \mathfrak{S}_m]$ -module obtained from $Q_k(\alpha|\beta)$ via the canonical surjection

$$(N_k/P_k) \wr \mathfrak{S}_m \rightarrow ((N_k/P_k) \wr \mathfrak{S}_m)/(N_{\mathfrak{A}_k}(P_k)/P_k)^m \cong C_2 \wr \mathfrak{S}_m.$$

We define the $F[C_2 \wr \mathfrak{S}_m]$ -module $\overline{Q}_1(\alpha|\beta)$ by

$$\overline{Q}_1(\alpha|\beta) = \text{Inf}_{\mathfrak{S}_m}^{C_2 \wr \mathfrak{S}_m} Q_1(\alpha|\beta).$$

The following lemma justifies the notation $\overline{Q}_k(\alpha|\beta)$ for the projective modules just defined.

Lemma 4.9. Let $k \geq 2$ and let $(\alpha|\beta) \in \mathcal{RP}^2(m)$ where $m \in \mathbb{N}_0$. Then

$$\overline{Q}_k(\alpha|\beta) \cong \overline{Q}(\alpha|\beta) \cong R_2(\alpha|\beta)$$

as $F[C_2 \wr \mathfrak{S}_m]$ -modules.

Proof. The first isomorphism is clear from the definitions and the second follows as in Lemma 3.7. \square

We pause to give a small example showing the exceptional behaviour when $k = 1$.

Example 4.10. Let $p = 3$, and let $k \geq 2$. Let $\varepsilon = \text{Inf}_{\mathfrak{S}_3}^{\mathfrak{S}_k \wr \mathfrak{S}_3}(\text{sgn}(3))$. There are four mutually non-isomorphic 1-dimensional simple $F[\mathfrak{S}_k \wr \mathfrak{S}_3]$ -modules, namely

$$\widehat{F(k)}^{\otimes 3}, \widehat{\text{sgn}(k)}^{\otimes 3}, \widehat{F(k)}^{\otimes 3} \otimes \varepsilon, \widehat{\text{sgn}(k)}^{\otimes 3} \otimes \varepsilon,$$

where the trivial module appears as $\widehat{F(k)}^{\otimes 3} \otimes \varepsilon$. The corresponding modules $R_k(\alpha|\beta)$ are $R_k((1, 1, 1)|\emptyset)$, $R_k(\emptyset|(2, 1))$, $R_k((2, 1)|\emptyset)$, $R_k(\emptyset|(1, 1, 1))$, respectively. Quotienting out by the trivial action of \mathfrak{A}_k , the corresponding modules $\overline{Q}(\alpha|\beta)$ for $F[C_2 \wr \mathfrak{S}_3]$ are precisely the projective covers of the four one-dimensional simple modules for $F[C_2 \wr \mathfrak{S}_3]$. For example, since the base group acts trivially, and the top group acts by sign, the projective cover of

$\widehat{F(2)}^{\otimes 3}$ is $\text{Inf}_{S_3}^{C_2 \wr S_3} P^{(1,1,1)}$, namely $\overline{Q}((1, 1, 1)|\emptyset)$. The four remaining simple modules for $F[C_2 \wr S_3]$, each projective; by Lemma 4.4, they are isomorphic to the modules $\overline{Q}(\alpha|\beta)$ where both α and β are non-empty. By contrast, when $k = 1$, identifying $S_1 \wr S_3$ with S_3 as described after Definition 4.6, we have $\overline{Q}_1((1, 1, 1)|\emptyset) \cong \overline{Q}_1(\emptyset|(2, 1)) \cong P^{(1,1,1)} \cong M^{(2,1)} \otimes \text{sgn}$ and $\overline{Q}_1((2, 1)|\emptyset) \cong \overline{Q}_1(\emptyset|(1, 1, 1)) \cong P^{(2,1)} \cong M^{(2,1)}$.

We are finally ready to define signed Young modules.

Definition 4.11. Let $(\lambda|p\mu) \in \mathcal{R}\mathcal{P}^2(n)$. Let $\lambda = \sum_{i \geq 0} p^i \lambda(i)$ and $\mu = \sum_{i \geq 0} p^i \mu(i)$ be the p -adic expansions of λ and μ , as defined in (2.1). Let $n_0 = |\lambda(0)|$ and let $n_i = |\lambda(i)| + |\mu(i-1)|$ for each $i \in \mathbb{N}$. Let r be maximal such that $n_r \neq 0$ and let $\rho = (1^{n_0}, p^{n_1}, \dots, (p^r)^{n_r})$. We define the *signed Young module* $Y(\lambda|p\mu)$ to be the unique (up to isomorphism) $F\mathfrak{S}_n$ -module V such that

$$V(P_\rho) \cong Q_1(\lambda(0)|\emptyset) \boxtimes Q_p(\lambda(1)|\mu(0)) \boxtimes \cdots \boxtimes Q_{p^r}(\lambda(r)|\mu(r-1)).$$

We define a *Young module* to be a signed Young module of the form $Y(\lambda|\emptyset)$.

The isomorphism above is an isomorphism of projective $F[\mathfrak{S}_{n_0} \times ((N_p/P_p) \wr \mathfrak{S}_{n_1}) \times \cdots \times ((N_{p^r}/P_{p^r}) \wr \mathfrak{S}_{n_r})]$ -modules. Observe that P_ρ is trivial if and only if λ is p -restricted and $\mu = \emptyset$; in this case $Q_1(\lambda(0)|\emptyset)$ is regarded as a \mathfrak{S}_n -module by identifying $N_1 \wr \mathfrak{S}_n$ with \mathfrak{S}_n , and since $\lambda = \lambda(0)$ we have $Y(\lambda|\emptyset) = Q_1(\lambda|\emptyset) = P^\lambda$.

The following proposition gives part of Theorem 1.1(i).

Proposition 4.12.

- (i) If $(\alpha|\beta) \in \mathcal{P}^2(n)$ then $M(\alpha|\beta)$ is a direct sum of signed Young modules.
- (ii) If $\alpha \in \mathcal{P}(n)$ then M^α is a direct sum of Young modules.

Proof. Let $(\alpha|\beta) \in \mathcal{P}^2(n)$ and let V be an indecomposable summand of $M(\alpha|\beta)$. By Proposition 4.2 there exists $\rho = (1^{m_0}, p^{m_1}, \dots, (p^r)^{m_r}) \in \mathcal{C}(n)$ such that P_ρ is a vertex of V . Recall that $N_\rho/P_\rho \cong \mathfrak{S}_{n_0} \times ((N_p/P_p) \wr \mathfrak{S}_{n_1}) \times \cdots \times ((N_{p^r}/P_{p^r}) \wr \mathfrak{S}_{n_r})$. By Proposition 3.12, there exists $(\gamma|\delta) \in \Lambda((\alpha|\beta), \rho)$ such that the projective $F[N_\rho/P_\rho]$ -module $V(P_\rho)$ is a direct summand of

$$W_1(\gamma^{(0)}|\delta^{(0)}) \boxtimes W_p(\gamma^{(1)}|\delta^{(1)}) \boxtimes \cdots \boxtimes W_{p^r}(\gamma^{(r)}|\delta^{(r)}).$$

By Lemmas 4.4 and 4.9 there exist partitions $\lambda(0), \dots, \lambda(r)$ and $\mu(0), \dots, \mu(r-1)$ such that

$$V(P_\rho) = Q_1(\lambda(0)) \boxtimes Q_p(\lambda(1)|\mu(0)) \boxtimes \cdots \boxtimes Q_{p^r}(\lambda(r)|\mu(r-1)).$$

By Theorem 2.5, $V \cong Y(\lambda|p\mu)$ where $\lambda = \sum_{i=0}^r p^i \lambda(i)$ and $\mu = \sum_{i=0}^{r-1} p^i \mu(i)$. This proves part (i). For part (ii), observe that if $\beta = \emptyset$ then we have $\delta^{(i)} = \emptyset$ for each i , and so $\mu(i) = \emptyset$ for each i . \square

4.4. Column symmetrization of $(\alpha|\beta)$ -tableaux. To deal with the projective summands of signed Young permutation modules we require the following corollary of the key lemma used by James to prove his Submodule Theorem in [18]. Given a tableau \mathfrak{t} with entries from a set \mathcal{O} , let $C_{\mathfrak{t}} \leq \mathfrak{S}_{\mathcal{O}}$ be the group of permutations which fix the columns of \mathfrak{t} setwise. Set $\kappa_{\mathfrak{t}} = \sum_{g \in C_{\mathfrak{t}}} \text{sgn}(g)g$.

Proposition 4.13. *Let $\lambda \in \mathcal{P}(n)$ and let \mathfrak{t} be a λ -tableau. In any direct sum decomposition of M^{λ} into indecomposable modules there is a unique summand U^{λ} such that $\kappa_{\mathfrak{t}}U^{\lambda} \neq 0$. Moreover if $\alpha \in \mathcal{P}(n)$ then $\kappa_{\mathfrak{t}}U^{\alpha} = 0$ unless $\lambda \trianglerighteq \alpha$.*

Proof. This follows immediately from [18, Lemma 4.6]. \square

By the Krull–Schmidt Theorem, the U^{λ} are well-defined up to isomorphism. It is clear that $U^{\alpha} \cong U^{\beta}$ if and only if $\alpha = \beta$.

We also need the following generalization of part of James' lemma.

Lemma 4.14. *Let $(\alpha|\beta) \in \mathcal{C}^2(n)$ and let $\mathbf{T} = (\mathbf{T}_+|\mathbf{T}_-)$ be an $(\alpha|\beta)$ -tableau. Let $\lambda \in \mathcal{P}(n)$ and let \mathfrak{t} be a λ -tableau. If $\kappa_{\mathfrak{t}}\{\mathbf{T}\} \neq 0$ then $(\lambda|\emptyset) \trianglerighteq (\alpha|\beta)$.*

Proof. Let \mathcal{O} be the set of entries of \mathbf{T}_+ . Let $H = \mathfrak{S}_{\mathcal{O}} \cap \mathfrak{S}_{\lambda'}$ and let $\mathcal{O}_1, \dots, \mathcal{O}_s$ be the orbits of H on \mathcal{O} , ordered so that $|\mathcal{O}_1| \geq \dots \geq |\mathcal{O}_s|$. Let $\nu = (|\mathcal{O}_1|, \dots, |\mathcal{O}_s|)' \in \mathcal{P}(|\alpha|)$. The j th largest orbit of H has size at most λ'_j . Therefore $\nu'_j \leq \lambda'_j$ for each $j \in \{1, \dots, s\}$, and so ν is a subpartition of λ . It immediately follows that

$$\sum_{i=1}^k \lambda_i \geq \sum_{i=1}^k \nu_i \quad (4.2)$$

for all $k \in \mathbb{N}$. (By our standing convention, $\nu_i = 0$ if $i > \ell(\nu)$).

Let \mathfrak{t}^* be a ν -tableau having the entries of \mathcal{O}_j in its j th column. Observe that $C_{\mathfrak{t}^*} \leq C_{\mathfrak{t}}$. Choose $g_1, \dots, g_s \in C_{\mathfrak{t}}$ such that $C_{\mathfrak{t}} = g_1 C_{\mathfrak{t}^*} \cup \dots \cup g_s C_{\mathfrak{t}^*}$ where the union is disjoint. We have

$$\kappa_{\mathfrak{t}} = (\text{sgn}(g_1)g_1 + \dots + \text{sgn}(g_s)g_s)\kappa_{\mathfrak{t}^*}.$$

Since $\kappa_{\mathfrak{t}}\{\mathbf{T}\} \neq 0$ we have $\kappa_{\mathfrak{t}^*}\{\mathbf{T}\} \neq 0$. Since $C_{\mathfrak{t}^*}$ fixes the entries in \mathbf{T}_- , it follows that $\kappa_{\mathfrak{t}^*}\{\mathbf{T}_+\} \neq 0$. The argument used to prove Lemma 4.6 in [18] now shows that any two entries in the same row of $\{\mathbf{T}_+\}$ lie in different columns of \mathfrak{t}^* , and so $\nu \trianglerighteq \alpha$. Hence, by (4.2), we have $(\lambda|\emptyset) \trianglerighteq (\alpha|\beta)$, as required. \square

4.5. Proof of Theorem 1.1. For convenience we repeat the statement of this theorem below.

Theorem 1.1 (Donkin [7]). *There exist indecomposable $F\mathfrak{S}_n$ -modules $Y(\lambda|p\mu)$ for $(\lambda|p\mu) \in \mathcal{P}^2(n)$ with the following properties:*

(i) *if $(\alpha|\beta) \in \mathcal{P}^2(n)$ then $M(\alpha|\beta)$ is isomorphic to a direct sum of modules $Y(\lambda|p\mu)$ for $(\lambda|p\mu) \in \mathcal{P}^2(n)$ such that $(\lambda|p\mu) \trianglerighteq (\alpha|\beta)$,*

(ii) $[M(\lambda|p\mu) : Y(\lambda|p\mu)] = 1$,

(iii) if $\lambda = \sum_{i=0}^r p^i \lambda(i)$ and $\mu = \sum_{i=0}^{r-1} p^i \mu(i)$ are the p -adic expansions of λ and μ , as defined in (2.1), then $Y(\lambda|p\mu)$ has as a vertex a Sylow p -subgroup of the Young subgroup \mathfrak{S}_ρ , where ρ is the partition of n having exactly $|\lambda(i)| + |\mu(i-1)|$ parts of size p^i for each $i \in \{0, \dots, r\}$.

We shall prove the theorem by showing that parts (i), (ii) and (iii) of Theorem 1.1 hold when $Y(\lambda|p\mu)$ is as defined in Definition 4.11. In fact part (iii) holds by definition, so we may concentrate on parts (i) and (ii).

Proof of Theorem 1.1. We work by induction on $n \in \mathbb{N}_0$. If $n < p$ then $F\mathfrak{S}_n$ is semisimple and the modules $Y(\lambda|\emptyset)$ for $\lambda \in \mathcal{P}(n)$ form a complete set of simple $F\mathfrak{S}_n$ -modules. Hence parts (i) and (ii) follow from Proposition 4.5. Now let $n \geq p$.

We first deal with non-projective summands. Let $(\lambda|p\mu) \in \mathcal{P}^2(n)$ and suppose that either λ is not p -restricted or $\mu \neq \emptyset$. Let $n_0 = |\lambda(0)|$ and let $n_i = |\lambda(i)| + |\mu(i-1)|$ for $i \in \mathbb{N}$. Let $\rho = (1^{n_0}, p^{n_1}, \dots, (p^r)^{n_r})$.

By Theorem 2.5 and Proposition 3.12, $[M(\alpha|\beta) : Y(\lambda|p\mu)]$ is equal to the sum of the products

$$[W_1(\gamma^{(0)}|\delta^{(0)}) : P^{\lambda(0)}] [W_p(\gamma^{(1)}|\delta^{(1)}) : Q_p(\lambda(1)|\mu(0))] \cdots [W_{p^r}(\gamma^{(r)}|\delta^{(r)}) : Q_{p^r}(\lambda(r)|\mu(r-1))]$$

over all $(\gamma|\delta) \in \Lambda((\alpha|\beta), \rho)$. Suppose the product is non-zero for $(\gamma|\delta) \in \Lambda((\alpha|\beta), \rho)$. Then $P^{\lambda(0)}$ is a direct summand of $W_1(\gamma^{(0)}|\delta^{(0)}) \cong M(\gamma^{(0)}|\delta^{(0)})$. Since $P^{\lambda(0)} = Y(\lambda(0)|\emptyset)$, it follows from the inductive hypothesis that $(\lambda(0)|\emptyset) \supseteq (\gamma^{(0)}|\delta^{(0)})$. Similarly $Q_{p^i}(\lambda(i)|\mu(i-1))$ is a direct summand of $W_{p^i}(\gamma^{(i)}|\delta^{(i)})$ for each $i \in \{1, \dots, r\}$. By Proposition 4.5, we have $\lambda(i) \supseteq \gamma^{(i)}$ and $\mu(i-1) \supseteq \delta^{(i)}$ for each such i . Hence

$$\lambda - \lambda(0) = \sum_{i=1}^r p^i \lambda(i) \supseteq \sum_{i=1}^r p^i \gamma^{(i)} = \alpha - \gamma^{(0)} \quad (4.3)$$

and

$$p\mu = \sum_{i=1}^r p^i \mu(i-1) \supseteq \sum_{i=1}^r p^i \delta^{(i)} = \beta - \delta^{(0)}. \quad (4.4)$$

Hence $\lambda \supseteq \alpha - \gamma^{(0)} + \lambda(0) \supseteq \alpha$ and

$$|\lambda| + \sum_{i=1}^j p\mu_i = |\alpha| + |\delta^{(0)}| + \sum_{i=1}^j p\mu_i \geq |\alpha| + |\delta^{(0)}| + \sum_{i=1}^j (\beta - \delta^{(0)})_i \geq |\alpha| + \sum_{i=1}^j \beta_i$$

for all $j \in \mathbb{N}$. Therefore $(\lambda|p\mu) \supseteq (\alpha|\beta)$. By Proposition 4.12, every summand of $M(\alpha|\beta)$ is isomorphic to some $Y(\lambda|p\mu)$, so this proves part (i) in the non-projective case. If $(\alpha|\beta) = (\lambda|p\mu)$ then, by divisibility considerations, $\gamma^{(0)} = \lambda(0)$ and $\delta^{(0)} = \emptyset$. Moreover, equality holds in both (4.3) and (4.4), so we have $\gamma^{(i)} = \lambda(i)$ and $\delta^{(i)} = \mu(i-1)$ for each $i \in \{1, \dots, r\}$. Conversely, if γ and δ are defined in this way, then the product is 1. This proves part (ii) in the non-projective case.

We now deal with the projective summands. By Proposition 4.12, if $\alpha \in \mathcal{P}(n)$ then M^α is a direct sum of modules $Y(\lambda|\emptyset)$ for $\lambda \in \mathcal{P}(n)$. The argument so far shows that if α is

not p -restricted then $Y(\alpha|\emptyset)$ is a summand of M^α , and $Y(\alpha|\emptyset)$ is a summand of M^γ only if $\alpha \supseteq \gamma$. Therefore, inductively working down the dominance order on partitions, we see that, for each such α , the submodule U^α in Proposition 4.13 is $Y(\alpha|\emptyset)$. By counting, the remaining U^α for $\alpha \in \mathcal{RP}(n)$ are the modules $Y(\lambda|\emptyset)$ for $\lambda \in \mathcal{RP}(n)$. Again working inductively down the dominance order of partitions, it follows from Proposition 4.5 that $U^\alpha = Y(\alpha|\emptyset)$ for each $\alpha \in \mathcal{RP}(n)$. This proves part (i) in the projective case when $\beta = \emptyset$, and also proves part (ii) in the projective case.

Finally suppose that λ is p -restricted and $Y(\lambda|\emptyset)$ is a direct summand of $M(\alpha|\beta)$. Let \mathfrak{t} be a λ -tableau. By Proposition 4.12, we have $\kappa_{\mathfrak{t}}M(\alpha|\beta) \neq 0$. Hence there exists an $(\alpha|\beta)$ -tabloid $\{\mathfrak{T}\}$ such that $\kappa_{\mathfrak{t}\{\mathfrak{T}\}} \neq 0$. By Lemma 4.14 we have $(\lambda|\emptyset) \supseteq (\alpha|\beta)$. This completes the proof of part (i) in the projective case. \square

5. APPLICATIONS OF THEOREM 1.1

5.1. Equivalent definitions. We observed in the introduction that since signed Young modules are characterized by Theorem 1.1, our definition of signed Young modules agrees with Donkin's in [7]. Similarly Theorem 1.1 characterizes the Young module $Y(\lambda|\emptyset)$ as the unique summand of M^λ appearing in M^μ only if $\lambda \supseteq \mu$. By [19, Theorem 3.1(i)], James' Young modules admit the same characterisation. The two definitions therefore agree. In [10], Erdmann and Schroll consider Young modules for finite general linear groups. Adapting their proof to symmetric groups (this is mentioned as a possibility in [10], as a way to correct [9]), their definition of the Young modules uses the characterization in Proposition 4.12. Our proof of Theorem 1.1 shows these definitions agree; of course this also follows from the alternative characterization just mentioned.

Remark 5.1.

- (1) The counting argument used in our proof of the projective case of Theorem 1.1 is motivated by similar counting arguments used in [10]; the authors of [10] thank Burkhard Külshammer for suggesting this approach.
- (2) We have assumed throughout that F has odd prime characteristic p . It is possible to construct Young modules when $p = 2$ and to prove the analogue of Theorem 1.1 by adapting (and simplifying) the approach herein.
- (3) The analogue of signed Young modules for the finite general linear group $\mathrm{GL}_n(\mathbb{F}_q)$ are the linear source modules induced from powers of the determinant representation of parabolic subgroups of $\mathrm{GL}_n(\mathbb{F}_q)$. These modules seem worthy of study, especially given the difficulty of working directly with Specht modules for $\mathrm{GL}_n(\mathbb{F}_q)$.

5.2. Klyachko's formula and other applications. The following corollary generalizes Klyachko's formula to signed Young modules. It is proved in the first step of our proof of Theorem 1.1; alternatively it follows from this theorem by taking Broué correspondents.

Corollary 5.2. *If $(\alpha|\beta)$ and $(\lambda|p\mu) \in \mathcal{P}^2(n)$ then*

$$\begin{aligned} & [M(\alpha|\beta) : Y(\lambda|p\mu)] \\ &= \sum_{(\gamma|\delta) \in \Lambda((\alpha|\beta), \rho)} [W_1(\gamma^{(0)}|\delta^{(0)}) : Y(\lambda(0)|\emptyset)] \prod_{i=1}^r [W_{p^i}(\gamma^{(i)}|\delta^{(i)}) : Q_{p^i}(\lambda(i)|\mu(i-1))]. \end{aligned}$$

We remark that the reduction formula for signed p Kostka numbers in Corollary 5.2 has previously been obtained by Danz, the first and the second authors in [5].

The proof of the following lemma is very easy and is left to the reader. Recall that the notation \bullet for the concatenation of two compositions was defined in 2.3.

Lemma 5.3. *Let*

$$\begin{aligned} \rho &= (1^{m_0}, p^{m_1}, \dots, (p^r)^{m_r}), \\ \gamma &= (1^{n_0}, p^{n_1}, \dots, (p^s)^{n_s}), \end{aligned}$$

be partitions of m and n respectively, and let $k > r$. Then

$$\begin{aligned} P_{\rho \bullet p^k \gamma} &= P_\rho \times P_{p^k \gamma}, \\ N_{\mathfrak{S}_{m+p^k n}}(P_\rho \times P_{p^k \gamma}) &= N_{\mathfrak{S}_m}(P_\rho) \times N_{\mathfrak{S}_{p^k n}}(P_{p^k \gamma}), \\ N_{\rho \bullet p^k \gamma} / P_{\rho \bullet p^k \gamma} &= (N_\rho / P_\rho) \times (N_{p^k \gamma} / P_{p^k \gamma}). \end{aligned}$$

Let $(\lambda|p\mu) \in \mathcal{P}^2(n)$. Suppose that the p -adic expansions of λ and μ are $\lambda = \sum_{i \geq 0} p^i \lambda(i)$ and $\mu = \sum_{i \geq 0} p^i \mu(i)$, respectively. Let $\mu(-1) = \emptyset$. If r is maximal such that $|\lambda(r)| + |\mu(r-1)| \neq 0$ then we set

$$\ell_p(\lambda|p\mu) = r. \quad (5.1)$$

Lemma 5.4. *Let $(\lambda|p\mu) \in \mathcal{P}^2(n)$ and let P_ρ be a vertex of the signed Young module $Y(\lambda|p\mu)$.*

- (i) *The signed Young module $Y(p\lambda|p^2\mu)$ has vertex $P_{p\rho}$.*
- (ii) *Suppose that $k > \ell_p(\lambda|p\mu)$ and let $(\alpha|\beta) \in \mathcal{P}^2(m)$ for some $m \in \mathbb{N}$. Then $Y(\lambda + p^k \alpha | p(\mu + p^k \beta))$ has vertex $P_\rho \times P_{p^k \gamma}$ where P_γ is a vertex of $Y(\alpha|p\beta)$. Moreover,*

$$Y(\lambda|p\mu)(P_\rho) \boxtimes Y(p^k \alpha | p^{k+1} \beta)(P_{p^k \gamma})$$

is isomorphic to the Broué correspondent $Y(\lambda + p^k \alpha | p(\mu + p^k \beta))(P_\rho \times P_{p^k \gamma})$.

Proof. Suppose that λ and μ have p -adic expansions $\sum_{i \geq 0} p^i \lambda(i)$ and $\sum_{i \geq 0} p^i \mu(i)$, respectively. It is clear that the partitions $p\lambda$ and $p\mu$ have p -adic expansions $p\lambda = \sum_{i \geq 1} p^i \lambda(i-1)$ and $p\mu = \sum_{i \geq 1} p^i \mu(i-1)$, respectively. So $|(p\lambda)(0)| = 0$, and $|(p\lambda)(i)| + |(p\mu)(i-1)| = |\lambda(i-1)| + |\mu(i-2)|$ for all $i \geq 1$, where we set $\mu(-1) = \emptyset$. By Definition 4.11, $Y(p\lambda|p^2\mu)$ has vertex $P_{p\rho}$, proving part (i).

Let $r = \ell_p(\lambda|p\mu)$. For part (ii), since $k > r$, the p -adic expansions of $\lambda + p^k\alpha$ and $\mu + p^k\beta$ are

$$\begin{aligned}\lambda + p^k\alpha &= \sum_{0 \leq i \leq r} p^i \lambda(i) + \sum_{i \geq k} p^i \alpha(i - k), \\ \mu + p^k\beta &= \sum_{0 \leq i \leq r} p^i \mu(i) + \sum_{i \geq k} p^i \beta(i - k),\end{aligned}$$

respectively. By Definition 4.11, $Y(\lambda + p^k\alpha | p(\mu + p^k\beta))$ has vertex P_η where

$$\eta = (1^{|\lambda(0)|}, p^{|\lambda(1)|+|\mu(0)|}, \dots, (p^r)^{|\lambda(r)|+|\mu(r-1)|}, (p^k)^{|\alpha(0)|}, (p^{k+1})^{|\alpha(1)|+|\beta(0)|}, \dots) = \rho \bullet p^k\gamma.$$

Thus $P_\eta = P_{\rho \bullet p^k\gamma} = P_\rho \times P_{p^k\gamma}$. By Definition 4.11 and Lemma 5.3, we have

$$\begin{aligned}Y(\lambda + p^k\alpha | p(\mu + p^k\beta))(P_{\rho \bullet p^k\gamma}) &= Q_1(\lambda(0)|\varnothing) \boxtimes Q_p(\lambda(1)|\mu(0)) \boxtimes \cdots \boxtimes Q_{p^r}(\lambda(r)|\mu(r-1)) \\ &\quad \boxtimes Q_{p^k}(\alpha(0)|\varnothing) \boxtimes Q_{p^{k+1}}(\alpha(1)|\beta(0)) \boxtimes \cdots \\ &\cong Y(\lambda|p\mu)(P_\rho) \boxtimes Y(p^k\alpha|p^{k+1}\beta)(P_{p^k\gamma}),\end{aligned}$$

as required. \square

The following result is an interesting special case of [6, Theorem 3.18]. It is included to illustrate a technique used again in the proof of Proposition 7.1.

Lemma 5.5. *Let $n \in \mathbb{N}$. If $n = mp + c$ where $m \in \mathbb{N}_0$ and $0 \leq c < p$ then $\text{sgn}(n) \cong Y((1^c)|(mp))$.*

Proof. Let $n = \sum_{i=0}^r p^i n_i$ be the p -adic expansion of n , and let $\rho = (1^{n_0}, p^{n_1}, \dots, (p^r)^{n_r})$. By Definition 4.11, the signed Young module $Y((1^c)|(mp))$ has P_ρ as a vertex and

$$Y((1^c)|(mp))(P_\rho) \cong Q_1((1^c)|\varnothing) \boxtimes Q_p(\varnothing|(n_1)) \boxtimes \cdots \boxtimes Q_{p^r}(\varnothing|(n_r))$$

as a module for $F[N_\rho/P_\rho]$. Since $n_i < p$ we have

$$Q_{p^i}(\varnothing|(n_i)) \cong \text{Inf}_{\mathfrak{S}_{n_i}}^{(N_{p^i}/P_{p^i}) \wr \mathfrak{S}_{n_i}}(F(n_i)) \otimes \widehat{\text{sgn}(N_{p^i})}^{\otimes n_i} \cong \text{Res}_{N_{p^i} \wr \mathfrak{S}_{n_i}}^{\mathfrak{S}_{p^i n_i}}(\text{sgn}(p^i n_i))$$

where the second isomorphism follows from (2.2), regarding the right-hand side as a representation of $(N_{p^i}/P_{p^i}) \wr \mathfrak{S}_{n_i}$. Hence there is an isomorphism of FN_ρ -modules,

$$Y((1^c)|(mp))(P_\rho) \cong \text{Res}_{N_\rho}^{\mathfrak{S}_n}(\text{sgn}(n)).$$

On the other hand, since P_ρ is a Sylow p -subgroup of \mathfrak{S}_n , it is a vertex of $\text{sgn}(n)$, and clearly $\text{sgn}(n)(P_\rho) \cong \text{Res}_{N_\rho}^{\mathfrak{S}_n}(\text{sgn}(n))$ as an FN_ρ -module. The Broué correspondence is bijective (see Theorem 2.5), so we have $Y((1^c)|(mp)) \cong \text{sgn}(n)$. \square

6. SIGNED p -KOSTKA NUMBERS

In this section we prove Theorem 1.2 and Theorem 1.3. We work mainly with the $F[(N_k/P_k) \wr \mathfrak{S}_m]$ -modules $W_k(\gamma|\delta)$ and $Q_k(\alpha|\beta)$ defined in Definitions 3.6 and 4.7, and the $F[C_2 \wr \mathfrak{S}_m]$ -modules $\overline{W}_k(\gamma|\delta)$ and $\overline{Q}_k(\alpha|\beta)$ obtained from them by factoring out the trivial action of the even permutations in the base group of the wreath product.

We begin with a key lemma for the proof of Theorem 1.2.

Lemma 6.1. *Let $n \in \mathbb{N}$. For any $(\gamma|\delta) \in \mathcal{C}^2(n)$ and $(\lambda|\mu) \in \mathcal{RP}^2(n)$ we have*

- (i) $[W_{p^{i+1}}(\gamma|\delta) : Q_{p^{i+1}}(\lambda|\mu)] = [W_{p^i}(\gamma|\delta) : Q_{p^i}(\lambda|\mu)]$ for all $i \geq 1$,
- (ii) $[W_p(\gamma|\emptyset) : Q_p(\lambda|\emptyset)] = [W_1(\gamma|\emptyset) : Q_1(\lambda|\emptyset)]$,
- (iii) $[W_p(\gamma|\delta) : Q_p(\lambda|\emptyset)] = 0$ if $\delta \neq \emptyset$.

Proof. By Lemma 3.7 and Lemma 4.9 we have $\overline{Q}_{p^j}(\lambda|\mu) \cong R_2(\lambda|\mu)$ and $\overline{W}_{p^j}(\gamma|\delta) \cong V_2(\gamma|\delta)$ for all $j \geq 1$. Part (i) now follows by applying Lemma 2.1. For (ii), if $\delta = \mu = \emptyset$ then

$$\begin{aligned} \overline{W}_p(\gamma|\emptyset) &\cong V_2(\gamma|\emptyset) = \text{Inf}_{\mathfrak{S}_m}^{C_2 \wr \mathfrak{S}_m}(M^\gamma) = \overline{W}_1(\gamma|\emptyset), \\ \overline{Q}_p(\lambda|\emptyset) &\cong R_2(\lambda|\emptyset) = \text{Inf}_{\mathfrak{S}_m}^{C_2 \wr \mathfrak{S}_m}(P^\lambda) = \overline{Q}_1(\lambda|\emptyset). \end{aligned}$$

So $[\overline{W}_p(\gamma|\emptyset) : \overline{Q}_p(\lambda|\emptyset)] = [\overline{W}_1(\gamma|\emptyset) : \overline{Q}_1(\lambda|\emptyset)]$. Now apply Lemma 2.1. Finally (iii) follows from Proposition 4.5 and Lemma 2.1. \square

We are now ready to prove Theorem 1.2.

Proof of Theorem 1.2. Let P_ρ be a vertex of $Y(\lambda|p\mu)$. By Definition 4.11 we have

$$\rho = (1^{n_0}, p^{n_1}, (p^2)^{n_2}, \dots, (p^r)^{n_r}),$$

where $n_0 = |\lambda(0)|$ and $n_i = |\lambda(i)| + |\mu(i-1)|$ for all $i \in \{1, \dots, r\}$. By the Broué correspondence (see Theorem 2.5) and the description of the Broué correspondents of signed Young modules in Lemma 5.4, it is equivalent to show that

$$[M(p\alpha|p\beta)(P_{p\rho}) : Y(p\lambda|p^2\mu)(P_{p\rho})] \leq [M(\alpha|\beta)(P_\rho) : Y(\lambda|p\mu)(P_\rho)].$$

Let $\Lambda = \Lambda((\alpha|\beta), \rho)$ and $\Lambda' = \Lambda((p\alpha|p\beta), p\rho)$ be as defined in Notation 3.8. Observe that Λ' consists of all compositions

$$(\emptyset, \gamma^{(0)}, \gamma^{(1)}, \dots, \gamma^{(r)}|\emptyset, \delta^{(0)}, \delta^{(1)}, \dots, \delta^{(r)})$$

where $(\gamma^{(0)}, \gamma^{(1)}, \dots, \gamma^{(r)}|\delta^{(0)}, \delta^{(1)}, \dots, \delta^{(r)}) \in \Lambda$. By Lemma 3.12 applied to $M(p\alpha|p\beta)(P_{p\rho})$, we have

$$\begin{aligned} M(p\alpha|p\beta)(P_{p\rho}) &\cong \bigoplus_{(\gamma'|\delta') \in \Lambda'} W_1(\gamma'^{(0)}|\delta'^{(0)}) \boxtimes W_p(\gamma'^{(1)}|\delta'^{(1)}) \boxtimes \dots \boxtimes W_{p^{r+1}}(\gamma'^{(r+1)}|\delta'^{(r+1)}) \\ &= \bigoplus_{(\gamma|\delta) \in \Lambda} W_1(\emptyset|\emptyset) \boxtimes W_p(\gamma^{(0)}|\delta^{(0)}) \boxtimes \dots \boxtimes W_{p^{r+1}}(\gamma^{(r)}|\delta^{(r)}). \end{aligned}$$

By Definition 4.11 and Lemma 5.4(i), we obtain both

$$\begin{aligned} [M(p\alpha|p\beta)(P_{p\rho}) : Y(p\lambda|p^2\mu)(P_{p\rho})] &= \sum_{(\gamma|\delta)\in\Lambda} \prod_{i=0}^r [W_{p^{i+1}}(\gamma^{(i)}|\delta^{(i)}) : Q_{p^{i+1}}(\lambda(i)|\mu(i-1))], \\ [M(\alpha|\beta)(P_\rho) : Y(\lambda|p\mu)(P_\rho)] &= \sum_{(\gamma|\delta)\in\Lambda} \prod_{i=0}^r [W_{p^i}(\gamma^{(i)}|\delta^{(i)}) : Q_{p^i}(\lambda(i)|\mu(i-1))], \end{aligned}$$

where, as usual, $\mu(-1) = \emptyset$. By Lemma 6.1, we have

$$[W_{p^{i+1}}(\gamma^{(i)}|\delta^{(i)}) : Q_{p^{i+1}}(\lambda(i)|\mu(i-1))] = [W_{p^i}(\gamma^{(i)}|\delta^{(i)}) : Q_{p^i}(\lambda(i)|\mu(i-1))]$$

for all $i \geq 1$, and for $i = 0$ whenever $\delta^{(0)} = \emptyset$. Otherwise, when $i = 0$ and $\delta^{(0)} \neq \emptyset$, we have

$$0 = [W_p(\gamma^{(0)}|\delta^{(0)}) : Q_p(\lambda(0)|\emptyset)] \leq [W_1(\gamma^{(0)}|\delta^{(0)}) : Q_1(\lambda(0)|\emptyset)].$$

This completes the proof. \square

Corollary 6.2. *Let $(\alpha|\beta), (\lambda|p\mu) \in \mathcal{P}^2(n)$. Suppose that $\lambda(0) = \emptyset$. Then*

$$[M(p\alpha|p\beta) : Y(p\lambda|p^2\mu)] = [M(\alpha|\beta) : Y(\lambda|p\mu)].$$

Proof. Let $\rho \in \mathcal{C}(n)$ be defined by

$$\rho = (1^{|\lambda(0)|}, p^{|\lambda(1)|+|\mu(0)|}, \dots, (p^r)^{|\lambda(r)|+|\mu(r-1)|}).$$

The vertex P_ρ of $Y(\lambda|p\mu)$ has no fixed points in $\{1, 2, \dots, n\}$. Hence $\delta^{(0)} = \emptyset$ for any $(\gamma|\delta) \in \Lambda((\alpha|\beta), \rho)$. The result now follows from Theorem 1.2. \square

It is now very easy to deduce the asymptotic stability of signed p -Kostka numbers mentioned in the introduction.

Corollary 6.3. *Let $(\alpha|\beta), (\lambda|p\mu) \in \mathcal{P}^2(n)$. Then, for every natural number $k \geq 2$, we have*

$$[M(p^k\alpha|p^k\beta) : Y(p^k\lambda|p^{k+1}\mu)] = [M(p\alpha|p\beta) : Y(p\lambda|p^2\mu)] \leq [M(\alpha|\beta) : Y(\lambda|p\mu)].$$

Proof. This follows immediately from Corollary 6.2 and Theorem 1.2. \square

Example 6.4. We present a family of examples where the inequality in Theorem 1.2 is strict. Let $0 < c < p$, let $m \in \mathbb{N}$ and let $n = mp + c$. Since $\mathfrak{S}_{mp} \times \mathfrak{S}_c$ has index coprime to p in \mathfrak{S}_n , the trivial module $Y((n)|\emptyset)$ is a direct summand of $M((mp, c)|\emptyset)$; the multiplicity is 1 since $M((mp, c)|\emptyset)$ comes from a transitive action of \mathfrak{S}_n . By Lemma 5.5 we have $\text{sgn}(n) \cong Y((1^c)|(mp))$. Thus

$$\begin{aligned} &[M(\emptyset|(mp, c) : Y((1^c)|(mp))] \\ &= [M(\emptyset|(mp, c)) \otimes \text{sgn}(n) : Y((1^c)|(mp)) \otimes \text{sgn}(n)] \\ &= [M((mp, c)|\emptyset) : Y((n)|\emptyset)] \\ &= 1. \end{aligned}$$

On the other hand, $[M((mp^2, cp)|\emptyset) : Y((mp^2)|(p(1^c)))] = 0$ because, by [7, 2.3(6)], the signed Young modules are pairwise non-isomorphic and so the signed Young module $Y((mp^2)|p(1^c))$ is not isomorphic to a Young module. Thus we have

$$\begin{aligned} & [M(\emptyset|p(mp, c)) : Y(p(1^c)|p(mp))] \\ &= [M(\emptyset|(mp^2, cp)) \otimes \text{sgn}(np) : Y(p(1^c)|p(mp)) \otimes \text{sgn}(np)] \\ &= [M((mp^2, cp)|\emptyset) : Y((mp^2)|p(1^c))] \\ &= 0, \end{aligned}$$

where the penultimate equation is obtained using [6, Theorem 3.18]. This shows that

$$[M(\emptyset|p(mp, c)) : Y(p(1^c)|p(mp))] < [M(\emptyset|(mp, c)) : Y((1^c)|(mp))].$$

We now turn to the proof of Theorem 1.3. We need a further result on the Brauer quotients of signed Young permutation modules.

Proposition 6.5. *Let $m, n \in \mathbb{N}$ and let $(\pi|\tilde{\pi}) \in \mathcal{C}^2(m)$ and $(\phi|\tilde{\phi}) \in \mathcal{C}^2(n)$. Let $\rho \in \mathcal{C}(m)$ and $\gamma \in \mathcal{C}(n)$ be compositions of the form*

$$\begin{aligned} \rho &= (1^{m_0}, p^{m_1}, \dots, (p^r)^{m_r}), \\ \gamma &= (1^{n_0}, p^{n_1}, \dots, (p^s)^{n_s}). \end{aligned}$$

For all $k \in \mathbb{N}$ such that $k > r$, we have that $M(\pi|\tilde{\pi})(P_\rho) \boxtimes M(p^k\phi|p^k\tilde{\phi})(P_{p^k\gamma})$ is isomorphic to a direct summand of $M(\pi + p^k\phi|\tilde{\pi} + p^k\tilde{\phi})(P_{\rho \bullet p^k\gamma})$. Furthermore, if $p^k > \max\{\pi_1, \tilde{\pi}_1\}$, then

$$M(\pi|\tilde{\pi})(P_\rho) \boxtimes M(p^k\phi|p^k\tilde{\phi})(P_{p^k\gamma}) \cong M(\pi + p^k\phi|\tilde{\pi} + p^k\tilde{\phi})(P_{\rho \bullet p^k\gamma}),$$

as $F[N_{\mathfrak{S}_{m+p^k n}}(P_{\rho \bullet p^k\gamma})/P_{\rho \bullet p^k\gamma}]$ -modules.

Note that, in Proposition 6.5, while $\sum_{i=0}^r m_i p^i = m$ and $\sum_{i=0}^r n_i p^i = n$, these need not be the base p expressions for either m or n .

Proof. Since $k > r$, by Lemma 5.3, we have

$$P_{\rho \bullet p^k\gamma} = P_\rho \times P_{p^k\gamma}.$$

To ease the notation, we denote by M , M_1 and M_2 the modules $M(\pi + p^k\phi|\tilde{\pi} + p^k\tilde{\phi})$, $M(\pi|\tilde{\pi})$ and $M(p^k\phi|p^k\tilde{\phi})$, respectively. Furthermore, let $P = P_{\rho \bullet p^k\gamma}$. By Corollary 3.4, $M(P)$ has as a basis the subset \mathcal{B} of $\Omega(\pi + p^k\phi|\tilde{\pi} + p^k\tilde{\phi})$ consisting of all $\{\mathbf{R}\}$ such that \mathbf{R} is a row standard $(\pi + p^k\phi|\tilde{\pi} + p^k\tilde{\phi})$ -tableau whose rows are unions of P -orbits. Similarly, we define bases \mathcal{B}_1 and \mathcal{B}_2 of $\Omega(\pi|\tilde{\pi})$ and $\Omega(p^k\phi|p^k\tilde{\phi})$ for $M_1(P_\rho)$ and $M_2(P_{p^k\gamma})$, respectively; here each $(p^k\phi|p^k\tilde{\phi})$ -tableau \mathbf{S} of \mathcal{B}_2 is filled with the numbers $m+1, m+2, \dots, m+p^k n$.

For $\{\mathbf{T}\} \in \mathcal{B}_1$ and $\{\mathbf{S}\} \in \mathcal{B}_2$, let

$$\psi : \mathcal{B}_1 \times \mathcal{B}_2 \longrightarrow \mathcal{B}$$

be the map defined by

$$\psi(\{\mathbf{T}\}, \{\mathbf{S}\}) = \{(\mathbf{R}_+|\mathbf{R}_-)\},$$

where R_+ is the row standard $(\pi + p^k\phi)$ -tableau such that row i of R_+ is the union of row i of T_+ and row i of S_+ , and R_- is the row standard $(\tilde{\pi} + p^k\tilde{\phi})$ -tableau such that row i of R_- is the union of row i of T_- and row i of S_- . Here we have used the convention row i of T_+ is empty if $i > \ell(\pi)$, and so on. The map ψ is well defined since the rows of $R = (R_+|R_-)$ are union of orbits of $P = P_\rho \times P_{p^k\gamma}$ on $\{1, 2, \dots, m + p^k n\}$.

Clearly ψ is injective and so it induces an injection of vector spaces

$$\theta : M_1(P_\rho) \boxtimes M_2(P_{p^k\gamma}) \longrightarrow M(P)$$

defined by $\theta(\{T\} \otimes \{S\}) = \psi(\{T\}, \{S\})$. By Lemma 5.3, we may regard the domain and codomain of θ as $FN_{\mathfrak{S}_{m+p^k n}}(P)$ -modules with trivial P -action. It is not difficult to check that

$$\theta(g(\{T\} \otimes \{S\})) = g\theta(\{T\} \otimes \{S\}),$$

for all $g \in N_{\mathfrak{S}_{m+p^k n}}(P)$, $\{T\} \in \mathcal{B}_1$ and $\{S\} \in \mathcal{B}_2$. Therefore θ is an injective homomorphism of $FN_{\mathfrak{S}_{m+p^k n}}(P)$ -modules, and hence an injective homomorphism of $F[N_{\mathfrak{S}_{m+p^k n}}(P)/P]$ -modules. Since both $M_1(P_\rho)$ and $M_2(P_{p^k\gamma})$ are projective and hence injective, their outer tensor product is also injective. Therefore, the map θ splits and we obtain that $M_1(P_\rho) \boxtimes M_2(P_{p^k\gamma})$ is a direct summand of $M(P)$.

The second assertion follows easily by observing that, if $p^k > \max\{\pi_1, \tilde{\pi}_1\}$, then the map ψ defined above is a bijection. \square

We are now ready to prove Theorem 1.3.

Proof of Theorem 1.3. Let $\rho \in \mathcal{C}(m)$ and $\gamma \in \mathcal{C}(n)$ be defined by

$$\begin{aligned} \rho &= (1^{|\lambda(0)|}, p^{|\lambda(1)|+|\mu(0)|}, \dots, (p^r)^{|\lambda(r)|+|\mu(r-1)|}), \\ \gamma &= (1^{|\alpha(0)|}, p^{|\alpha(1)|+|\beta(0)|}, \dots, (p^s)^{|\alpha(s)|+|\beta(s-1)|}), \end{aligned}$$

where $r = \ell_p(\lambda|p\mu)$ and $s = \ell_p(\alpha|p\beta)$, respectively. By Definition 4.11, P_ρ is a vertex of $Y(\lambda|p\mu)$ and P_γ is a vertex of $Y(\alpha|p\beta)$. Since $k > r$, by Lemma 5.3, we have $P_{\rho \bullet p^k\gamma} = P_\rho \times P_{p^k\gamma}$ and

$$N_{\mathfrak{S}_{m+p^k n}}(P_\rho \times P_{p^k\gamma}) = N_{\mathfrak{S}_m}(P_\rho) \times N_{\mathfrak{S}_{p^k n}}(P_{p^k\gamma}).$$

By Lemma 5.4, $Y(p^k\alpha|p^{k+1}\beta)$ has vertex $P_{p^k\gamma}$ and $Y(\lambda + p^k\alpha|p(\mu + p^k\beta))$ has vertex $P_{\rho \bullet p^k\gamma}$. Moreover, the Broué correspondent of $Y(\lambda + p^k\alpha|p(\mu + p^k\beta))$ is

$$Y(\lambda|p\mu)(P_\rho) \boxtimes Y(p^k\alpha|p^{k+1}\beta)(P_{p^k\gamma}).$$

By Proposition 6.5, we have

$$M(\pi|\tilde{\pi})(P_\rho) \boxtimes M(p^k\phi|p^k\tilde{\phi})(P_{p^k\gamma}) \mid M(\pi + p^k\phi|\tilde{\pi} + p^k\tilde{\phi})(P_{\rho \bullet p^k\gamma}).$$

Therefore, using Theorem 2.5(ii), we deduce that

$$\begin{aligned}
& [M(\pi + p^k \phi | \tilde{\pi} + p^k \tilde{\phi}) : Y(\lambda + p^k \alpha | p(\mu + p^k \beta))] \\
&= [M(\pi + p^k \phi | \tilde{\pi} + p^k \tilde{\phi})(P_{\rho \bullet p^k \gamma}) : Y(\lambda + p^k \alpha | p(\mu + p^k \beta))(P_{\rho \bullet p^k \gamma})] \\
&\geq [M(\pi | \tilde{\pi})(P_\rho) \boxtimes M(p^k \phi | p^k \tilde{\phi})(P_{p^k \gamma}) : Y(\lambda | p\mu)(P_\rho) \boxtimes Y(p^k \alpha | p^{k+1} \beta)(P_{p^k \gamma})] \\
&= [M(\pi | \tilde{\pi})(P_\rho) : Y(\lambda | p\mu)(P_\rho)] [M(p^k \phi | p^k \tilde{\phi})(P_{p^k \gamma}) : Y(p^k \alpha | p^{k+1} \beta)(P_{p^k \gamma})] \\
&= [M(\pi | \tilde{\pi}) : Y(\lambda | p\mu)] [M(p^k \phi | p^k \tilde{\phi}) : Y(p^k \alpha | p^{k+1} \beta)] \\
&= [M(\pi | \tilde{\pi}) : Y(\lambda | p\mu)] [M(p \phi | p \tilde{\phi}) : Y(p \alpha | p^2 \beta)],
\end{aligned}$$

where the final equality follows from Corollary 6.3. If $p^k > \max\{\pi_1, \tilde{\pi}_1\}$, then Proposition 6.5 implies that we have equalities throughout. \square

7. INDECOMPOSABLE SIGNED YOUNG PERMUTATION MODULES

In this section, in the spirit of Gill's result [12, Theorem 2], we classify all indecomposable signed Young permutation modules over the field F and determine their endomorphism algebras and their labels as signed Young modules. By [12], any indecomposable Young permutation module is of the form $M^{(m)}$ or $M^{(kp-1,1)}$. It is immediate from the definition of signed Young permutation modules in (1.1) that

$$M(\alpha|\beta) \cong \text{Ind}_{\mathfrak{S}_{|\alpha|} \times \mathfrak{S}_{|\beta|}}^{\mathfrak{S}_{|\alpha|+|\beta|}} (M^\alpha \boxtimes (M^\beta \otimes \text{sgn}(|\beta|))).$$

As such, by Gill's result, any indecomposable signed Young permutation module is of one of the forms $M((m)|(n))$, $M((m)|(kp-1,1))$, $M((kp-1,1)|(m))$ or $M((kp-1,1)|(\ell p-1,1))$. Since $M((m)|(kp-1,1)) \otimes \text{sgn}(m+kp) \cong M((kp-1,1)|(m))$, there are essentially three different forms to consider.

Proof of Theorem 1.4. Let $M_1 = M((m)|(n))$. If $m = 0$ then M_1 is the sign representation, and if $n = 0$ then M_1 is the trivial representation. In these cases, M_1 is simple with 1-dimensional endomorphism ring. Suppose that both m, n are non-zero. By the Littlewood–Richardson rule, the module M_1 has a Specht series with top Specht factor $S^{(m+1,1^{n-1})}$ and bottom Specht factor $S^{(m,1^n)}$. If $m+n$ is not divisible by p then the p -cores of $(m+1,1^{n-1})$ and $(m,1^n)$ are non-empty and distinct and so $S^{(m+1,1^{n-1})}$ and $S^{(m,1^n)}$ lie in different blocks. Consequently, M_1 is decomposable. Now suppose that $m+n$ is divisible by p . In this case, by Peel's result [25],

$$S^{(m+1,1^{n-1})} = \begin{cases} F & n = 1, \\ \begin{bmatrix} D^\lambda \\ D^\gamma \end{bmatrix} & n \geq 2, \end{cases} \quad S^{(m,1^n)} = \begin{cases} \text{sgn}(m+n) & m = 1, \\ \begin{bmatrix} D^\mu \\ D^\lambda \end{bmatrix} & m \geq 2, \end{cases}$$

where μ, λ, γ are the p -regularization of the partitions $(m,1^n)$, $(m+1,1^{n-1})$ and $(m+2,1^{n-2})$ respectively (see [20, 6.3.48]). If $m = 1$ then $M((1)|(n)) \cong M(\emptyset|(n,1))$ is indecomposable. Similarly, if $n = 1$ then $M((m)|(1)) \cong M((m,1)|\emptyset)$ is indecomposable. Moreover, since $M((m,1)|\emptyset)$ has a Loewy series with factors $F, D^{(m,1)}, F$, the endomorphism algebra

$\text{End}_{F\mathfrak{S}_{m+1}}M((m, 1)|\emptyset)$ is 2-dimensional. Tensoring by the sign representation we obtain the same result for $\text{End}_{F\mathfrak{S}_{n+1}}(\emptyset|(n, 1))$.

We now study the case when $m, n \geq 2$. In this case, both the head and socle of M_1 contain the simple module D^λ . Also, as a signed Young permutation module, M_1 is self-dual. Suppose that D^γ is not isomorphic to a composition factor of any direct summand of M_1 containing D^λ in its head (and hence in its socle). Then D^γ is necessarily isomorphic to a direct summand of M_1 . From the Specht series, there is a surjection ψ from M_1 onto the Specht module $S = S^{(m+1, 1^{n-1})}$. Since S has composition factors D^γ and D^λ , we have $\psi(D^\gamma) \neq 0$ and so $\psi(D^\gamma) \cong D^\gamma$. Let Y be an indecomposable direct summand of M_1 such that $\psi(Y)$ contains a composition factor D^λ . This shows that $\psi(Y) \cong D^\lambda$ and hence

$$S = \psi(D^\gamma \oplus Y) \cong D^\gamma \oplus Y/(Y \cap \ker \psi) \cong D^\gamma \oplus D^\lambda.$$

This is absurd since S is indecomposable. Hence there exists an indecomposable direct summand of M_1 containing D^λ in its head and that does not contain D^γ in its head or in its socle. Dually, there exists an indecomposable direct summand of M_1 containing D^λ in its head, that does not contain D^μ in its head or in its socle. Thus the only possibility is that M_1 is indecomposable with the Loewy structure

$$\begin{bmatrix} D^\lambda \\ D^\mu & D^\gamma \\ D^\lambda \end{bmatrix}$$

and has 2-dimensional endomorphism ring.

Let $M_2 = M((kp - 1, 1)|(m))$. By Gill's result, if $m = 0$ then M_2 is indecomposable and if $m = 1$ then $M_2 \cong M((kp - 1, 1^2)|\emptyset)$ is decomposable. Suppose that $m \geq 2$. By the Young and Littlewood–Richardson rules, M_2 has a Specht series with Specht factors $S_1 = S^{(kp+1, 1^{m-1})}$, $S_2 = S^{(kp, 2, 1^{m-2})}$, $S_3 = S^{(kp, 1^m)}$, $S_4 = S^{(kp-1, 2, 1^{m-1})}$, $S_5 = S^{(kp-1, 1^{m+1})}$, with S_3 occurring twice. If $m \not\equiv 0 \pmod{p}$, then S_1 and S_3 lie in different blocks. If $m \equiv 0 \pmod{p}$ then S_3 and S_4 belong to different blocks. Thus we conclude that M_2 is decomposable whenever $m \geq 2$.

Let $M_3 = M((kp - 1, 1)|(\ell p - 1, 1))$. Then $M_3 \cong M((kp - 1, 1^2)|(\ell p - 1))$. By Gill's result, since $M^{(kp-1, 1^2)}$ is decomposable, we have that

$$M((kp - 1, 1^2)|(\ell p - 1)) = \text{Ind}_{\mathfrak{S}_{kp+1} \times \mathfrak{S}_{\ell p-1}}^{\mathfrak{S}_{kp+\ell p}} (M^{(kp-1, 1^2)} \boxtimes (M^{(\ell p-1)} \otimes \text{sgn}(\ell p - 1)))$$

is decomposable. □

We end by determining the labels of the indecomposable signed Young permutation modules. By the remark immediately following the statement of Theorem 1.4, it suffices to consider the modules $M((m)|(n))$ where either $m = 0$, $n = 0$ or $m + n$ is divisible by p .

Proposition 7.1. *Let $m, n \in \mathbb{N}$. Let $n = n_0 + pn'$ where $0 \leq n_0 < p$. There are isomorphisms $M((m)|\emptyset) \cong Y((m)|\emptyset)$, $M(\emptyset|(n)) \cong Y((1^{n_0})|(pn'))$ and, provided $m + n$ is divisible by p , $M((m)|(n)) \cong Y((m, 1^{n_0})|(pn'))$.*

Proof. Clearly $M((n)|\emptyset) \cong Y((n)|\emptyset) \cong F(n)$. The second isomorphism follows from Lemma 5.5. In the remaining case, $m, n > 0$ and $m + n$ is divisible by p . Let $m = \sum_{i \geq 0} m_i p^i$ and let $n = \sum_{i \geq 0} n_i p^i$ be the p -adic expansions. Let r be the greatest integer such that $m_r + n_r \neq 0$. Let P be a Sylow p -subgroup of $\mathfrak{S}_m \times \mathfrak{S}_n$. By Proposition 3.12 we have an isomorphism of $F[N_{\mathfrak{S}_{m+n}}(P)/P]$ -modules

$$M((m)|(n))(P) \cong W_1((m_0)|(n_0)) \boxtimes W_p((m_1)|(n_1)) \boxtimes \cdots \boxtimes W_{p^r}((m_r)|(n_r)).$$

By Theorem 3.3, the signed Young module $Y((m, 1^{n_0})|(pn'))$ satisfies

$$Y((m, 1^{n_0})|(pn'))(P) = Y((m_0, 1^{n_0})|\emptyset) \boxtimes Q_p((m_1)|(n_1)) \boxtimes \cdots \boxtimes Q_{p^r}((m_r)|(n_r)),$$

where $Q_{p^i}((m_i)|(n_i))$ is the $F[(N_k/P_k) \wr \mathfrak{S}_m]$ -module defined in Definition 4.7. The Broué correspondence is bijective (see Theorem 2.5), so it suffices to prove that the tensor factors in these two modules agree.

Observe that $m_0 + n_0$ is a multiple of p and $m_0 + n_0 < 2p$. If $m_0 = n_0 = 0$ we have $W_1(\emptyset|\emptyset) = Y(\emptyset|\emptyset)$. Next, we assume that $m_0 + n_0 = p$. The $F\mathfrak{S}_p$ -module $W_1((m_0)|(n_0)) \cong M((m_0)|(n_0))$ is indecomposable by Theorem 1.4. The only signed Young module for $F\mathfrak{S}_p$ that is not a Young module is the sign representation. Since $n_0 < p$ we see that $M((m_0)|(n_0))$ is a Young module. The proof of Theorem 1.4 shows that it has a Specht filtration with $S^{(m_0, 1^{n_0})}$ at the bottom and $S^{(m_0+1, 1^{n_0-1})}$ at the top. Therefore $W_1((m_0)|(n_0)) = M((m_0)|(n_0)) \cong Y((m_0, 1^{n_0})|\emptyset)$, as required.

Finally suppose that $i \geq 1$. By Definition 3.6(ii) $W_{p^i}((m_i)|(n_i))$ is the $F[(N_{p^i}/P_{p^i}) \wr \mathfrak{S}_{m_i+n_i}]$ -module obtained from

$$\mathrm{Ind}_{N_{p^i} \wr (\mathfrak{S}_{m_i} \times \mathfrak{S}_{n_i})}^{N_{p^i} \wr \mathfrak{S}_{m_i+n_i}} \left(\mathrm{Inf}_{\mathfrak{S}_{m_i}}^{N_{p^i} \wr \mathfrak{S}_{m_i}}(F(m_i)) \boxtimes \left(\mathrm{Inf}_{\mathfrak{S}_{n_i}}^{N_{p^i} \wr \mathfrak{S}_{n_i}}(F(n_i)) \otimes \widehat{\mathrm{sgn}}(N_{p^i})^{\otimes n_i} \right) \right).$$

by the canonical surjection $(N_{p^i} \wr \mathfrak{S}_{m_i+n_i})/(P_{p^i})^{m_i+n_i} \cong (N_{p^i}/P_{p^i}) \wr \mathfrak{S}_{m_i+n_i}$. Since $m_i, n_i < p$ the projective covers $P^{(m_i)}$ and $P^{(n_i)}$ are the trivial $F\mathfrak{S}_{m_i}$ - and $F\mathfrak{S}_{n_i}$ -modules, respectively. Therefore, by (4.1), we have $W_{p^i}((m_i)|(n_i)) \cong Q_{p^i}((m_i)|(n_i))$, again as required. \square

Acknowledgements. We thank Kai Meng Tan for his valuable suggestions made at the start of this work. We thank an anonymous referee for a careful reading of this paper.

REFERENCES

- [1] ALPERIN, J. L. *Local representation theory*, vol 11 of *Cambridge Studies in Advanced Mathematics*, Cambridge University Press, 1986.
- [2] BENSON, D. J. *Representations and cohomology I*, vol 30 of *Cambridge studies in advanced mathematics*, Cambridge University Press 1995.
- [3] BROUÉ, M. On Scott modules and p -permutation modules: an approach through the Brauer homomorphism. *Proc. Amer. Math. Soc.*, 93 no. 3 (1985), 401–408.
- [4] BRUNDAN, J. AND KUJAWA, J. A new proof of the Mullineux Conjecture. *J. Alg. Combinat.*, 18 (2003), 13–39.
- [5] DANZ, S., GIANNELLI, E. AND LIM, K. J. On signed p -Kostka matrices, in preparation.
- [6] DANZ, S. AND LIM, K. J. Signed Young modules and simple Specht modules. *Adv. Math.*, 307 (2017), 369–416.

- [7] DONKIN, S. Symmetric and exterior powers, linear source modules and representations of Schur superalgebras. *Proc. London Math. Soc.* (3), 83 (2001), 647–680.
- [8] DONKIN, S. Some remarks on Gill's theorems on Young modules *J. Algebra*, 430, (2015), 1–14.
- [9] ERDMANN, K. Young modules for symmetric groups. *J. Aust. Math. Soc.*, 71 no. 2 (2001), 201–210.
- [10] ERDMANN, K. AND SCHROLL, S. On Young modules of general linear groups. *J. Algebra*, 310 no. 1, (2007), 434–451.
- [11] FANG, M., HENKE, A. AND KOENIG, S. Comparing $GL(n)$ -representations by characteristic-free isomorphisms between generalized Schur algebras. *Forum Math.*, 20 no. 1 (2008), 45–79. With an appendix by Stephen Donkin.
- [12] GILL, C. Young module multiplicities, decomposition numbers and the indecomposable Young permutation modules. *J. Algebra Appl.*, 13, no. 5 (2014) 1350147, 23 pp.
- [13] GRABMEIER, J. Unzerlegbare Moduln mit trivialer Youngquelle und Darstellungstheorie der Schuralgebra, *Bayreuth. Math. Schr.*, 20 (1985), 9–152.
- [14] GREEN, J. A. *Polynomial representations of GL_n* , vol. 830 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1980.
- [15] HEMMER, D. J. Irreducible Specht modules are signed Young modules. *J. Algebra*, 305 (2006) 433–441.
- [16] HENKE, A. On p -Kostka numbers and Young modules, *European J. Combin.*, 26 (2005), 923–942.
- [17] HENKE, A. AND KOENIG, S. Relating polynomials $GL(n)$ -representations in different degrees. *J. Reine Angew. Math.*, 551 (2002) 219–235.
- [18] JAMES, G. D. *The representation theory of the symmetric groups*, vol. 682 of *Lecture Notes in Mathematics*. Springer-Verlag, 1978.
- [19] JAMES, G. D. Trivial source modules for symmetric groups. *Arch. Math. (Basel)*, 41 (1983), 294–300.
- [20] JAMES, G. D. AND KERBER, A. *The representation theory of the symmetric group*. Addison-Wesley, 1981.
- [21] KLESHCHEV, A. Representation theory of symmetric groups and related Hecke algebras *Bull. Amer. Math. Soc.*, 47 (2009), 419–481.
- [22] KLYACHKO, A. A. Direct summand of permutation modules. *Sel. Math. Sov.*, 3 (1983/84) 45–55.
- [23] KUJAWA, J. The Steinberg tensor product theorem for $GL(m|n)$. *Contemp. Math.*, 413 (2006) 123–132.
- [24] PAGET, R. A family of modules with Specht and dual Specht filtrations. *J. Algebra*, 312 (2007), 880–890.
- [25] PEEL, M. H. Hook representations of the symmetric groups. *Glasgow Math. J.*, 12 (1971), 136–149.
- [26] SERRE, J. P. *Linear representations of finite groups*. GTM, vol. 42, Springer, 1977.
- [27] WILDON, M. Multiplicity-free representations of symmetric groups. *J. Pure Appl. Alg.*, 213 (2009), 1464–1477.

(E. Giannelli) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF KAISERSLAUTERN, P.O. Box 3049, 67655 KAISERSLAUTERN, GERMANY

E-mail address: `giannelli@mathematik.uni-kl.de`

(K. J. Lim) DIVISION OF MATHEMATICAL SCIENCES, NANYANG TECHNOLOGICAL UNIVERSITY, SPMS-MAS-03-01, 21 NANYANG LINK, SINGAPORE 637371.

E-mail address: `limkj@ntu.edu.sg`

(W. O'Donovan) DEPARTMENT OF MATHEMATICS, ROYAL HOLLOWAY, UNIVERSITY OF LONDON, UNITED KINGDOM.

E-mail address: `william.odonovan.2014@live.rhul.ac.uk`

(M. Wildon) DEPARTMENT OF MATHEMATICS, ROYAL HOLLOWAY, UNIVERSITY OF LONDON, UNITED KINGDOM.

E-mail address: `mark.wildon@rhul.ac.uk`