

PLETHYSMS OF SYMMETRIC FUNCTIONS AND HIGHEST WEIGHT REPRESENTATIONS

MELANIE DE BOECK, ROWENA PAGET, AND MARK WILDON

ABSTRACT. Let $s_\nu \circ s_\mu$ denote the plethystic product of the Schur functions s_ν and s_μ . In this article we define an explicit polynomial representation corresponding to $s_\nu \circ s_\mu$ with basis indexed by certain ‘plethystic’ semistandard tableaux. Using these representations we prove generalizations of four results on plethysms due to Bruns–Conca–Varbaro, Brion, Ikenmeyer and the authors. In particular, we give a sufficient condition for the multiplicity $\langle s_\nu \circ s_\mu, s_\lambda \rangle$ to be stable under insertion of new parts into μ and λ . We also characterize all maximal and minimal partitions λ in the dominance order such that s_λ appears in $s_\nu \circ s_\mu$ and determine the corresponding multiplicities using plethystic semistandard tableaux.

1. INTRODUCTION

Let s_λ denote the Schur function labelled by the partition λ . Expressing a general plethysm $s_\nu \circ s_\mu$ as a non-negative linear combination of Schur functions has been identified by Stanley as a fundamental open problem in symmetric combinatorics [28, Problem 9]. While many partial results are known, often obtained by deep combinatorial, algebraic or geometric arguments, a satisfying general solution remains out of reach.

In this article we generalize four results on plethysms due to Bruns–Conca–Varbaro [5], Brion [4], Ikenmeyer [17], and the authors [8, 25], and give them unified proofs in the setting of polynomial representations of general linear group. Our proofs are essentially elementary, requiring little more than basic multilinear algebra and the background recalled in §2. The article is intended to be readable by non-experts: in particular, we define the Schur functions s_λ and the plethysm product \circ in §2.2 and include several examples in this introduction.

To state our main results we need the following notation. Let $\text{Par}(r)$ denote the set of partitions of $r \in \mathbf{N}_0$ and let $\ell(\lambda)$ denote the number of parts of the partition λ . Given partitions λ and μ , let $\lambda \sqcup \mu$ be the partition whose multiset of parts is the disjoint union of the multisets of parts of λ and μ . Define $\lambda + \mu$ by $(\lambda + \mu)_i = \lambda_i + \mu_i$. (As a standing convention, if $i > \ell(\lambda)$ then we set $\lambda_i = 0$.) For $N \in \mathbf{N}$ define $N\lambda$ by $(N\lambda)_i = N\lambda_i$ for each i . Let λ' denote the conjugate of the partition λ .

Fix $\mu \in \text{Par}(m)$, $\nu \in \text{Par}(n)$ and $\lambda \in \text{Par}(mn)$.

2010 *Mathematics Subject Classification*. Primary 05E05, Secondary: 05E10, 17B10, 20C30, 22E47.

Theorem 1.1. *If r is at least the greatest part of μ then*

$$\langle s_\nu \circ s_{(r)\sqcup\mu}, s_{(nr)\sqcup\lambda} \rangle = \langle s_\nu \circ s_\mu, s_\lambda \rangle.$$

Theorem 1.2. *If $r \in \mathbf{N}$ then*

$$\langle s_\nu \circ s_{\mu+(1^r)}, s_{\lambda+(n^r)} \rangle \geq \langle s_\nu \circ s_\mu, s_\lambda \rangle.$$

Moreover $\langle s_\nu \circ s_{\mu+N(1^r)}, s_{\lambda+N(n^r)} \rangle$ is constant for

$$N \geq n(\mu_1 + \cdots + \mu_{r-1}) + (n-1)\mu_r + \mu_{r+1} - (\lambda_1 + \cdots + \lambda_r).$$

In particular, $\langle s_\nu \circ s_{\mu+(1^r)}, s_{\lambda+(n^r)} \rangle = \langle s_\nu \circ s_\mu, s_\lambda \rangle$ if $r \geq \ell(\mu)$ and $r \geq \ell(\lambda)$. We give an upper bound for the stable multiplicity at the end of §5.

Theorem 1.3. *If $n^* \in \mathbf{N}$, $\lambda^* \in \text{Par}(mn^*)$ and $\langle s_{(n^*)} \circ s_\mu, s_{\lambda^*} \rangle \geq 1$ then*

$$\langle s_{(n+n^*)} \circ s_\mu, s_{\lambda+\lambda^*} \rangle \geq \langle s_{(n)} \circ s_\mu, s_\lambda \rangle.$$

For our final theorem we need some further combinatorial definitions. Semistandard tableaux, with entries from an arbitrary totally ordered set, are defined in §2.1 below. In particular, $\text{SSYT}_{\mathbf{N}}(\mu)$ denotes the set of semistandard μ -tableaux with entries from \mathbf{N} . We order $\text{SSYT}_{\mathbf{N}}(\mu)$ by the total order defined in Definition 2.1.

Definition 1.4.

- (i) A *plethystic semistandard tableau of shape μ^ν* is a semistandard ν -tableau whose entries are tableaux in $\text{SSYT}_{\mathbf{N}}(\mu)$.
- (ii) Let T be a plethystic semistandard tableau and let M be the greatest entry of the tableau entries of T . The *weight* of T , denoted $\text{wt}(T)$, is the composition $(\beta_1, \dots, \beta_M)$ such that for each b , the total number of occurrences of b in the tableau entries of T is β_b .

These objects are illustrated in Example 1.6.

Theorem 1.5. *The maximal partitions λ in the dominance order such that s_λ is a constituent of $s_\nu \circ s_\mu$ are precisely the maximal weights of the plethystic semistandard tableaux of shape μ^ν . Moreover if λ is such a maximal partition then $\langle s_\nu \circ s_\mu, s_\lambda \rangle$ is the number of plethystic semistandard tableaux of shape μ^ν and weight λ .*

Applying the sign twist in Lemma 2.2 to the main theorems gives equivalent results that are also noteworthy. In particular, Theorem 1.1 implies that if $r \geq \ell(\mu)$ then

$$(1.1) \quad \langle s_\kappa \circ s_{\mu+(1^r)}, s_{\lambda+(1^{nr})} \rangle = \langle s_\nu \circ s_\mu, s_\lambda \rangle,$$

where $\kappa = \nu$ if r is even and $\kappa = \nu'$ if r is odd. The sign-twist of Theorem 1.5 characterizes the minimal partitions λ such that s_λ appears in a general plethysm $s_\nu \circ s_\mu$.



In the setting of polynomial representations of general linear groups, the Schur function s_λ corresponds to the Schur functor ∇^λ . We develop this background in §2. In §3 we construct an explicit model for the module $\nabla^\nu(\nabla^\mu(E))$, where

E is a complex vector space. By Proposition 3.3, the formal character of this module is $(s_\nu \circ s_\mu)(x_1, \dots, x_d)$, where $d = \dim E$. Using this model we prove Theorems 1.1, 1.2, 1.3, 1.5 in §4, §5, §6 and §7, respectively.

Example 1.6. As an illustration of our main theorems, we determine the plethysm $s_{(3)} \circ s_{(3)}$. By Lemma 2.2, its sign twist is $s_{(1^3)} \circ s_{(1^3)}$. Using the closure condition in Definition 1.8 it is easy to show that there are two plethystic semistandard tableaux of shape $(1^3)^{(1^3)}$ and maximal weight, as shown in margin. The weights are $(3, 3, 1, 1, 1)$ and $(3, 2, 2, 2)$ respectively. Hence, by Theorem 1.5 and Lemma 2.2,

$$\langle s_{(3)} \circ s_{(3)}, s_{(5,2,2)} \rangle = \langle s_{(3)} \circ s_{(3)}, s_{(4,4,1)} \rangle = 1.$$

Since $s_{(1^3)} \circ s_\emptyset = s_\emptyset$, it follows from (1.1), applied with $r = 3$, that $\langle s_{(1^3)} \circ s_{(1^3)}, s_{(1^9)} \rangle = 1$, and so $\langle s_{(3)} \circ s_{(3)}, s_{(9)} \rangle = 1$. (This also follows from Theorem 1.5, since the unique plethystic semistandard tableau of shape $(3)^{(3)}$ and maximal weight is



and can, of course, be seen in many other ways.)

By Example 1.10, $s_{(3)} \circ s_{(2)} = s_{(6)} + s_{(4,2)} + s_{(2,2,2)}$. Applying the final part of Theorem 1.2 to the second summand, we have $\langle s_{(3)} \circ s_{(3)}, s_{(7,2)} \rangle = 1$. Moreover, by the Cayley–Sylvester formula (see for instance [13, Exercise 6.18, solution]), $\langle s_{(3)} \circ s_{(3)}, s_{(6,3)} \rangle$ is the number of partitions of 3 contained in the 3×3 box, minus the number of partitions of 2 satisfying the same restriction. Therefore $\langle s_{(3)} \circ s_{(3)}, s_{(6,3)} \rangle = 1$. Hence

$$(1.2) \quad s_{(3)} \circ s_{(3)} = s_{(9)} + s_{(7,2)} + s_{(6,3)} + s_{(5,2,2)} + s_{(4,4,1)} + f$$

for some symmetric function f with non-negative coefficients in the Schur basis. Under the characteristic isometry (see §2.2), $s_{(3)} \circ s_{(3)}$ is the image of the permutation character of S_9 acting on the set partitions of $\{1, \dots, 9\}$ into 3 sets, each of size 3. This character has degree $9!/3!^3 = 280$, which equals the sum of the degrees of the irreducible characters of S_9 labelled by the partitions appearing in (1.2). Hence $f = 0$.

We now explain the antecedents of the main theorems, before giving a more detailed outline of the strategy of our proofs. For general background on symmetric functions, including the Young and Pieri rules, we refer the reader to [27, Ch. 7] or [22, Ch. 1]. Further background on plethysms is given in [21] and the introduction to [26].

Antecedents of the main theorems.

Theorem 1.1. Stated in the language of symmetric functions, Proposition 1.16 of [5] becomes

$$(1.3) \quad \langle s_\nu \circ s_{(1^{m+1})}, s_{(n) \sqcup \lambda} \rangle = \langle s_\nu \circ s_{(1^m)}, s_\lambda \rangle,$$

provided that n is at least the first part of λ . By Remark 4.4, both sides in Theorem 1.1 are zero if $\lambda_1 > nr$. Therefore this proposition is equivalent to the case $\mu = (1^m)$ and $r = 1$ of Theorem 1.1. (The statement in [5] replaces $(n) \sqcup \lambda$ with its conjugate partition $\lambda' + (1^n)$; the conjugation arises because the functor L_λ in [5] is our $\nabla^{\lambda'}$.) The proof in [5] gives a bijection between the invariants in $\otimes^n (\wedge^m E)$ and $\otimes^n (\wedge^{m+1} E)$ for a Borel subgroup of $\mathrm{GL}(E)$. Our proof establishes a corresponding bijection between highest-weight vectors, with \wedge^m replaced with an arbitrary Schur functor. We remark after this proof on the connection with the later proof of (1.3) given in [20, Lemma 3.2].

Specializing (1.3) by taking $\nu = (n)$ or $\nu = (1^n)$ gives two results first proved in [23]. Since Newell's paper is not easy to read, we explain the connection. Taking the inner product of both sides of Theorem 1.1 of [23] with s_λ , where $\lambda \in \mathrm{Par}(mn - k)$, gives

$$(1.4) \quad \langle (s_{(1^k)} \circ s_{(m-1)})(s_{(n-k)} \circ s_{(m)}), s_\lambda \rangle = \langle s_{(n)} \circ s_{(m)}, s_{(1^k)s_\lambda} \rangle.$$

(This corrects a typographical error in [23]: as can be seen from the correctly stated and analogous Theorem 1A, $g_{(1^k)\xi\nu}\{\nu\}$ should be $g_{(1^k)\xi\nu}\{\xi\}$; note that, by definition, $g_{(1^k)\xi\nu} = \langle s_{(1^k)s_\xi}, s_\nu \rangle$.) Taking $k = n$ we obtain $\langle s_{(1^n)} \circ s_{(m-1)}, s_\lambda \rangle = \langle s_{(n)} \circ s_{(m)}, s_{(1^n)s_\lambda} \rangle$. By Pieri's rule, $s_{(1^n)s_\lambda}$ is the sum of all the Schur functions labelled by partitions obtained from λ by adding n boxes, no two in the same row. On the other hand, by Young's rule, $s_{(n)}$ is a summand of $s_{(1)}^n$, so the plethysm $s_{(n)} \circ s_{(m)}$ is contained in $s_{(1)}^n \circ s_{(m)} = (s_{(1)} \circ s_{(m)})^n = s_{(m)}^n$. Another application of Young's rule now shows that if s_λ is a constituent of $s_{(n)} \circ s_{(m)}$ then $\ell(\lambda) \leq n$. Hence

$$(1.5) \quad \langle s_{(1^n)} \circ s_{(m-1)}, s_\lambda \rangle = \langle s_{(n)} \circ s_{(m)}, s_{\lambda+(1^n)} \rangle.$$

Similarly Theorem 1A in [23] implies that

$$(1.6) \quad \langle s_{(n)} \circ s_{(m-1)}, s_\lambda \rangle = \langle s_{(1^n)} \circ s_{(m)}, s_{\lambda+(1^n)} \rangle.$$

These are the special case of the equivalent form of Theorem 1.1 stated in (1.1) when $\mu = (1^m)$, $r = 1$ and $\nu = (n)$ or $\nu = (1^n)$.

It follows from our Theorem 1, or by combining (1.5) and (1.6), that $\langle s_{(n)} \circ s_{(m+2)}, s_{\lambda+(2^n)} \rangle = \langle s_{(n)} \circ s_{(m)}, s_\lambda \rangle$ for all $\lambda \in \mathrm{Par}(mn)$. This was proved by Dent in [9, Theorem 3.8] using the symmetric group.

Theorem 1.2. The special case of part (i) of the theorem on page 354 of [4] when G is $\mathrm{GL}(E)$ asserts that if ρ is any partition then $\langle s_\nu \circ s_{\mu+N\rho}, s_{\lambda+Nn\rho} \rangle$ is a non-decreasing function of $N \in \mathbf{N}_0$. Part (ii) gives a condition in terms of simple roots for the values to stabilise: in the special case of $\mathrm{GL}(E)$, it becomes

$$\mu_i - \mu_{i+1} + N(\rho_i - \rho_{i+1}) \geq n(\mu_1 + \cdots + \mu_i) - (\lambda_1 + \cdots + \lambda_i)$$

for every i such that $\rho_i > \rho_{i+1}$, as stated in [4, page 362, Corollary 1]. Part (iii) gives a technical formula for the stable multiplicity.

Our Theorem 1.2 is Brion's Theorem in the case $\rho = (1^r)$. Brion's theorem for a general partition ρ follows by repeatedly applying our theorem to each column of ρ in turn. Brion's proof uses $\mathrm{GL}(E)$ -invariant vector bundles on the Grassmannian variety of full flags in E and the long exact cohomology sequence. The more elementary proof given here, which leads to a combinatorial upper bound on the stable multiplicity (see Proposition 5.9), is therefore of interest.

Taking $\nu = (n)$, $\mu = (m)$ and $r = 1$ in Theorem 1.2 we obtain $\langle s_{(n)} \circ s_{(m)}, s_\lambda \rangle \leq \langle s_{(n)} \circ s_{(m+1)}, s_{\lambda+(n)} \rangle$. This is Foulkes' Second Conjecture, stated as a working hypothesis at the end of §1 of [12], and proved by Brion in [4].

Theorem 1.3. Proposition 4.3.4 of [17] is equivalent to the special case of Theorem 1.3 when $\mu = (m)$. The proof in [17] uses polynomial representations of $\mathrm{GL}(E)$, where E is a complex vector space. The key idea is to multiply highest-weight vectors in $\mathrm{Sym}^n(\mathrm{Sym}^m E)$ and $\mathrm{Sym}^{n^*}(\mathrm{Sym}^m E)$ to get a highest-weight vector in $\mathrm{Sym}^{n+n^*}(\mathrm{Sym}^m E)$. This generalizes to prove Theorem 1.3. To motivate this proof we digress briefly to illustrate the geometric interpretation of this multiplication, basing our discussion on the examples in [13, §11.3]. (This example is not logically essential.)

Example 1.7. Let E have basis e_1, \dots, e_d . Then $\mathrm{Sym}^2 E$ has basis

$$\{e_i^2 : 1 \leq i \leq d\} \cup \{2e_i e_j : 1 \leq i < j \leq d\}.$$

Let $X_{ii} = (e_i^2)^*$ and $X_{ij} = (2e_i e_j)^*$ be the corresponding elements of the dual space $(\mathrm{Sym}^2 E)^*$, regarded as a polynomial representation of $\mathrm{GL}(E)$ by the contravariant duality in [14, §2.7]. (Thus if $\rho(g)$ is the matrix representing g in its action on a polynomial representation V of $\mathrm{GL}(E)$ then $\rho(g^{\mathrm{tr}})^{\mathrm{tr}}$ represents g in its action on V^* .) Let \mathcal{C} be the image of E under the map $E \rightarrow \mathrm{Sym}^2 E$ defined by $v \mapsto v^2$, so

$$\mathcal{C} = \left\{ \sum_{i=1}^d \alpha_i^2 e_i^2 + \sum_{1 \leq i < j \leq d} 2\alpha_i \alpha_j e_i e_j : \alpha_1, \dots, \alpha_d \in \mathbf{C} \right\}.$$

Thinking of \mathcal{C} as an affine variety contained in $\mathrm{Sym}^2 E$, we see that the vanishing ideal of \mathcal{C} in $\mathcal{O}(\mathrm{Sym}^2 E)$ contains $X_{11}X_{22} - X_{12}^2$ in degree 2. This is a highest-weight vector for $\mathrm{GL}(E)$ of weight $(2, 2)$, so by Proposition 2.15(ii) it generates a submodule of $\mathcal{O}(\mathrm{Sym}^2 E)$ isomorphic to $\nabla^{(2,2)}(E)$. This submodule is the kernel of the map $\mathcal{O}(\mathrm{Sym}^2 E) \rightarrow \mathcal{O}(E)$ induced by restricting a coordinate function on $\mathrm{Sym}^2 E$ to \mathcal{C} and then pulling it back to E using the squaring map $E \mapsto \mathcal{C}$. Thus if $\mathcal{O}(E) = \mathbf{C}[Y_1, \dots, Y_d]$ then $X_{ij} \mapsto Y_i Y_j$. This defines a homomorphism of $\mathrm{GL}(E)$ -modules $\mathrm{Sym}^2((\mathrm{Sym}^2 E)^*) \rightarrow \mathrm{Sym}^4(E^*)$ with kernel $\nabla^{(2,2)}(E)$. Since $X_{11}^2 \in \mathcal{O}(\mathrm{Sym}^2 E)_2$ maps to $Y_1^4 \in \mathcal{O}(E)_4$, which is highest-weight of weight (4) , this $\mathrm{GL}(E)$ -homomorphism is surjective. Since all irreducible $\mathrm{GL}(E)$ -modules are self-dual under contravariant duality, we obtain

$$\mathrm{Sym}^2(\mathrm{Sym}^2 E) \cong \mathrm{Sym}^4 E \oplus \nabla^{(2,2)}(E).$$

Multiplying highest-weight vectors in the coordinate ring $\mathcal{O}(\mathrm{Sym}^2 E)$, we see that for $r, s \in \mathbf{N}_0$, the product $(X_{11}X_{22} - X_{12}^2)^r X_{11}^s$ vanishes on \mathcal{C} with multiplicity r and is highest-weight of weight $(2s + 2r, 2r)$. It follows that

$$\mathrm{Sym}^n(\mathrm{Sym}^2 E) \cong \nabla^{2n}(E) \oplus \nabla^{(2n-2,2)}(E) \oplus \nabla^{(2n-4,4)}(E) \oplus \dots \oplus W$$

where if $\nabla^\lambda(E)$ appears in W then $\ell(\lambda) \geq 3$. In particular, if $d = 2$ then $W = 0$, and every summand in the decomposition of $\mathrm{Sym}^n(\mathrm{Sym}^2 E)$ has a simple geometric interpretation. The decomposition for general d is obtained in Example 1.10 below.

In [8], the first author constructed explicit homomorphisms between modules for the symmetric group that prove the special cases of Theorem 1.3 when $\mu = (1^m)$ and $n^* = 1$ (and necessarily $\lambda^* = (1^m)$) and, subject to the conditions that m is even and λ has at most $2m$ parts, when $\mu = (1^m)$ and $\lambda^* = (1^{mn^*})$.

Theorem 1.5. This theorem strengthens the main result of [26]. The proof in [26] is entirely within the symmetric group, and constructs an explicit homomorphism corresponding to each maximal partition λ such that s_λ appears in $s_\nu \circ s_\mu$. This requires a lengthy and quite intricate argument, so again we believe that the shorter proof presented here, which also gives a precise result on the multiplicity, is of interest. As a corollary (see Corollary 7.3) we show that if T is a plethystic semistandard tableau of maximal weight for its shape, under the dominance order, then $\mathrm{wt}(T)$ is a partition. This fact was mentioned in [26], where we noted that it has a non-trivial combinatorial proof using a variation on the Bender–Knuth involution (see [2, page 47]).

Although it is not logically essential to the proof of Theorem 1.5, it is often useful in calculations (see for instance Example 1.6) that the μ -tableau entries of a plethystic semistandard tableau of shape $\mu^{(1^n)}$ and maximal weight satisfy the following closure condition.

Definition 1.8. Let \mathcal{T} be a set of semistandard μ -tableaux. We say that \mathcal{T} is *closed* if whenever $t \in \mathcal{T}$ and s is a semistandard μ -tableau obtained from t by changing a single entry c to $c - 1$, then $s \in \mathcal{T}$.

Proposition 1.9. *Let T be a plethystic semistandard tableau of shape $\mu^{(1^n)}$. If the weight of T is maximal in the dominance order for its shape then the set of μ -tableau entries of T is closed.*

Proof. Let \mathcal{T} be the set of μ -tableau entries of T . Suppose that \mathcal{T} is not closed, and let $t \in \mathcal{T}$, $s \notin \mathcal{T}$ and c be as in Definition 1.8. Let S be the plethystic semistandard tableau of shape $\mu^{(1^n)}$ obtained from T by deleting t and inserting s , and then reordering (if necessary) the μ -tableau entries within the single column of S so that S is column-standard in the total order $<$ on semistandard μ -tableau in

Definition 2.1. Then S is a plethystic semistandard tableau of shape $\mu^{(1^n)}$ and

$$\text{wt}(S)_b = \begin{cases} \text{wt}(T)_b + 1 & \text{if } b = c - 1 \\ \text{wt}(T)_b - 1 & \text{if } b = c \\ \text{wt}(T)_b & \text{otherwise.} \end{cases}$$

Therefore $\text{wt}(S) \triangleright \text{wt}(T)$. \square

Polynomial functors and highest-weight vectors. Let $d \in \mathbf{N}$ and let E be a d -dimensional complex vector space. Let V be an D -dimensional representation of $\text{GL}(E)$ corresponding, under some choice of bases of E and V , to the homomorphism $\rho : \text{GL}_d(\mathbf{C}) \rightarrow \text{GL}_D(\mathbf{C})$. Recall that V is a *polynomial representation* of degree r if for all $a, b \in \{1, \dots, D\}$ the matrix coefficient $\rho(g)_{ab}$ is a polynomial of degree r in the matrix coefficients of $g \in \text{GL}_d(\mathbf{C})$. Let $\text{GL}(E)\text{-mod}$ be the additive category of finitely generated polynomial representations of $\text{GL}(E)$ and let $\text{GL}(E)\text{-mod}_r$ be its full subcategory of representations of polynomial degree r . For each $\lambda \in \text{Par}(r)$ let $\nabla^\lambda : \text{GL}(E)\text{-mod}_p \rightarrow \text{GL}(E)\text{-mod}_{pr}$ be the Schur functor, as defined in §2.3. (Our construction and proofs have some novel features, but this section will be background for most readers.)

By Proposition 2.15 every polynomial representation $V \in \text{GL}(E)\text{-mod}$ decomposes as a direct sum of submodules each isomorphic to some $\nabla^\lambda(E)$. Let $[W : \nabla^\lambda(E)]$ denotes the number of irreducible summands of the polynomial representation W that are isomorphic to $\nabla^\lambda(E)$. By Proposition 3.3, composition of Schur functors corresponds to plethysm of Schur functions. Hence, by Proposition 2.15(iv),

$$(1.7) \quad \langle s_\nu \circ s_\mu, s_\lambda \rangle = [\nabla^\nu(\nabla^\mu E) : \nabla^\lambda E]$$

for all partitions λ, μ and ν with at most d parts. Thus each of the main theorems has an equivalent restatement as a result on polynomial representations of $\text{GL}(E)$. To prove these restatements, we use the model for $\nabla^\nu(\nabla^\mu(E))$ constructed in §3 and the following key fact about highest-weight vectors, as characterized in Lemma 2.13 using the Lie algebra action of $\mathfrak{gl}(E)$, and proved in Proposition 2.15: *if V is a polynomial $\text{GL}(E)$ -module then V contains a highest-weight vector v ; moreover, if the weight of v is λ then the submodule of V generated by v is isomorphic to $\nabla^\lambda(E)$.*

To illustrate the power of this property we end this introduction by giving a very short proof that if $\dim E \geq n$ then

$$(1.8) \quad \text{Sym}^n(\text{Sym}^2 E) \cong \bigoplus_{\lambda \in \text{Par}(n)} \nabla^{2\lambda}(E).$$

By much more lengthy arguments, Boffi shows in [3] that, over an arbitrary field, $\text{Sym}^n \text{Sym}^2$ has a filtration by the functors ∇^λ ; he reports that this result had previously been obtained in [1]. The analogous result for the symmetric group was proved independently by the second author in [24]. Boffi's result was generalized to arbitrary commutative rings in [7].

Example 1.10. Let E have basis e_1, \dots, e_n . For each $\ell \in \{1, \dots, n\}$ and each function $\sigma : \{1, \dots, \ell\} \rightarrow \{1, \dots, \ell\}$, define

$$w(\sigma) = (e_1 e_{1\sigma}) \dots (e_\ell e_{\ell\sigma}) \in \text{Sym}^\ell(\text{Sym}^2 E).$$

Let $w_\ell = \sum_{\sigma \in S_\ell} w(\sigma) \text{sgn}(\sigma)$. Since the coefficient of $e_1^2 \dots e_\ell^2$ in w_ℓ is 1, we have $w_\ell \neq 0$. Let $X^{(c)} \in \mathfrak{gl}(E)$ be defined, as in Lemma 2.13, by $X^{(c)} \cdot e_c = e_{c-1}$ and $X^{(c)} \cdot e_b = 0$ if $b \neq c$. Define $\delta : \{1, \dots, \ell\} \rightarrow \{1, \dots, \ell\}$ by $\delta(c) = c - 1$ and $\delta(b) = b$ if $b \neq c$. Thus $\delta(b) = c - 1$ if and only if $b \in \{c - 1, c\}$. The action of $\mathfrak{gl}(E)$ on symmetric powers is recalled immediately before the proof of Proposition 2.14. Using this action and $w(\sigma) = w(\sigma^{-1})$, we find that

$$X^{(c)} \cdot w(\sigma) = (e_{c-1} e_{c\sigma}) \prod_{i \neq c} (e_i e_{i\sigma}) + (e_{c\sigma-1} e_{c-1}) \prod_{j \neq c} (e_j e_{j\sigma^{-1}}).$$

The summands are the products of all $e_{b\sigma^{-1}\delta} e_b$ and $e_b e_{b\sigma\delta}$, respectively. Hence

$$X^{(c)} \cdot w(\sigma) = w(\sigma^{-1}\delta) + w(\sigma\delta).$$

Let τ be the transposition $(c - 1, c)$. Using that $\tau\delta = \delta$ we get

$$\sum_{\sigma \in S_\ell} w(\sigma\delta) \text{sgn}(\sigma) = \sum_{\sigma' \in S_\ell} w(\sigma'\tau\delta) \text{sgn}(\sigma'\tau) = - \sum_{\sigma' \in S_\ell} w(\sigma'\delta) \text{sgn}(\sigma')$$

and so $\sum_{\sigma \in S_\ell} w(\sigma\delta) \text{sgn}(\sigma) = 0$. Similarly $\sum_{\sigma \in S_\ell} w(\sigma^{-1}\delta) \text{sgn}(\sigma) = 0$, and so $X^{(c)} \cdot w_\ell = 0$. (This cancellation has an attractive combinatorial interpretation, shown in Figure 1 overleaf.) By Lemma 2.13, $w_\ell \in \text{Sym}^\ell \text{Sym}^2 E$ is a highest-weight vector of weight (2^ℓ) . Now take $\lambda \in \text{Par}(n)$. Multiplying highest-weight vectors shows that if k is the first part of λ then $w(\lambda_1) \dots w(\lambda_k) \in \text{Sym}^n(\text{Sym}^2 E)$ is a highest-weight vector of weight 2λ . By the key property $\text{Sym}^n(\text{Sym}^2 E)$ has $\nabla^{2\lambda}(E)$ as a summand.

We use the Frobenius–Schur involution count to show that there are no further summands. For $\lambda \in \text{Par}(r)$, let χ^λ be the irreducible character of the symmetric group S_r corresponding to $\nabla^\lambda(E)$ (and to the Schur function s_λ). By the involution count, $\sum_{\lambda \in \text{Par}(r)} \chi^\lambda(1) = T_r$, where T_r is the number of permutations of order at most two in S_r . On the other hand, setting

$$M_k = (\text{Sym}^n(\text{Sym}^2 E)) \otimes \bigwedge^k E,$$

it follows from Pieri’s rule and the result already proved that M_k contains $\bigoplus_\lambda \nabla^\lambda(E)$, where the sum is over all partitions $\lambda \in \text{Par}(r)$ having exactly k odd parts. By (2.2) in §2.2 below, the character of the symmetric group S_r corresponding to $\bigoplus_{2n+k=r} M_k$ is $\sum_{2n+k=r} (1_{C_{2\ell} S_n} \times \text{sgn}_{S_k}) \uparrow_{C_{2\ell} S_n \times S_k}^{S_r}$. For $n \geq 1$, the degree of $(1_{C_{2\ell} S_n} \times \text{sgn}_{S_k}) \uparrow_{C_{2\ell} S_n \times S_k}^{S_r}$ is the number of involutions in S_{2n} having precisely k fixed points. Therefore this character has degree T_r and so $\bigoplus_{2n+k=r} M_k \cong \bigoplus_{\lambda \in \text{Par}(r)} \nabla^\lambda(E)$ and $M_0 = \text{Sym}^n(\text{Sym}^2 E)$ has no summands other than those already found.




FIGURE 1. Example 1.10 for $\ell = 3$. Writing permutations $\sigma \in S_3$ in one-line form, the six summands in w_3 are $w(123)$, $w(213)$, $w(231)$, $w(321)$, $w(312)$, $w(132)$, clockwise from the top. Identifying 1 and 2 in the image of σ (shown as the bottom row in the permutation diagrams) the contributions from σ and $\sigma(12)$ cancel. Cancelling pairs are connected by edges marked (12). Thus $\sum_{\sigma \in S_3} w(\sigma\delta) \operatorname{sgn}(\sigma) = 0$.

2. BACKGROUND

2.1. Partitions and tableaux. Let λ be a partition of $r \in \mathbf{N}$. Recall that $\ell(\lambda)$ denotes the number of parts of λ . The *Young diagram* of λ is the set $[\lambda] = \{(i, j) : 1 \leq i \leq \ell(\lambda), 1 \leq j \leq \lambda_i\}$. Let \mathcal{B} be a set. A λ -*tableau with entries from \mathcal{B}* is a function $t : [\lambda] \rightarrow \mathcal{B}$. We write the image of (i, j) under t as $t_{(i,j)}$. If $t_{(i,j)} = b \in \mathcal{B}$ then we say that b is the *entry* of t in row i and column j . When \mathcal{B} is a set of natural numbers, this corresponds to the usual diagrammatic representation of tableaux (see §2.3 for a small example).

Now suppose that \mathcal{B} is totally ordered by an order denoted $<$. We say that a tableau t with entries from \mathcal{B} is *row-semistandard* if its rows are weakly increasing from left to right, *column-standard* if its columns are strictly increasing from top to bottom, both under the order $<$. We say that t is *semistandard* if it is both row-semistandard and column-standard. The terms *row-standard* and *standard* are defined analogously, requiring in addition that the rows are strictly increasing. Let $\operatorname{SSYT}_{\mathcal{B}}(\lambda)$ denote the set of semistandard λ -tableaux with entries from \mathcal{B} . When

$\mathcal{B} \subseteq \mathbf{N}$, we refer to elements of $\text{SSYT}_{\mathcal{B}}(\lambda)$ simply as semistandard λ -tableaux. If $\mathcal{B} = \{1, \dots, d\}$ and t has exactly β_b entries for each $b \in \{1, \dots, d\}$, then we say that t has *content* β . Let $\text{SSYT}(\lambda, \beta)$ denote the set of semistandard λ -tableaux of content β . Let t^λ be the unique element of $\text{SSYT}(\lambda, \lambda)$, as defined by $t_{(i,j)}^\lambda = i$ for each $(i, j) \in [\lambda]$. In the total order on column-standard λ -tableau defined below, t^λ is the least element.

Definition 2.1. Given column-standard λ -tableaux t and u with entries in a totally ordered set, set $t < u$ if and only if in the rightmost column that differs between t and u , the greatest entry not appearing in both columns lies in u .

In particular, if μ and ν are partitions then the semistandard μ -tableaux are totally ordered by $<$. A plethystic semistandard tableau of shape μ' , as defined in Definition 1.4, is a semistandard ν -tableau with entries from $\text{SSYT}_{\mathbf{N}}(\mu)$.

The symmetric group on $[\lambda]$ acts on λ -tableaux on the right by place permutation: thus if $\sigma \in S_{[\lambda]}$ and t is a λ -tableau then $(t\sigma)_{(i,j)} = t_{(i,j)\sigma^{-1}}$. Note that the entry in position (i, j) of t is found in position $(i\sigma, j\sigma)$ of $t\sigma$. Let $\text{CPP}(\lambda)$ denote the Young subgroup of $S_{[\lambda]}$ having as its orbits the columns of $[\lambda]$.

2.2. Schur functions and plethysm. Let Λ be the ring of symmetric functions as defined in [27, §7]. Let $\lambda \in \text{Par}(r)$. Given a λ -tableau t of content β , let x^t denote the monomial $x_1^{\beta_1} \dots x_{\ell(\beta)}^{\beta_{\ell(\beta)}}$. The *Schur function* $s_\lambda \in \Lambda$ is defined by $s_\lambda = \sum_{t \in \text{SSYT}_{\mathbf{N}}(\lambda)} x^t$. By [27, Theorem 7.10.2], s_λ is a symmetric function. Thus

$$(2.1) \quad s_\lambda = \sum_{\kappa \in \text{Par}(r)} |\text{SSYT}(\lambda, \kappa)| \text{mon}_\kappa$$

where $\text{mon}_\kappa = x_1^{\kappa_1} \dots x_{\ell(\kappa)}^{\kappa_{\ell(\kappa)}} + \dots$ denotes the monomial symmetric function corresponding to κ . For example, $s_{(2)} = x_1^2 + x_2^2 + \dots + x_1x_2 + x_1x_3 + x_2x_3 + \dots = \text{mon}_{(2)} + \text{mon}_{(1,1)}$.

Given $f(x_1, x_2, \dots) \in \Lambda$, the plethysm $f \circ s_\lambda$ may be defined by substituting the monomials x^t , where t ranges over all semistandard λ -tableaux with entries from \mathbf{N} , for the variables x_1, x_2, \dots of f . The general definition of $f \circ g$ is given in [27, Ch. 7, Appendix 2], [22, Ch. 1, Appendix A] or [21]. For example, the combinatorial analogue of the case $n = 2$ of (1.8) is

$$s_{(2)} \circ s_{(2)} = s_{(2)}(x_1^2, x_1x_2, \dots) = \text{mon}_{(4)} + \text{mon}_{(3,1)} + 2\text{mon}_{(2,2)} = s_{(4)} + s_{(2,2)}.$$

The coefficient of $\text{mon}_{(2,2)}$ is 2 since $x_1^2x_2^2$ may be obtained as both $(x_1^2)(x_2^2)$ and $(x_1x_2)^2$ when multiplying out $s_{(2)}(x_1^2, x_1x_2, \dots)$.

There is an involutory ring homomorphism $\omega : \Lambda \rightarrow \Lambda$ defined by $\omega(s_\lambda) = s_{\lambda'}$. We call ω the *sign twist*. Its effect on plethysms is as follows.

Lemma 2.2. *Let μ and ν be partitions. If μ is a partition of m then*

$$\omega(s_\nu \circ s_\mu) = \begin{cases} s_\nu \circ s_{\mu'} & \text{if } m \text{ is even} \\ s_{\nu'} \circ s_{\mu'} & \text{if } m \text{ is odd.} \end{cases}$$

Proof. See [22, Ch. I, Equation (2.7)]. □

In Examples 1.6 and 1.10 we used the characteristic isometry, which sends the irreducible character χ^λ of S_r to the Schur function s_λ where $\lambda \in \text{Par}(r)$. By [27, A2.8] or [22, Ch. 1, Appendix A, (6.2)], if $\mu \in \text{Par}(m)$ and $\nu \in \text{Par}(n)$ then under this isometry,

$$(2.2) \quad ((\chi^\mu)^{\widetilde{\times} n} \text{Inf}_{S_n}^{S_m \wr S_n} \chi^\nu) \uparrow_{S_m \wr S_n}^{S_{mn}} \mapsto s_\nu \circ s_\mu$$

where $(\chi^\mu)^{\widetilde{\times} n}$ is the character of the irreducible representation of $S_m \wr S_n$ defined in [18, (4.3.8)] and $\text{Inf}_{S_n}^{S_m \wr S_n}$ is the inflation map. Thus all the main theorems have immediate translations into results on characters of symmetric groups. We shall not use the characteristic isometry any further below.

2.3. Polynomial representations of $\text{GL}_n(K)$. In the following three subsections we construct the Schur functors ∇^λ used in the proofs of the main theorems. We end with Remark 2.16 which explains the connection with an earlier construction due to James [19, Ch. 26]. Let K be a commutative ring, let \mathcal{B} be a totally ordered set as in §2.1, and let V be a free K -module with basis $\{v_b : b \in \mathcal{B}\}$. Except in the proof of Lemma 2.4, the case $K = \mathbf{C}$ suffices. The set \mathcal{B} is either $\{1, \dots, d\}$ or, when we compose Schur functors, the set of semistandard tableaux with entries from $\{1, \dots, d\}$, with the order from Definition 2.1.

Fix $\lambda \in \text{Par}(r)$. Recall that $\text{CPP}(\lambda)$ is defined after Definition 2.1.

Definition 2.3. Let $\text{Sym}^\lambda V = \bigotimes_{i=1}^{\ell(\lambda)} \text{Sym}^{\lambda_i} V$. Given a λ -tableau t with entries from \mathcal{B} , the *GL-tabloid* corresponding to t is the element $f(t) \in \text{Sym}^\lambda V$ defined by

$$f(t) = \bigotimes_{i=1}^{\ell(\lambda)} \prod_{j=1}^{\lambda_i} v_{t_{(i,j)}} \in \text{Sym}^\lambda V.$$

The *GL-polytabloid* corresponding to t is

$$F(t) = \sum_{\sigma \in \text{CPP}(\lambda)} f(t\sigma) \text{sgn}(\sigma) \in \text{Sym}^\lambda V.$$

We define $\nabla^\lambda(V)$ to be the K -submodule of $\text{Sym}^\lambda V$ spanned by the GL-polytabloids $F(t)$ for t a λ -tableau with entries from \mathcal{B} .

Since $f(t) = f(t')$ if and only if the rows of t and t' are equal as multisets,

$$(2.3) \quad \{f(t) : t \text{ a row-semistandard } \lambda\text{-tableau with entries from } \mathcal{B}\}$$

is a basis of $\text{Sym}^\lambda(V)$. It is also useful to note that if $\sigma \in \text{CPP}(\lambda)$ then

$$(2.4) \quad F(t\sigma) = F(t) \text{sgn}(\sigma).$$

In particular, $F(t) = 0$ if t has a column with a repeated entry, and so $\nabla^\lambda(V) = 0$ if $\ell(\lambda) > \dim V$, as used in Example 1.7. A potential trap is that $F(t)$ depends on the tableau t , not just on the GL-tabloid $f(t)$. For example, if $\mathcal{B} = \{1, 2, 3\}$ and

$$t = \begin{array}{|c|c|} \hline 1 & 3 \\ \hline 2 & 1 \\ \hline \end{array}, \quad t' = \begin{array}{|c|c|} \hline 1 & 3 \\ \hline 1 & 2 \\ \hline \end{array}$$

then $f(t) = f(t')$ but $F(t') = 0$ whereas $F(t) = v_1 v_3 \otimes v_2 v_1 - v_2 v_3 \otimes v_1^2 - v_1^2 \otimes v_2 v_3 + v_2 v_1 \otimes v_1 v_3 \neq 0$. It is clear that $\nabla^\lambda(V)$ is functorial in V , so ∇^λ is an endofunctor of the category of free K -modules of finite rank.

Postponing the action of the general linear group for the moment, we find an explicit basis for $\nabla^\lambda(V)$, introducing two results that are critical to the proofs of the main theorems. The following lemma is the analogue for GL-polytabloids of part of the proof of Theorem 8.4 in [19]. There are some subtle differences between the proofs because of our use of place permutations.

Lemma 2.4. *Let $1 \leq j < j' < \lambda_1$ and let $1 \leq i \leq \lambda'_j$. Define subsets of $[\lambda]$ by $A_\lambda(i, j) = \{(i, j), \dots, (\lambda'_j, j)\}$ and $B_\lambda(i, j') = \{(1, j'), \dots, (i, j')\}$. If t is a λ -tableau then*

$$\sum_{\tau} F(t\tau) \operatorname{sgn}(\tau) = 0$$

where the sum is over all $\tau \in S_{A_\lambda(i, j) \cup B_\lambda(i, j')}$.

Proof. Let A and B denote $A_\lambda(i, j)$ and $B_\lambda(i, j')$, respectively. By definition, the left-hand side is $\sum_{\tau \in S_{A \cup B}} \sum_{\sigma \in \operatorname{CPP}(\lambda)} f((t\tau)\sigma) \operatorname{sgn}(\tau) \operatorname{sgn}(\sigma)$. Therefore it suffices to show that $\sum_{\tau \in S_{A \cup B}} f((t\tau)\sigma) \operatorname{sgn}(\tau) = 0$ for each $\sigma \in \operatorname{CPP}(\lambda)$. Since $|A \cup B| = \lambda'_j + 1$, there exist boxes $(h, j)\sigma \in A\sigma$ and $(h', j')\sigma \in B\sigma$ such that $(h, j)\sigma$ and $(h', j')\sigma$ are in the same row of $[\lambda]$. Let $\delta = ((h, j)\sigma, (h', j')\sigma) \in S_{A\sigma \cup B\sigma}$ be the transposition swapping these boxes. Let $\vartheta_1, \dots, \vartheta_\ell$ be representatives for the left cosets of $\langle \delta \rangle$ in $S_{A\sigma \cup B\sigma}$. Thus $S_{A\sigma \cup B\sigma} = \bigcup_{c=1}^{\ell} \vartheta_c \langle \delta \rangle$ and

$$\begin{aligned} \sum_{\tau \in S_{A \cup B}} f((t\tau)\sigma) \operatorname{sgn}(\tau) &= \sum_{\tau \in S_{A \cup B}} f((t\sigma)\tau^\sigma) \operatorname{sgn}(\tau) \\ &= \sum_{\tau^* \in S_{A\sigma \cup B\sigma}} f((t\sigma\tau^*) \operatorname{sgn}(\tau^*)) \\ &= \sum_{c=1}^{\ell} (f(t\sigma\vartheta_c) - f(t\sigma\vartheta_c\delta)) \operatorname{sgn}(\vartheta_c) \\ &= 0 \end{aligned}$$

where the final equality holds because δ swaps two boxes in the same row of $[\lambda]$, and so the tableaux $t\sigma\vartheta_h$ and $(t\sigma\vartheta_h)\delta$ have equal rows. \square

If $\tau \in S_{A_\lambda(i, j)} \times S_{B_\lambda(i, j')}$ then, by (2.4), $F(t\tau) \operatorname{sgn}(\tau) = F(t)$. Let ϕ_1, \dots, ϕ_ℓ be representatives for the left cosets of $S_{A_\lambda(i, j)} \times S_{B_\lambda(i, j')}$ in S_C , where $C = A_\lambda(i, j) \cup B_\lambda(i, j')$, chosen so that $\phi_1 = \operatorname{id}$. Thus $S_C = \bigcup_{c=1}^{\ell} \phi_c (S_{A_\lambda(i, j)} \times S_{B_\lambda(i, j')})$. By (2.4), $F(t\phi_c\tau) = F(t\phi_c) \operatorname{sgn}(\tau)$ for each c and each $\tau \in S_{A_\lambda(i, j)} \times S_{B_\lambda(i, j')}$. Therefore Lemma 2.4 implies that

$$|A_\lambda(i, j)|! |B_\lambda(i, j')|! \sum_{c=1}^{\ell} F(t\phi_c) \operatorname{sgn}(\phi_c) = 0.$$

When $K = \mathbf{Z}$ we may cancel the factorials since $\nabla^\lambda(V)$ is a submodule of the free \mathbf{Z} -module $\text{Sym}^\lambda(V)$. Thus the relation

$$(2.5) \quad F(t) = - \sum_{c=2}^{\ell} F(t\phi_c) \text{sgn}(\phi_c).$$

holds over an arbitrary commutative ring K . We call (2.5) a *snake relation*, because of the shape formed by the boxes in $A_\lambda(i, j) \cup B_\lambda(i, j')$ when $j' = j + 1$. It is critical to the proofs of Theorem 1.1, 1.2 and 1.5.

It is convenient to choose the coset representatives ϕ_1, \dots, ϕ_ℓ so that each ϕ_c is a product of transpositions swapping boxes in $A_\lambda(i, j)$ and $B_\lambda(i, j)$, preserving the relative vertical order of boxes in each set.

Example 2.5. The snake relation for $A_{(2,2,1)}(2, 1) = \{(2, 1), (3, 1)\}$ and $B_{(2,2,1)}(2, 2) = \{(1, 2), (2, 2)\}$ has five summands on its right-hand side. Some of these summands may vanish. For example

$$F \left(\begin{array}{|c|c|} \hline 1 & 1 \\ \hline 3 & 2 \\ \hline 4 & \\ \hline \end{array} \right) = F \left(\begin{array}{|c|c|} \hline 1 & 1 \\ \hline 2 & 3 \\ \hline 4 & \\ \hline \end{array} \right) + F \left(\begin{array}{|c|c|} \hline 1 & 1 \\ \hline 3 & 4 \\ \hline 2 & \\ \hline \end{array} \right)$$

because, taking coset representatives as suggested above, the tableaux obtained by the transpositions $(1, 2) \leftrightarrow (2, 1)$ and $(1, 2) \leftrightarrow (3, 1)$, and the double transposition $(1, 2) \leftrightarrow (2, 1), (2, 2) \leftrightarrow (3, 1)$ have a repeated 1 in their first column.

By (2.4), applying the transposition $(2, 1) \leftrightarrow (3, 1)$ to the second summand switches its sign and expresses the left-hand side as a linear combination of semistandard GL-polytabloids.

Corollary 2.6. *If t is a λ -tableau with entries from \mathcal{B} then $F(t)$ may be expressed as a K -linear combination of GL-polytabloids $F(s)$ for semistandard λ -tableaux s by applying finitely many snake relations.*

Proof. By (2.4), we may assume that t is column-standard. If t is not standard then there exist $(i, j), (i, j + 1) \in [\lambda]$ such that $t_{(i,j)} > t_{(i,j+1)}$. Let $A, B \subseteq [\lambda]$ be as in Lemma 2.4, taking $j' = j + 1$. By (2.5), $F(t) = - \sum_{c=2}^{\ell} F(t\phi_c) \text{sgn}(\phi_c)$ where each ϕ_c swaps certain boxes in A with certain boxes, necessarily having smaller entries, in B . Thus $t < t\phi_c$ for each ϕ_c , where $<$ refers to the order in Definition 2.1. The result now follows by induction. \square

We say that $F(t)$ is *straightened* by snake relations. A related result to (2.5) gives some control over the $F(s)$ that may appear in the straightening of $F(t)$; it is needed in the proof of Theorem 1.2. To state it, we require two further orders.

Define a *composition* of $n \in \mathbf{N}_0$ of length ℓ to be an element $\beta \in \mathbf{N}_0^\ell$ such that $\sum_{i=1}^{\ell} \beta_i = n$. We set $\ell(\beta) = \ell$.

Definition 2.7. If β and γ are compositions of n , we say that β *dominates* γ , and write $\beta \succeq \gamma$ if $\beta_1 + \dots + \beta_i \geq \gamma_1 + \dots + \gamma_i$ for all $i \in \mathbf{N}$. (As usual, if i exceeds the number of parts of β or γ then the corresponding part is taken to be 0.) Given

a row-semistandard tableau t with entries from $\{1, \dots, d\}$, and $b \in \{1, \dots, d\}$, let $t^{\leq b}$ be the composition γ defined by $\gamma_i = |\{j : 1 \leq j \leq \lambda_i, t_{(i,j)} \leq b\}|$. If u is a row-semistandard tableau with entries also with entries from $\{1, \dots, d\}$ and of the same shape as t , we say that t *dominates* u , and write $t \triangleright u$ if $t^{\leq b} \triangleright u^{\leq b}$ for all $b \in \{1, \dots, d\}$. We extend this order to tableaux with entries from an arbitrary totally ordered set \mathcal{B} by the unique order-preserving bijection between $\{1, \dots, |\mathcal{B}|\}$ and \mathcal{B} .

Definition 2.8. Given a tableau t with entries from \mathbf{N} , let \bar{t} be the row-semistandard tableau obtained by sorting the rows of t into non-decreasing order.

Proposition 2.9. *Let t be a column-standard λ -tableau with entries from $\{1, \dots, d\}$. Then \bar{t} is semistandard and*

$$F(t) = F(\bar{t}) + w$$

where w is an integral linear combination of GL-polytabloids $F(s)$ for semistandard λ -tableaux s such that $\bar{t} \triangleright s$.

Proof. We reduce to the analogous result for tableaux with distinct entries proved in [29, Proposition 4.1]. Let t have content β . Let t_\star be the tableau obtained from t by replacing each of the β_b entries of t equal to b with a *symbol* $b^{(1)}, \dots, b^{(\beta_b)}$, for each $b \in \{1, \dots, d\}$. We say that $b^{(i)}$ has *number* b and *exponent* i . We order symbols lexicographically, by number then exponent, so $b^{(i)} < c^{(j)}$ if and only if $b < c$ or $b = c$ and $i < j$. Thus t_\star has distinct entries and is column-standard. By Proposition 4.1 of [29], \bar{t}_\star is standard. Let V_\star be the free K -module with basis vectors $v_{b^{(i)}}$ in bijection with symbols. Again by [29], now working in $\nabla^\lambda(V_\star)$, we have $F(t_\star) = F(\bar{t}_\star) + w$ where w is an integral linear combination of GL-polytabloids $F(s)$ for standard λ -tableaux s (each having symbol entries). The proposition now follows from functoriality: the quotient map $V_\star \rightarrow V$ sending $v_{b^{(i)}}$ to v_b for each symbol $b^{(i)}$ corresponds to replacing each symbol with its number. \square

Corollary 2.6 also does most of the work to prove a well-known basis theorem for $\nabla^\lambda(V)$. We include the details since the following lemma is also needed in the proof of Theorem 1.2. As a notational convenience, we extend the dominance order \triangleright on row-semistandard tableaux to GL-tabloids by setting $f(t) \triangleright f(u)$ if and only if $\bar{t} \triangleright \bar{u}$.

Lemma 2.10. *Let t be a column-standard λ -tableau with entries from \mathcal{B} . Let $F(t) = f(t) + w$ where $w \in \text{Sym}^\lambda(V)$. If u is a row-semistandard λ -tableau such that $f(u)$ appears with non-zero coefficient when w is written in the canonical basis (2.3) of $\text{Sym}^\lambda(V)$, then $f(t) \triangleright f(u)$.*

Proof. By permuting the rows of u we may assume that $u = t\tau$ for some non-identity permutation $\tau \in \text{CPP}(\lambda)$. Since t is column-standard, there exist boxes $(i, j), (i', j)$ of λ such that $i < i'$ and $u_{(i,j)} > u_{(i',j)}$. Say that two such boxes form a *column inversion*. Let $c = u_{(i,j)}$, let $c' = u_{(i',j)}$ and let $u^\star = u((i, j), (i', j))$. It

is easily seen that if $c' \leq b < c$ then the Young diagrams of $u^{\leq b}$ and $u^{\star \leq b}$ differ by a single box, moved from row i' up to row i . For all other b , we have $u^{\leq b} = u^{\star \leq b}$. Hence $\bar{u}^{\star} \triangleright \bar{u}$. By induction on the number of column inversions, $\bar{t} \triangleright \bar{u}^{\star}$. Hence $\bar{t} \triangleright \bar{u}$ as required. \square

We note this proof is essentially the same as that of Lemma 8.3 in [19]. It is also possible to reduce to this result by distinguishing equal entries of t by formal symbols, as in the proof of Proposition 2.9.

Proposition 2.11. *The set $\{F(s) : s \in \text{SSYT}_{\mathcal{B}}(\lambda)\}$ is a K -basis for $\nabla^{\lambda}(V)$.*

Proof. By Corollary 2.6, $\nabla^{\lambda}(V)$ is spanned by $\{F(s) : s \in \text{SSYT}_{\mathcal{B}}(\lambda)\}$. Let $v = \sum_{s \in \text{SSYT}_{\mathcal{B}}(\lambda)} \alpha_s F(s)$ where not every coefficient is zero. Take s maximal in the dominance order on semistandard λ -tableaux such that $\alpha_s \neq 0$. By Lemma 2.10, the coefficient of $f(s)$ in $F(s)$ is 1. Again by this lemma, if s' is a semistandard λ -tableau other than s such that $\alpha_{s'} \neq 0$, then $f(s') \triangleright f(t')$ for every $f(t')$ appearing in $F(s')$. Hence, by maximality of s , we have $f(t') \not\triangleright f(s)$. Therefore the coefficient of $f(s)$ in v is α_s , and so $v \neq 0$. \square

2.4. Action of $\text{GL}(E)$. We now suppose that K is an infinite field. Thus, with the notation from the previous section, V is a K -vector space with basis $\{v_b : b \in \mathcal{B}\}$. Let E be a K -vector space and suppose that V is a polynomial $\text{GL}(E)$ -module. Let $t \in \text{SSYT}_{\mathcal{B}}(\lambda)$. The action of $g \in \text{GL}(E)$ on $F(t) \in \nabla^{\lambda}(V)$, where t is a λ -tableau with entries from \mathcal{B} , is determined by the multilinear construction in Definition 2.3. The following method is convenient in calculations: formally replace each entry b in t with gv_b , expressed as a K -linear combination of $\{v_b : b \in \mathcal{B}\}$, and then expand multilinearly. For example, suppose that E is 3-dimensional and V is the natural representation of E , so we take $\mathcal{B} = \{1, 2, 3\}$. Let

$$g = \begin{pmatrix} \alpha & 0 & 0 \\ \delta & \beta & 0 \\ \varepsilon & 0 & \gamma \end{pmatrix} \in \text{GL}(E).$$

Then in its action on $\nabla^{(2,2)}(V)$ we have

$$\begin{aligned} gF\left(\begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & 1 \\ \hline \end{array}\right) &= F\left(\begin{array}{|c|c|} \hline \alpha v_1 + \delta v_2 + \varepsilon v_3 & \beta v_2 \\ \hline \gamma v_3 & \alpha v_1 + \delta v_2 + \varepsilon v_3 \\ \hline \end{array}\right) \\ &= \alpha^2 \beta \gamma F\left(\begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & 1 \\ \hline \end{array}\right) + \alpha \beta \gamma \delta F\left(\begin{array}{|c|c|} \hline 2 & 2 \\ \hline 3 & 1 \\ \hline \end{array}\right) \\ &\quad + \alpha \beta \gamma \varepsilon F\left(\begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & 3 \\ \hline \end{array}\right) + \beta \gamma \delta \varepsilon F\left(\begin{array}{|c|c|} \hline 2 & 2 \\ \hline 3 & 3 \\ \hline \end{array}\right) \end{aligned}$$

where the first line should be interpreted entirely formally. One may then use snake relations to express the right-hand side in the standard basis of $\nabla^{(2,2)}(V)$.

In this example, V had polynomial degree 1. In general, if the $\text{GL}(E)$ -module V has degree p , then, identifying $\text{GL}(E)$ with $\text{GL}_d(\mathbf{C})$, the coefficients in $gF(t)$ have degree pr in the matrix coefficients of $g \in \text{GL}_d(\mathbf{C})$. Thus if $|\lambda| = r$ then ∇^{λ} :

$\mathrm{GL}(E)\text{-mod}_p \rightarrow \mathrm{GL}(E)\text{-mod}_{pr}$ is an endofunctor of $\mathrm{GL}(E)\text{-mod}$. (This was seen for the decomposition of $\mathrm{Sym}^n(\mathrm{Sym}^2 E)$ in Example 1.7.) For a further example, if $\dim E = 2$, $V = \mathrm{Sym}^2 E$, $\lambda = (2, 1)$ and $v_1 = e_1^2, v_2 = e_1 e_2, v_3 = e_2^2$ then working in $\nabla^{(2,1)}(\mathrm{Sym}^2(E))$ we have

$$\begin{aligned} & \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} F \left(\begin{array}{|c|c|} \hline 1 & 3 \\ \hline 2 & \\ \hline \end{array} \right) \\ &= F \left(\begin{array}{|c|c|} \hline \alpha^2 v_1 + 2\alpha\gamma v_2 + \gamma^2 v_3 & \beta^2 v_1 + 2\beta\delta v_2 + \delta^2 v_3 \\ \hline \alpha\gamma v_1 + (\alpha\delta + \beta\gamma)v_2 + \beta\delta v_3 & \\ \hline \end{array} \right) \\ &= \alpha^2 \beta^2 (\alpha\delta + \beta\gamma) F \left(\begin{array}{|c|c|} \hline 1 & 1 \\ \hline 2 & \\ \hline \end{array} \right) + 2\alpha^2 \beta\delta (\alpha\delta + \beta\gamma) F \left(\begin{array}{|c|c|} \hline 1 & 2 \\ \hline 2 & \\ \hline \end{array} \right) \\ &\quad + \alpha^2 \delta^2 (\alpha\delta + \beta\gamma) F \left(\begin{array}{|c|c|} \hline 1 & 3 \\ \hline 2 & \\ \hline \end{array} \right) + \dots \end{aligned}$$

where the coefficients have degree 6.

2.5. Action of $\mathfrak{gl}(E)$ and highest-weight vectors. As in the previous subsection, let K be an infinite field and let E be a K -vector space. Let $\dim E = d$. Recall that if V is a polynomial representation of $\mathrm{GL}(E)$ and β is a composition with $\ell(\beta) = d$ then a non-zero vector $v \in V$ is a *weight* vector of weight β if

$$(2.6) \quad gv = g_{11}^{\beta_1} \dots g_{dd}^{\beta_d} v$$

for all diagonal matrices $g \in \mathrm{GL}(E)$. Let V_β be the subspace of V of weight vectors of weight β , together with 0. The *formal character* of a polynomial representation V of $\mathrm{GL}(E)$ is the polynomial

$$\Phi_V(x_1, \dots, x_d) = \sum_{\beta} \dim(V_\beta) x_1^{\beta_1} \dots x_d^{\beta_d}$$

where the sum is over all compositions β such that $\ell(\beta) = d$.

Lemma 2.12.

- (i) If $s \in \mathrm{SSYT}(\lambda, \beta)$ then $F(s) \in \nabla^\lambda(E)$ has weight β .
- (ii) The formal character of $\nabla^\lambda(E)$ is $s_\lambda(x_1, \dots, x_d)$.

Proof. If $g \in \mathrm{GL}(E)$ is a diagonal matrix then $gf(s) = g_{11}^{\beta_1} \dots g_{dd}^{\beta_d} f(s)$. Since all the λ -tableaux appearing in $F(s)$ have the same content as s , it follows that $F(s)$ has weight β , proving (i). Part (ii) follows from (i), Proposition 2.11 and the definition of Schur functions (see (2.1) in §2.2). \square

In the proofs of the main theorems, it is easiest to consider $\nabla^\lambda(E)$ as a module for the Lie algebra $\mathfrak{gl}(E)$ of $\mathrm{GL}(E)$. Let \mathfrak{b} be the Borel subalgebra of $\mathfrak{gl}(E)$ of upper-triangular matrices. Let β be a composition of n with $\ell(\beta) = d$. Recall that if V is a $\mathfrak{gl}(E)$ -module then $v \in V$ is a *weight vector* of weight β if

$$X \cdot v = (\beta_1 X_{11} + \dots + \beta_d X_{dd})v$$

for all diagonal matrices $X \in \mathfrak{gl}(E)$. If this equation holds for all diagonal $X \in \mathfrak{b}$, and $X \cdot v = 0$ whenever X is strictly upper-triangular, then we say that v is a *highest-weight vector* of weight β . For $c \in \{2, \dots, d\}$, let $X^{(c)} \in \mathfrak{b}$ be the strictly upper-triangular matrix having 1 in position $(c-1, c)$ and 0 in all other positions. Since the Lie subalgebra of \mathfrak{b} of strictly upper-triangular matrices is generated by $X^{(2)}, \dots, X^{(c)}$, we have the following lemma.

Lemma 2.13. *Let V be a $\mathfrak{gl}(E)$ -module. The vector $v \in V$ is a highest-weight vector of weight β if and only if $X \cdot v = (\beta_1 X_{11} + \dots + \beta_d X_{dd})v$ for all diagonal matrices $X \in \mathfrak{gl}(E)$ and $X^{(c)} \cdot v = 0$ for each $c \in \{2, \dots, d\}$. \square*

Using Lemma 2.13 we establish the remaining basic properties of $\nabla^\lambda(E)$. The main novel feature is the use of Proposition 2.9 to prove (v) and (vi).

Recall that if V and W are $\mathfrak{gl}(E)$ -modules then the action of $\mathfrak{gl}(E)$ on $V \otimes W$ and $\text{Sym}^r V$ is defined by linear extension of $x \cdot (v \otimes w) = (x \cdot v) \otimes w + v \otimes (x \cdot w)$ and $x \cdot (v_{i_1} \dots v_{i_r}) = (x \cdot v_{i_1})v_{i_2} \dots v_{i_r} + \dots + v_{i_1}v_{i_2} \dots v_{i_{r-1}}(x \cdot v_{i_r})$. The λ -tableau t^λ in (iv) was defined in §2.1 by $t_{(i,j)} = i$ for each $(i, j) \in [\lambda]$.

Proposition 2.14. *Let $s \in \text{SSYT}_{\{1, \dots, d\}}(\lambda)$ and let $c \in \{2, \dots, d\}$.*

- (i) *If s has content β then $F(s)$ is a weight vector of weight β .*
- (ii) *$X^{(c)} \cdot F(s) = \sum_t F(t)$ where the sum is over all tableaux t obtained from s by changing a single entry from c to $c-1$.*
- (iii) *If every c in s has a $c-1$ immediately above it then $X^{(c)} \cdot F(s) = 0$.*
- (iv) *$F(t^\lambda)$ is a highest-weight vector of weight λ .*
- (v) *Suppose that s has a c not having a $c-1$ immediately above it. Find the highest row of s containing such an entry, and let t be the tableau obtained by changing its leftmost c to $c-1$. Then t is semistandard and $X^{(c)} \cdot F(s) = \varepsilon F(t) + y$ where $\varepsilon \in \mathbf{N}$ and y is an integral linear combination of $F(u)$ for semistandard λ -tableaux u such that $t \triangleright u$.*
- (vi) *If K has characteristic zero and $v \in \nabla^\lambda(E)$ is such that $X^{(c)} \cdot v = 0$ for all $c \in \{2, \dots, d\}$ then v is a multiple of $F(t^\lambda)$.*
- (vii) *If $K = \mathbf{C}$ then $\nabla^\lambda(E)$ is irreducible.*

Proof. Part (i) follows easily from the rules for the action of $\mathfrak{gl}(E)$ and the definition of $f(t)$ as a tensor product of symmetric powers, in analogy with Lemma 2.12(i). Since $X^{(c)} \cdot v_c = v_{c-1}$ and $X^{(c)} \cdot v_b = 0$ if $b \neq c$, we also have

$$X^{(c)} \cdot f(s) = X^{(c)} \cdot \bigotimes_{i=1}^{\ell(\lambda)} \prod_{j=1}^{\lambda_i} v_{s(i,j)} = \sum_t \bigotimes_{i=1}^{\ell(\lambda)} \prod_{j=1}^{\lambda_i} v_{t(i,j)} = \sum_t f(t).$$

where the sums are over all tableaux t obtained from s by changing a single entry from c to $c-1$. This proves the analogue of (ii) for GL-tabloids, and (ii) now follows from the definition of $F(s)$. By (2.4), $F(t) = 0$ whenever t has a repeated entry in a column, so (iii) follows from (ii). Now (iv) follows from the definition of t^λ in §2.1, (i), (iii) and Lemma 2.13.

For (v), let row a be the row of s containing the chosen entry c . By choice of a , the tableau t is semistandard. By (ii), $X^{(c)} \cdot F(s) = F(t) + \sum_{t'} F(t')$ where the sum is over all tableaux t' obtained from s by changing a different c to a $c - 1$. If this c has a $c - 1$ above it then $F(t') = 0$. If not, and this c is in row a , then by Proposition 2.9, $F(t') = F(t) + z$ where z is an integral linear combination of $F(u)$ for semistandard λ -tableaux u such that $t \triangleright u$. In the remaining case c is in a lower row than row a , and t' is column standard with $t \triangleright \bar{t}'$. Therefore, by Proposition 2.9,

$$X^{(c)} \cdot F(s) = \varepsilon F(t) + y$$

where ε is the number of entries c in row a of t not having $c - 1$ immediately above them and y is as required.

For (vi), let $v = \sum_{s \in \mathcal{S}} \gamma_s F(s)$ where $\mathcal{S} \subseteq \text{SSYT}_{\{1, \dots, d\}}(\lambda)$ and $\gamma_s \neq 0$ for each $s \in \mathcal{S}$. Suppose that v is not a multiple of $F(t^\lambda)$. Choose a minimal such that some $s \in \mathcal{S}$ has an entry c in row a with $c > a$. Choose c minimal with this property and suppose that s has ε such entries c , where ε is maximal. Thus rows $a - 1$ and a of s have the form

$$\begin{array}{cccccccc} a^- & \dots & a^- & a^- & \dots & a^- & a^- & \dots & a^- \\ a & \dots & a & \underbrace{c \quad \dots \quad c}_{\varepsilon} & \dots & & & & \end{array}$$

where a^- denotes $a - 1$. Replacing any c in row a of s with $c - 1$ gives a column-standard tableau; moreover, replacing the leftmost such c gives a semistandard λ -tableau. Let t be this tableau. Applying (v) to the summands of $X^{(c)} \cdot v$ we see that

$$X^{(c)} \cdot F(s) = \varepsilon F(t) + y$$

where y is an integral linear combination of $F(u)$ for semistandard λ -tableaux such that $t \triangleright u$. Now suppose that $\gamma_{s'} \neq 0$. Let t' be obtained from s' by changing a single c to $c - 1$; we may suppose this c has no $c - 1$ above it, and so t' is column-standard. Then, comparing $\bar{t}'^{\leq c}$ and $t^{\leq c}$ on row a , our choice of s implies that $\bar{t}' \not\triangleright t$. Therefore, by Proposition 2.9, the coefficient of $F(t)$ in $F(\bar{t}')$ is zero. Hence the coefficient of $F(t)$ in $X^{(c)} \cdot v$ is $\varepsilon \gamma_s$. Since K has characteristic zero, this is a contradiction.

Finally, by Proposition 14.13 of [13], the $\text{gl}(E)$ -submodule V of $\nabla^\lambda(E)$ generated by $F(t^\lambda)$ is irreducible. If V is a proper submodule of $\nabla^\lambda(E)$ then, by Weyl's Theorem, V has a complementary submodule. By Proposition 14.13 of [13] this complement containing a highest-weight vector, contradicting (vi). Hence $\nabla^\lambda(E)$ is irreducible. \square

We end with a result summarizing the remaining properties we need of polynomial $\text{GL}(E)$ -modules.

Proposition 2.15. *Let V be a polynomial $\text{GL}(E)$ -module of degree r .*

- (i) *V contains a highest-weight vector.*

- (ii) If $v \in V$ is a highest-weight vector of weight λ then v generates a submodule of V isomorphic to $\nabla^\lambda(E)$.
- (iii) V is isomorphic to a direct sum of certain $\nabla^\lambda(E)$ for $\lambda \in \text{Par}(r)$.
- (iv) Let Φ_V be the formal character of V . If $\lambda \in \text{Par}(r)$ has at most $\dim E$ parts then $\langle \Phi_V, s_\lambda \rangle = [V : \nabla^\lambda(E)]$.

Proof. Parts (i) and (ii) follow from Proposition 14.13 of [13] and Proposition 2.14(vii); (iii) then follows from Weyl's Theorem (see [16, §6.3] or [11, Appendix B]). By (iii) it suffices to prove (iv) when V is irreducible, in which case it is immediate from the orthogonality of Schur functions and Lemma 2.12. \square

Remark 2.16. Our definition of GL-tabloids and GL-polytabloids is in deliberate analogy with the definitions of tabloids and polytabloids in [19, (3.9), (4.3)]. In [19, Ch. 26], James uses his results on symmetric groups to construct a polynomial representation W^λ of $\text{GL}(E)$, defined over an arbitrary field K .

Let $\mathcal{G}_\otimes : S_r\text{-mod} \rightarrow \text{GL}(E)\text{-mod}_r$ be the inverse Schur functor sending a S_r -module U to $E^{\otimes r} \otimes_{KS_r} U$. (Here $E^{\otimes r}$ is regarded as a right KS_r -module by place permutation on tensors.) James' module W^λ may be defined as the image of $\mathcal{G}_\otimes(S^\lambda)$ under the canonical map $\mathcal{G}_\otimes S^\lambda \rightarrow \mathcal{G}_\otimes M^\lambda$ induced by the inclusion $S^\lambda \rightarrow M^\lambda$ of the Specht module S^λ into the Young permutation module M^λ . It follows from the definition of GL-polytabloids and the isomorphism $\mathcal{G}_\otimes(M^\lambda) \cong \bigotimes_{i=1}^{\ell(\lambda)} \text{Sym}^{\lambda_i} E$ that $W^\lambda \cong \nabla^\lambda(E)$. By working throughout in $\bigotimes_{i=1}^{\ell(\lambda)} \text{Sym}^{\lambda_i} E$ we avoid the nasty technicality that, in general, $\mathcal{G}_\otimes(S^\lambda) \not\cong W^\lambda$. For example, since $S^{(2)} \cong S^{(1,1)} \cong M^{(2)} \cong K$ when K has characteristic 2, and $\mathcal{G}_\otimes(K) = (E^{\otimes 2})_{\otimes KS_2} K \cong \text{Sym}^2 E$, we have $\mathcal{G}_\otimes(S^{(2)}) \cong \mathcal{G}_\otimes(S^{(1,1)}) \cong \text{Sym}^2 E \cong W^{(2)}$. But $W^{(2)} \not\cong W^{(1,1)} \cong \wedge^2 E$.

For more recent work on inverse Schur functions we refer the reader to [15]. A related homological remark, which explains our notation, is that $\nabla^\lambda(E)$ is a co-standard module in the sense of quasi-hereditary algebras: see [10].

3. A MODEL FOR $\nabla^\nu(\nabla^\mu(E))$

Fix $\mu \in \text{Par}(m)$ and $\nu \in \text{Par}(n)$. Let E be a complex vector space with basis e_1, \dots, e_d . Throughout this section let $\mathcal{B} = \text{SSYT}_{\{1, \dots, d\}}(\mu)$ ordered by the total order $<$ in Definition 2.1.

Lemma 3.1. *The $\text{GL}(E)$ -module $\nabla^\nu(\nabla^\mu(E))$ has as a canonical basis the set*

$$\{F(S) : S \in \text{SSYT}_{\mathcal{B}}(\nu)\}.$$

of plethystic semistandard tableaux of shape μ^ν whose μ -tableau entries each have entries from $\{1, \dots, d\}$.

Proof. By Proposition 2.11, applied with $V = E$ and $\mathcal{B} = \{1, \dots, d\}$, $\nabla^\mu(E)$ has $\{F(t) : t \in \text{SSYT}_{\{1, \dots, d\}}(\mu)\}$ as a basis. The lemma now follows from another application of Proposition 2.11, this time with $V = \nabla^\mu(E)$ and $\mathcal{B} = \text{SSYT}_{\{1, \dots, d\}}(\mu)$. \square

As a notational guide, we use upper case letters to denote ν -tableaux whose entries are μ -tableaux and upper case indices I and J to refer to their rows and columns.

By Definition 2.3,

$$(3.1) \quad F(S) = \sum_{\tau \in \text{CPP}(\nu)} f(S\tau) \text{sgn}(\sigma).$$

Since a μ -tableau entry $s = S_{(I,J)\tau^{-1}} \in \mathcal{B}$ corresponds to the basis vector $F(s)$ of $\nabla^\mu(E)$, we have

$$f(S\tau) = \bigotimes_{I=1}^{\ell(\nu)} \prod_{J=1}^{\nu_I} F(S_{(I,J)\tau^{-1}}) \text{sgn}(\tau).$$

In turn,

$$F(S_{(I,J)\tau^{-1}}) = \sum_{\sigma \in \text{CPP}(\mu)} f(s\sigma) \text{sgn}(\sigma) = \sum_{\sigma \in \text{CPP}(\mu)} \bigotimes_{i=1}^{\ell(\mu)} \prod_{j=1}^{\mu_i} u_{s_{(i,j)\sigma^{-1}}} \text{sgn}(\sigma).$$

Thus $F(S) \in \bigotimes_{I=1}^{\ell(\nu)} \text{Sym}^{\nu_I} \left(\bigotimes_{J=1}^{\ell(\mu)} \text{Sym}^{\mu_J}(E) \right)$. It will be convenient to define the *weight* of a tableau $S \in \text{SSYT}_{\mathcal{B}}(\nu)$, denoted $\text{wt}(S)$, to be the sum of the contents of its μ -tableau entries.

For example, take $\mathcal{B} = \text{SSYT}_{\{1,2,3\}}((2,1))$. If $\nu = (2,2)$ and $\mu = (2,1)$ and

$$S = \begin{array}{|c|c|} \hline \begin{array}{|c|c|} \hline 1 & 1 \\ \hline 2 & \\ \hline \end{array} & \begin{array}{|c|c|} \hline 1 & 1 \\ \hline 2 & \\ \hline \end{array} \\ \hline \begin{array}{|c|c|} \hline 1 & 1 \\ \hline 3 & \\ \hline \end{array} & \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 2 & \\ \hline \end{array} \\ \hline \end{array} \in \text{SSYT}_{\mathcal{B}}((2,2))$$

then $\text{wt}(T) = (7, 4, 1)$ and

$$\begin{aligned} F(S) = & F\left(\begin{array}{|c|c|} \hline 1 & 1 \\ \hline 2 & \\ \hline \end{array}\right) F\left(\begin{array}{|c|c|} \hline 1 & 1 \\ \hline 2 & \\ \hline \end{array}\right) \otimes F\left(\begin{array}{|c|c|} \hline 1 & 1 \\ \hline 3 & \\ \hline \end{array}\right) F\left(\begin{array}{|c|c|} \hline 1 & 2 \\ \hline 2 & \\ \hline \end{array}\right) - F\left(\begin{array}{|c|c|} \hline 1 & 1 \\ \hline 3 & \\ \hline \end{array}\right) F\left(\begin{array}{|c|c|} \hline 1 & 1 \\ \hline 2 & \\ \hline \end{array}\right) \otimes F\left(\begin{array}{|c|c|} \hline 1 & 1 \\ \hline 2 & \\ \hline \end{array}\right) F\left(\begin{array}{|c|c|} \hline 1 & 2 \\ \hline 2 & \\ \hline \end{array}\right) \\ & - F\left(\begin{array}{|c|c|} \hline 1 & 1 \\ \hline 2 & \\ \hline \end{array}\right) F\left(\begin{array}{|c|c|} \hline 1 & 2 \\ \hline 2 & \\ \hline \end{array}\right) \otimes F\left(\begin{array}{|c|c|} \hline 1 & 1 \\ \hline 3 & \\ \hline \end{array}\right) F\left(\begin{array}{|c|c|} \hline 1 & 1 \\ \hline 2 & \\ \hline \end{array}\right) + F\left(\begin{array}{|c|c|} \hline 1 & 1 \\ \hline 3 & \\ \hline \end{array}\right) F\left(\begin{array}{|c|c|} \hline 1 & 2 \\ \hline 2 & \\ \hline \end{array}\right) \otimes F\left(\begin{array}{|c|c|} \hline 1 & 1 \\ \hline 2 & \\ \hline \end{array}\right) F\left(\begin{array}{|c|c|} \hline 1 & 1 \\ \hline 2 & \\ \hline \end{array}\right). \end{aligned}$$

where $F\left(\begin{array}{|c|c|} \hline 1 & 1 \\ \hline 2 & \\ \hline \end{array}\right) = e_1^2 \otimes e_2 - e_2 e_1 \otimes e_1$, and so on.

Proposition 3.2. *Let $S \in \text{SSYT}_{\mathcal{B}}(\nu)$.*

- (i) *If $\text{wt}(S) = \beta$ then $F(S) \in \nabla^\nu(\nabla^\mu(E))$ is a weight vector of weight β .*
- (ii) *$X^{(c)} \cdot F(S) = \sum_T F(T)$ where the sum is over all ν -tableaux T obtained from S by changing a single c to $c-1$ in a single μ -tableau entry.*

Proof. Apply Proposition 2.14(i) and (ii) to (3.1). □

In particular, the canonical basis defined in Lemma 3.1 for $\nabla^\nu(\nabla^\mu(E))$ consists of weight vectors.

Proposition 3.3. *The formal character of $\nabla^\nu(\nabla^\mu(E))$ is $(s_\nu \circ s_\mu)(x_1, \dots, x_d)$.*

Proof. By the definition of Schur functions in (2.1) and the definition of plethysm given shortly afterwards, $(s_\nu \circ s_\mu)(x_1, \dots, x_d)$ is obtained by evaluating s_ν at the monomials x^t for $t \in \mathcal{B}$. Thus

$$(s_\nu \circ s_\mu)(x_1, \dots, x_d) = \sum_{S \in \text{SSYT}_{\mathcal{B}}(\nu)} x^{\text{wt}(S)}.$$

It follows that if β is a composition of mn with $\ell(\beta) = d$ then the coefficient x^β in $s_\nu \circ s_\mu$ is the number of $S \in \text{SSYT}_{\mathcal{B}}(\nu)$ of weight β . By Lemma (3.1) and Proposition 3.2(i), this is the dimension of the β -weight space in $\nabla^\nu(\nabla^\mu(E))$. \square

4. PROOF OF THEOREM 1.1

We use the model for $\nabla^\nu(\nabla^\mu(E))$ in §3, taking $\dim E = d$ where $d \geq mn$. It will be convenient to number the rows of $[(r) \sqcup \mu]$ from 0, so that

$$[(r) \sqcup \mu] = \{(0, j) : 1 \leq j \leq r\} \cup [\mu].$$

Given a μ -tableau t with entries from $\{1, \dots, d\}$, let \tilde{t} be the $((r) \sqcup \mu)$ -tableau with entries from $\{1, \dots, d, d+1\}$ defined by

$$\tilde{t}_{(i,j)} = \begin{cases} t_{(i,j)} + 1 & \text{if } i \geq 1 \\ 1 & \text{if } i = 0. \end{cases}$$

Thus \tilde{t} is obtained from t by increasing each entry by 1 and then inserting a new row of 1s of length r at the top.

The following technical lemma shows that each snake relation satisfied by $F(t)$ gives a very similar relation satisfied by $F(\tilde{t})$. We use this to show in the proof of Proposition 4.3 that $F(\tilde{t})$ can be straightened in essentially the same way as $F(t)$.

Lemma 4.1. *Let t be a μ -tableau with entries from $\{1, \dots, d\}$. Let $(i, j) \in [\mu]$ with $j < \mu_1$. Let $F(t) = -\sum_{c=2}^\ell F(t\phi_c) \text{sgn}(\phi_c)$ be a snake relation as in (2.5), with $j' = j + 1$. Then*

$$F(\tilde{t}) = -\sum_{c=2}^\ell F(\tilde{t}\phi_c) \text{sgn}(\phi_c).$$

Proof. Let $A = A_\mu(i, j)$ and $B = B_\mu(i, j + 1)$ be as in (2.5). Let $B^+ = B \cup \{(0, j + 1)\}$. (Recall that the rows of $[(r) \sqcup \mu]$ are numbered from 0.) Let $C^+ = A \cup B^+$. By hypothesis, ϕ_2, \dots, ϕ_ℓ are representatives for the proper left cosets of $S_A \times S_B$ in $S_{A \cup B}$. The permutations in $\phi_c(S_A \times S_{B^+})$ fixing $(0, j + 1)$ are precisely the elements of $\phi_c(S_A \times S_B)$. Therefore the cosets $\phi_c(S_A \times S_{B^+})$ for $2 \leq c \leq \ell$ are disjoint. Let $\phi_1 = \text{id}$ and choose further coset representatives $\phi_{\ell+1}, \dots, \phi_{\ell^+}$ such that $\phi_1, \dots, \phi_{\ell^+}$ is a full set of representatives for the left cosets of $S_A \times S_{B^+}$ in S_{C^+} . By (2.5) we have

$$(4.1) \quad F(\tilde{t}) = -\sum_{c=2}^{\ell^+} F(\tilde{t}\phi_c) \text{sgn}(\phi_c).$$

Suppose that $c > \ell$. Then $(0, j+1)\phi_c \in A$, and since $(0, j)\tilde{t} = 1$ and $(0, j+1)\tilde{t} = 1$, it follows that $\tilde{t}\phi_c$ has two entries equal to 1 in column j . Therefore $F(\tilde{t}\phi_c) = 0$ by (2.4). We may therefore replace the upper limit in the sum in (4.1) with ℓ . After making this change, (4.1) is precisely the relation we require. \square

Recall from §3 that $\mathcal{B} = \text{SSYT}_{\{1, \dots, d\}}(\mu)$ ordered by the total order $<$ in Definition 2.1. Let $\mathcal{B}^+ = \text{SSYT}_{\{1, \dots, d, d+1\}}((r) \sqcup \mu)$. Given $S \in \text{SSYT}_{\mathcal{B}}(\nu)$, let \tilde{S} be the ν -tableau defined by replacing each μ -tableau entry s of S with \tilde{s} . For $s, t \in \mathcal{B}$ we have $s < t$ if and only if $\tilde{s} < \tilde{t}$. Hence $\tilde{S} \in \text{SSYT}_{\mathcal{B}^+}(\nu)$.

Let $E^+ = E \oplus \langle e_{d+1} \rangle$ be a $(d+1)$ -dimensional complex vector space. Recall that V_λ denotes the λ -weight space of a $\mathfrak{gl}(E)$ -module V .

Lemma 4.2. *The map $F(S) \mapsto F(\tilde{S})$ defines a \mathbf{C} -linear isomorphism*

$$\nabla^\nu(\nabla^\mu(E))_\lambda \rightarrow \nabla^\nu(\nabla^{(r) \sqcup \mu}(E^+))_{(nr) \sqcup \lambda}.$$

Proof. Suppose that $v \in \nabla^\nu(\nabla^{(r) \sqcup \mu}(E^+))$ is a weight vector of weight $(nr) \sqcup \lambda$. Let $T \in \text{SSYT}_{\mathcal{B}^+}(\nu)$ and suppose that the coefficient of $F(T)$ in v is non-zero. By Proposition 3.2(i), there are nr entries equal to 1 in the μ -tableau entries of T . Since (r) is the largest part of $((r) \sqcup \mu)$, each $T_{(g,h)}$ for $(g,h) \in [\nu]$ has at most r entries equal to 1. Therefore each $T_{(g,h)}$ has exactly r entries equal to 1, necessarily lying in its longest row. Hence $T = \tilde{S}$ for a unique $S \in \text{SSYT}_{\mathcal{B}}(\nu)$, and so the map is surjective. Since it sends basis elements to basis elements, it is injective. \square

Let $\tilde{v} \in \nabla^\nu(\nabla^{(r) \sqcup \mu}(E^+))_{(nr) \sqcup \lambda}$ denote the image of $v \in \nabla^\nu(\nabla^\mu(E))_\lambda$ under the map in the previous lemma.

Proposition 4.3. *The map $v \mapsto \tilde{v}$ restricts to a bijection between highest-weight vectors in $\nabla^\nu(\nabla^\mu(E))$ of weight λ and highest-weight vectors in $\nabla^\nu(\nabla^{(r) \sqcup \mu}(E^+))$ of weight $(nr) \sqcup \lambda$.*

Proof. Let

$$v = \sum_{S \in \text{SSYT}_{\mathcal{B}}(\nu)} \alpha_S F(S) \in \nabla^\nu(\nabla^\mu(E))_\lambda.$$

By definition,

$$\tilde{v} = \sum_{S \in \text{SSYT}_{\mathcal{B}}(\nu)} \alpha_S F(\tilde{S}) \in \nabla^\nu(\nabla^{(r) \sqcup \mu}(E^+))_{\lambda+(nr)}.$$

Let $S \in \text{SSYT}_{\mathcal{B}}(\nu)$. Since changing any 2 to 1 in a $((r) \sqcup \mu)$ -tableau entry \tilde{s} of \tilde{S} gives a $((r) \sqcup \mu)$ -tableau with two 1s in the same column, Proposition 3.2(ii) implies that $X^{(2)} \cdot F(\tilde{S}) = 0$. Now let $c \in \{2, \dots, d\}$. Again by Proposition 3.2(ii), $X^{(c)} \cdot F(S) = \sum_T F(T)$ where the sum is over all $T \in \text{SSYT}_{\mathcal{B}}(\nu)$ obtained from S by changing a single c to $c-1$. Moreover, $X^{(c+1)} \cdot F(\tilde{S}) = \sum_T F(\tilde{T})$ with the same conditions on the sum.

Suppose that v is a highest-weight vector. By the previous paragraph and (3.1), each summand $F(T)$ appearing in $X^{(c)} \cdot F(S)$ (respectively, each $F(\tilde{T})$ appearing in

$X^{(c+1)} \cdot F(\tilde{S})$) is a sum of tensor products of symmetric products of $F(u)$ (respectively $F(\tilde{u})$) for certain μ -tableaux u (respectively $((r) \sqcup \mu)$ -tableaux \tilde{u}), at most one of which, say t (respectively \tilde{t}), is non-semistandard. By Corollary 2.6, we may straighten $F(t)$ to a linear combination of $F(s)$ for $s \in \text{SSYT}_{\{1, \dots, d\}}(\mu)$ by a sequence of snake relations (2.5) swapping boxes between adjacent columns. By multilinearity, this expresses $F(T)$ as a linear combination of $F(S)$ for $S \in \text{SSYT}_{\mathcal{B}}(\nu)$. Recall that the rows of $[(r) \sqcup \mu]$ are labelled from 0. By Lemma 4.1 if we apply *exactly the same sequence of relations* to straighten $F(\tilde{t})$, we express $F(\tilde{T})$ as a linear combination of $F(\tilde{S})$ for $S \in \text{SSYT}_{\mathcal{B}}(\nu)$ *with the same coefficients*. Hence $X^{(c)} \cdot v = 0$ implies $X^{(c+1)} \cdot \tilde{v} = 0$. By Lemma 2.13, \tilde{v} is a highest-weight vector.

Conversely, if \tilde{v} is a highest-weight vector then running this argument in reverse shows that $X^{(c)} \cdot v = 0$ for $c \in \{2, \dots, d\}$, and so v is a highest-weight vector. \square

By Proposition 4.3 and Proposition 2.15 we have $[\nabla^\nu(\nabla^\mu E) : \nabla^\lambda E] = [\nabla^\nu(\nabla^{(r) \sqcup \mu})(E) : \nabla^{(nr) \sqcup \lambda}(E)]$. Theorem 1.1 now follows using (1.7).

Remark 4.4. We remark that since any plethystic semistandard tableau of shape μ^ν has at most $n\mu_1$ integer entries of 1, if $\lambda_1 > n\mu_1$ then both sides of the equation in Theorem 1.1 are zero.

Remark 4.5. In [20, Lemma 3.2] a proof of the special case $\mu = (1^m)$ and $r = 1$ is indicated. In our notation, the authors consider $\nabla^\nu(\bigwedge^{m+1}(E))$ as a submodule of $(\bigwedge^{m+1}(E))^{\otimes n}$ and observe that each tensor summand in a highest-weight vector v of weight $(n) \sqcup \lambda$ is of the form $(e_1 \wedge \dots) \otimes \dots \otimes (e_1 \wedge \dots)$. They define a map into $(\bigwedge^m(E))^{\otimes n}$ by removing e_1 from each tensor factor of v and reducing the indices. This is essentially the inverse map to ours, in this special case.

5. PROOF OF THEOREM 1.2

We adapt the strategy used to prove Theorem 1.1, again working in the model $\nabla^\nu(\nabla^\mu(E))$ from §3, now taking $\dim E = d$ where $d \geq r$ and $d \geq \ell(\mu)$.

If $r \geq \ell(\mu)$ then set $e = 1$. Otherwise let $e = \mu_{r+1} + 1$. To relate $[\mu]$ and $[\mu + (1^r)]$ we use the following notation. Let $[i, 0] = (i, e)$ for $1 \leq i \leq r$ and for $(i, j) \in [\mu]$, let

$$[i, j] = \begin{cases} (i, j) & \text{if } j < e \\ (i, j + 1) & \text{if } j \geq e. \end{cases}$$

As illustrated in Example 5.1, we have

$$[\mu + (1^r)] = \{[i, 0] : 1 \leq i \leq r\} \cup \{[i, j] : (i, j) \in [\mu]\}.$$

Given a μ -tableau t with entries from $\{1, \dots, d\}$, let t^* be the $(\mu + (1^r))$ -tableau defined by $t^*_{[i, 0]} = i$ for $1 \leq i \leq r$ and $t^*_{[i, j]} = t_{(i, j)}$ if $j > 0$. Thus t^* is obtained from t by inserting a new column e with entries $1, \dots, r$, moving the existing column e and other later numbered columns one position right.

Example 5.1. If $r = 2$ and $\mu = (4, 2, 1)$ then $e = 2$; the labels for the boxes in $[(4, 2, 1)]$ and $[(4, 2, 1) + (1, 1)]$ are

(1,1)	(1,2)	(1,3)	(1,4)	,	[1,1]	[1,0]	[1,2]	[1,3]	[1,4]	
(2,1)	(2,2)					[2,1]	[2,0]	[2,2]		
(3,1)						[3,1]				

Two pairs of a μ -tableau t and the corresponding $(\mu + (1^r))$ -tableau t^* are

$$\left(\begin{array}{|c|c|c|c|} \hline 1 & 1 & 2 & 3 \\ \hline 2 & 3 & & \\ \hline 4 & & & \\ \hline \end{array}, \begin{array}{|c|c|c|c|} \hline 1 & 1 & 1 & 2 & 3 \\ \hline 2 & 2 & 3 & & \\ \hline 4 & & & & \\ \hline \end{array} \right), \quad \left(\begin{array}{|c|c|c|c|} \hline 1 & 1 & 2 & 3 \\ \hline 3 & 3 & & \\ \hline 4 & & & \\ \hline \end{array}, \begin{array}{|c|c|c|c|} \hline 1 & 1 & 1 & 2 & 3 \\ \hline 3 & 2 & 3 & & \\ \hline 4 & & & & \\ \hline \end{array} \right).$$

Given $\phi \in S_{[\mu]}$, let $\phi^* \in S_{[\mu] + (1^r)}$ be defined by $[i, 0]\phi^* = [i, 0]$ for $1 \leq i \leq r$ and $[i, j]\phi^* = [i', j'] \iff (i, j)\phi = (i', j')$. In analogy with Lemma 4.1, we now show that $F(t^*)$ satisfies the appropriate conjugate of each snake relation (see (2.5) after Lemma 2.4) satisfied by $F(t)$.

Lemma 5.2. *Let t be a μ -tableau with entries from $\{1, \dots, d\}$. Let $(i, j) \in [\mu]$ with $j < \mu_1$. Let $F(t) = -\sum_{c=2}^{\ell} F(t\phi_c) \operatorname{sgn}(\phi_c)$ be a snake relation as in (2.5), with $j' = j + 1$. Then*

$$F(t^*) = -\sum_{c=2}^{\ell} F(t^*\phi_c^*) \operatorname{sgn}(\phi_c^*).$$

Proof. The claimed relation is an instance of (2.5) for t^* , with respect to the boxes $[i, j]$ and $[i, j + 1] \in [\mu] + (1^r)$. \square

If $e = 1$ the added column in t^* is at the far left, and Lemma 5.2 may be compared with Lemma 4.1. If $e > 1$ and $j = e - 1$ then $[i, j]$ and $[i, j + 1]$ lie in the non-adjacent columns $e - 1$ and $e + 1$ of $[\mu] + (1^r)$; this is the only case where we need the freedom in (2.5) to take $j' \neq j + 1$.

Recall from §3 that $\mathcal{B} = \operatorname{SSYT}_{\{1, \dots, d\}}(\mu)$. Let \mathcal{C}^* be the set of column-standard $(\mu + (1^r))$ -tableaux with entries from $\{1, \dots, d\}$ and let $\mathcal{B}^+ = \operatorname{SSYT}_{\{1, \dots, d\}}(\mu + (1^r))$. Thus $\mathcal{B}^+ \subseteq \mathcal{C}^*$. Both \mathcal{B} and \mathcal{C}^* are ordered by the total order $<$ in Definition 2.1. Given $T \in \operatorname{SSYT}_{\mathcal{B}}(\nu)$, let T^* be the ν -tableau defined by replacing each μ -tableau entry s of T with s^* . For $s, t \in \mathcal{B}$ we have $s < t$ if and only if $s^* < t^*$, since the inserted column is the same in s^* and t^* . Hence $T^* \in \operatorname{SSYT}_{\mathcal{C}^*}(\nu)$ and $F(T^*) \in \nabla^{\nu}(\nabla^{\mu + (1^r)}(E))$. By the case $N = 1$ of the following definition, $T^* \in \operatorname{SSYT}_{\mathcal{B}^+}(\nu)$ if and only if T^* is r -saturated.

Definition 5.3. Recall that if $r \geq \ell(\mu)$ then $e = 1$ and otherwise $e = \mu_{r+1} + 1$. Let $N \in \mathbf{N}_0$ and let U be a ν -tableau whose entries are certain $\mu + N(1^r)$ -tableaux. We say that U is r -saturated if whenever u is a $\mu + N(1^r)$ -tableau entry of U , we have $u_{(i,j)} = i$ for $1 \leq i \leq r$ and $1 \leq j \leq e$.

Equivalently, U is r -saturated if the first e columns of each $\mu + N(1^r)$ -tableau entry of U each begin $1, \dots, r$ when read from top to bottom. For example, when $r = 2$ and $\mu = (4, 2, 1)$ we saw in Example 5.1 that $e = 2$. Taking $N = 1$, of the two

$((4, 2, 1) + (1^2))$ -tableaux t^* shown, only the first could be an entry of a 2-saturated tableau, since for the second $(2, 1)t^* = 3$.

Given $v \in \nabla^\nu(\nabla^\mu(E))$ let $v^* \in \nabla^\nu(\nabla^{\mu+(1^r)})$ denote the image of v under the \mathbf{C} -linear map defined on the canonical basis in Lemma 3.1 of $\nabla^\nu(\nabla^\mu(E))$ by $F(T) \mapsto F(T^*)$ for each $T \in \text{SSYT}_{\mathcal{B}}(\nu)$.

Example 5.4. Let $r = 2$ and let $\mu = (2, 1, 1)$ so $e = 2$. Let $d = 4$, so $\mathcal{B} = \text{SSYT}_{\{1,2,3,4\}}((2, 1, 1))$. Take $\nu = (2)$. A calculation, either in MAGMA or by hand using the domino tableau rule in [6, Theorem 4.1], shows that $\langle s_{(2)} \circ s_{(2,1,1)}, s_{(3,2,2,1)} \rangle = 1$ and so the space of highest-weight vectors of weight $(3, 2, 2, 1)$ in $\text{Sym}^2(\nabla^{(2,1,1)}(E))$ is 1-dimensional. Computing the images of the $F(s)$ for $s \in \mathcal{B}$ using Proposition 2.14(ii) one finds that if $T_{(1)}, T_{(2)}, T_{(3)}, T_{(4)}$ are the four tableaux in $\text{SSYT}_{\mathcal{B}}((2))$ shown below

$$\begin{array}{|c|c|} \hline 1 & 1 \\ \hline 2 & \\ \hline 3 & \\ \hline \end{array}, \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & \\ \hline 4 & \\ \hline \end{array}, \begin{array}{|c|c|} \hline 1 & 1 \\ \hline 2 & \\ \hline 3 & \\ \hline \end{array}, \begin{array}{|c|c|} \hline 1 & 3 \\ \hline 2 & \\ \hline 4 & \\ \hline \end{array}, \begin{array}{|c|c|} \hline 1 & 1 \\ \hline 3 & \\ \hline 4 & \\ \hline \end{array}, \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 2 & \\ \hline 3 & \\ \hline \end{array}, \begin{array}{|c|c|} \hline 1 & 1 \\ \hline 2 & \\ \hline 4 & \\ \hline \end{array}, \begin{array}{|c|c|} \hline 1 & 3 \\ \hline 2 & \\ \hline 3 & \\ \hline \end{array}$$

then, by Lemma 2.13,

$$v = F(T_{(1)}) - F(T_{(2)}) - F(T_{(3)}) + F(T_{(4)}) \in \text{Sym}^2(\nabla^{(2,1,1)}(E)).$$

is a highest-weight vector of weight $(3, 2, 2, 1)$. For example, by Proposition 2.14(ii), we have $X^{(2)} \cdot F(T_{(2)}) = 0$, $X^{(2)} \cdot F(T_{(4)}) = 0$ and

$$X^{(2)} \cdot F(T_{(1)}) = F\left(\begin{array}{|c|c|c|} \hline 1 & 1 & 1 \\ \hline 2 & & \\ \hline 3 & & \\ \hline \end{array}\right) F\left(\begin{array}{|c|c|} \hline 1 & 1 \\ \hline 3 & \\ \hline 4 & \\ \hline \end{array}\right) = X^{(2)} \cdot F(T_{(3)});$$

since $T_{(1)}$ and $T_{(3)}$ appear with opposite signs in v , this implies $X^{(2)} \cdot v = 0$. Essentially the same calculation shows that

$$v^* = F(T_{(1)}^*) - F(T_{(2)}^*) - F(T_{(3)}^*) + F(T_{(4)}^*) \in \text{Sym}^2(\nabla^{(2,1,1)+(1,1)}(E))$$

is a highest-weight vector of weight $(3, 2, 2, 1) + 2(1, 1)$. The tableaux $T_{(1)}^*$ and $T_{(3)}^*$ are not 2-saturated; they lie in $\text{SSYT}_{\mathcal{C}^*}((2))$ but not in $\text{SSYT}_{\mathcal{B}^+}((2))$. For example

$$\begin{aligned} F(T_{(1)}^*) &= F\left(\begin{array}{|c|c|c|} \hline 1 & 1 & 1 \\ \hline 2 & 2 & \\ \hline 3 & & \\ \hline \end{array}\right) F\left(\begin{array}{|c|c|c|} \hline 1 & 1 & 2 \\ \hline 3 & 2 & \\ \hline 4 & & \\ \hline \end{array}\right) \\ &= F\left(\begin{array}{|c|c|c|} \hline 1 & 1 & 1 \\ \hline 2 & 2 & \\ \hline 3 & & \\ \hline \end{array}\right) F\left(\begin{array}{|c|c|c|} \hline 1 & 1 & 2 \\ \hline 2 & 3 & \\ \hline 4 & & \\ \hline \end{array}\right) - F\left(\begin{array}{|c|c|c|} \hline 1 & 1 & 1 \\ \hline 2 & 2 & \\ \hline 3 & & \\ \hline \end{array}\right) F\left(\begin{array}{|c|c|c|} \hline 1 & 1 & 2 \\ \hline 2 & 4 & \\ \hline 3 & & \\ \hline \end{array}\right) \end{aligned}$$

expressed in the canonical basis of $\text{Sym}^2(\nabla^{(3,2,1)}(E))$. We leave it as an exercise to show that if $U_{(1)}, U_{(2)}, U_{(3)}, U_{(4)}$ are the four tableaux in $\text{SSYT}_{\mathcal{C}^*}((2))$ shown

below

$$\begin{array}{|c|c|c|} \hline 1 & 1 & 1 \\ \hline 2 & 3 & \\ \hline 3 & & \\ \hline \end{array} \quad \begin{array}{|c|c|c|} \hline 1 & 1 & 2 \\ \hline 2 & 2 & \\ \hline 4 & & \\ \hline \end{array}, \quad \begin{array}{|c|c|c|} \hline 1 & 1 & 1 \\ \hline 2 & 2 & \\ \hline 4 & & \\ \hline \end{array} \quad \begin{array}{|c|c|c|} \hline 1 & 1 & 2 \\ \hline 2 & 3 & \\ \hline 3 & & \\ \hline \end{array}, \quad \begin{array}{|c|c|c|} \hline 1 & 1 & 1 \\ \hline 2 & 4 & \\ \hline 3 & & \\ \hline \end{array} \quad \begin{array}{|c|c|c|} \hline 1 & 1 & 2 \\ \hline 2 & 2 & \\ \hline 3 & & \\ \hline \end{array}, \quad \begin{array}{|c|c|c|} \hline 1 & 1 & 1 \\ \hline 2 & 2 & \\ \hline 3 & & \\ \hline \end{array} \quad \begin{array}{|c|c|c|} \hline 1 & 1 & 2 \\ \hline 2 & 4 & \\ \hline 3 & & \\ \hline \end{array}$$

then

$$w = F(U_{(1)}) - F(U_{(2)}) - F(U_{(3)}) + F(U_{(4)}) - F(T_{(2)}^*) + F(T_{(4)}^*)$$

is a highest-weight vector in $\text{Sym}^2(\nabla^{(2,1,1)+(1,1)}(E))$ of weight $(3, 2, 2, 1) + 2(1, 1)$, linearly independent of v^* . By Theorem 1.2, the multiplicity

$$[\text{Sym}^2(\nabla^{(2,1,1)+N(1,1)}(E)) : \nabla^{(3,2,2,1)+N(2,2)}]$$

is constant for $N \geq 1$. A further domino tableau calculation shows that $\langle s_{(2)} \circ s_{(3,2,1)}, s_{(5,4,2,1)} \rangle = 2$. Therefore v^{**} and w^* span the subspace of $\text{Sym}^2(\nabla^{(2,1,1)+2(1,1)}(E))$ of highest-weight vectors of weight $(3, 2, 2, 1) + 2(2, 2)$, and a similar result holds for all further additions.

Lemma 5.5. *The map $v \mapsto v^*$ defines a \mathbf{C} -linear injection*

$$\nabla^\nu(\nabla^\mu(E))_\lambda \rightarrow \nabla^\nu(\nabla^{\mu+(1^r)}(E))_{\lambda+n(1^r)}.$$

Proof. Let $T \in \text{SSYT}_{\mathcal{B}}(\nu)$. Since $\text{wt}(T^*) = \text{wt}(T) + n(1^r)$, Proposition 3.2(i) implies that $F(T^*) \in \nabla^\nu(\nabla^{\mu+(1^r)}(E))_{\lambda+n(1^r)}$. By definition

$$(5.1) \quad F(T^*) = \sum_{\tau \in \text{CPP}(\nu)} \bigotimes_{I=1}^{\ell(\nu)} \prod_{J=1}^{\nu_I} F(T_{(I,J)\tau^{-1}}^*) \text{sgn}(\tau) \in \text{SSYT}_{\mathcal{C}^*}(\nu).$$

The row-standardization \bar{t} of a tableau t was defined in Definition 2.8. Applying this operation to each entry of T^* , we define $S(T^*) \in \text{SSYT}_{\mathcal{B}^+}(\nu)$ by $S(T^*)_{(I,J)} = \overline{T_{(I,J)}^*}$. By Proposition 2.9,

$$F(T_{(I,J)\tau^{-1}}^*) = F(S(T^*)_{(I,J)\tau^{-1}}) + u_{(I,J)\tau^{-1}}$$

where $u_{(I,J)\tau^{-1}} \in \nabla^{\mu+(1^r)}(E)$ is a linear combination of GL-polytabloids $F(s)$ for $s \in \mathcal{B}^+$ such that $\overline{T_{(I,J)\tau^{-1}}^*} \triangleright s$. Define $V \in \nabla^\nu(\nabla^{\mu+(1^r)}(E))$ by

$$F(T^*) = F(S(T^*)) + V.$$

By the previous sentence, V is a linear combination of basis elements of $\text{Sym}^\nu(\nabla^{\mu+(1^r)}(E))$ each of the form

$$\bigotimes_{I=1}^{\ell(\nu)} \prod_{J=1}^{\nu_I} F(u_{(I,J)}),$$

where the tableaux $u_{(I,J)} \in \mathcal{B}^+$ can be relabelled by a permutation τ so that $S(T^*)_{(I,J)} \triangleright u_{(I,J)\tau}$ for each $(I, J) \in [\nu]$, with at least one of these dominance relations strict. It follows that the coefficient of $f(S(T^*))$ in $F(T^*)$ comes entirely from $F(S(T^*))$. By Lemma 2.10, this coefficient is 1.

Each $\mu + (1^r)$ -tableau entry of $S(T^*)$ is of the form $\overline{t^*}$ where $t \in \mathcal{B}$. Given $s = \overline{t^*}$ one may reconstruct t as follows: choose, for each $i \in \{1, \dots, r\}$, a box (i, j_i)

containing i in row i of s ; now erase the entry in this box, and move each entry to the right of the now empty box one place to the left; finally delete the box at the end of row i . More formally,

$$t_{(i,h)} = \begin{cases} s_{(i,h+1)} & \text{if } i \in \{1, \dots, r\} \text{ and } h \geq j_i \\ s_{(i,h)} & \text{otherwise.} \end{cases}$$

Therefore the map $T \mapsto S(T^*)$ is injective.

Let

$$v = \sum_{T \in \text{SSYT}_{\mathcal{B}}(\nu)} \alpha_T F(T^*)$$

where not every coefficient is zero. Choose T so that $S(T^*)$ is a maximal element of $\{S(T^*) : \alpha_T \neq 0\}$ in the dominance order. By the previous two paragraphs, the coefficient of $f(S(T^*))$ in v is α_T . Hence the map $F(T) \mapsto F(T^*)$ is injective. \square

Proposition 5.6. *The map $v \mapsto v^*$ restricts to an injective \mathbf{C} -linear map from highest-weight vectors in $\nabla^\nu(\nabla^\mu(E))$ of weight λ to highest-weight vectors in $\nabla^\nu(\nabla^{\mu+(1^r)}(E))$ of weight $\lambda + (n^r)$. Moreover if every $U \in \text{SSYT}_{\mathcal{B}^+}(\nu)$ of weight $\lambda + (n^r)$ is r -saturated then the map is bijective.*

Proof. The first part follows by combining Proposition 3.2(ii), Lemma 5.2 and Lemma 5.5, in the same way as Proposition 4.3. If the final hypothesis holds then every $U \in \text{SSYT}_{\mathcal{B}^+}(\nu)$ is of the form T^* for some $T \in \text{SSYT}_{\mathcal{B}}(\nu)$ and so, by Lemma 3.1 and Proposition 3.2(i), the map $F(T) \mapsto F(T^*)$ defines a linear isomorphism $\nabla^\nu(\nabla^\mu(E))_\lambda \rightarrow \nabla^\nu(\nabla^{\mu+(1^r)}(E))_{\lambda+(n^r)}$. Therefore in this case the restricted map is bijective on highest-weight vectors. \square

We need the following sufficient condition for the r -saturation condition.

Lemma 5.7. *Let $\mathcal{B}^{+M} = \text{SSYT}_{\{1, \dots, d\}}(\mu + M(1^r))$. Every element of $\text{SSYT}_{\mathcal{B}^{+M}}(\nu)$ of weight λ is r -saturated if*

$$M > n(\mu_1 + \dots + \mu_{r-1}) + (n-1)\mu_r + \mu_{r+1} - (\lambda_1 + \dots + \lambda_r).$$

Proof. Let u be a $\mu + M(1^r)$ -tableau entry of $U \in \text{SSYT}_{\mathcal{B}^{+M}}(\nu)$. The entries of u in $\{1, \dots, r\}$ lie in its first r rows. Therefore u has at most $\mu_1 + \dots + \mu_r + Mr$ such entries. If U is not saturated then it has a $\mu + M(1^r)$ -tableau entry t such that $t_{(r,e)} > r$. This t has at most $\mu_1 + \dots + \mu_{r-1} + M(r-1) + (e-1)$ entries in $\{1, \dots, r\}$. Since $e = \mu_{r+1} + 1$, this shows that U has at most $(n-1)(\mu_1 + \dots + \mu_r + Mr) + (\mu_1 + \dots + \mu_{r-1} + M(r-1) + \mu_{r+1})$ entries in $\{1, \dots, r\}$. The number of such entries is $\lambda_1 + \dots + \lambda_r + Mnr$. Therefore

$$n(\mu_1 + \dots + \mu_{r-1} + Mr) + (n-1)\mu_r + \mu_{r+1} - M \geq \lambda_1 + \dots + \lambda_r + Mnr.$$

Rearranging, this implies the lemma. \square

Proof of Theorem 1.2. By Proposition 5.6 and Proposition 2.15 we have

$$[\nabla^\nu(\nabla^\mu(E)) : \nabla^\lambda(E)] \leq [\nabla^\nu(\nabla^{\mu+(1^r)}(E)) : \nabla^{\lambda+n(1^r)}(E)].$$

The first part of Theorem 1.2 now follows from (1.7). Now suppose that $N \geq n(\mu_1 + \cdots + \mu_{r-1}) + (n-1)\mu_r + \mu_{r+1} - (\lambda_1 + \cdots + \lambda_r)$. By Lemma 5.7, taking $M = N + 1$, every element of $\text{SSYT}_{\mathcal{B}+(N+1)}(\nu)$ is r -saturated. Therefore, by Proposition 5.6, the map $v \mapsto v^*$ from highest-weight vectors in $\nabla^\nu(\nabla^{\mu+N(1^r)}(E))$ to highest-weight vectors in $\nabla^\nu(\nabla^{\mu+(N+1)(1^r)}(E))$ is a bijection. The stability result now follows from Proposition 2.15. \square

Example 5.8. Example 5.4 shows that the stability bound in Theorem 1.2 may be sharp. We give an example of the opposite case. Fix $n \in \mathbf{N}$. It is known (see for example [25, §8.5]) that $\bigwedge^n(\text{Sym}^2 E)$ is multiplicity-free. Moreover, the partitions λ such that $[\bigwedge^n(\text{Sym}^2 E) : \nabla^\lambda(E)] = 1$ are all incomparable under the dominance order, and correspond, by Theorem 1.5, to the maximal weights of the plethystic semistandard tableaux of shape $(2)^{(1^n)}$. For example, $\bigwedge^3(\text{Sym}^2 E) = \nabla^{(4,1,1)}(E) \oplus \nabla^{(3,3)}(E)$, corresponding to the plethystic tableau whose single column has (2)-tableau entries

$$\left\{ \boxed{1 \ 1}, \boxed{1 \ 2}, \boxed{1 \ 3} \right\}, \left\{ \boxed{1 \ 1}, \boxed{1 \ 2}, \boxed{2 \ 2} \right\},$$

respectively. More generally, for each $\ell \in \mathbf{N}$, provided that $\dim E \geq \ell$, $\bigwedge^{\binom{\ell+1}{2}}(\text{Sym}^2 E)$ has $\nabla^{((\ell+1)^\ell)}(E)$ as an irreducible constituent, corresponding to the plethystic semistandard tableau of shape $(2)^{(1^n)}$ where $n = \binom{\ell+1}{2}$, defined using all 2-multisubsets of $\{1, \dots, \ell\}$.

Let λ be a partition of $2n$ such that $[\bigwedge^n(\text{Sym}^2 E) : \nabla^\lambda(E)] = 1$. Let $\{\boxed{a_1 \ b_1}, \dots, \boxed{a_n \ b_n}\}$ be the entries in the corresponding plethystic semistandard tableau of shape $(2)^{(1^n)}$. Let $N \in \mathbf{N}$. Then the unique plethystic semistandard tableau of shape $(2+N)^{(1^n)}$ and weight $\lambda + (nN)$ has $(2+N)$ -tableau entries $\{u_1, \dots, u_n\}$ where for each i ,

$$u_i = \boxed{1 \ \cdots \ 1 \ a_i \ b_i}.$$

Hence, by Theorem 1.5, $[\bigwedge^n(\text{Sym}^{2+N} E) : \nabla^{\lambda+(nN)}(E)] = 1$ for all $N \in \mathbf{N}_0$. This stability follows from Theorem 1.2 for $N \geq 2(n-1) - \lambda_1$. In the case $[\bigwedge^{\binom{\ell+1}{2}} \text{Sym}^{2+N}(E) : \nabla^{((\ell+1)^\ell)+(nN)}(E)] = 1$ this bound becomes $N \geq 2\binom{\ell+1}{2} - 1 - (\ell+1) = \ell^2 - 3$; clearly this can be arbitrarily large.

We end with a combinatorial upper bound for the stable multiplicity. Example 5.8 shows that the bound is sharp in infinitely many cases.

Proposition 5.9. *Let L be the greater of $n(\mu_1 + \cdots + \mu_{r-1}) + (n-1)\mu_r + \mu_{r+1} - (\lambda_1 + \cdots + \lambda_r)$ and 0. Then*

$$|\nabla^\nu(\nabla^{\mu+N(1^r)}(E)) : \nabla^{\lambda+N(n^r)}(E)| \leq |\{T \in \text{SSYT}_{\mathcal{B}+L}(\nu) : \text{wt}(T) = \lambda + L(n^r)\}|$$

for all $N \geq L$.

Proof. The bound holds when $N = L$ since the right-hand side is

$$\dim \nabla^\nu(\nabla^{\mu+L(1^r)}(E))_{\lambda+L(n^r)}$$

and by Proposition 2.15 this is an upper bound for the left-hand side. By Theorem 1.2 the bound holds for all $N \geq L$. \square

6. PROOF OF THEOREM 1.3

It is equivalent to show that if $n^* \in \mathbf{N}_0$, $\lambda^* \in \text{Par}(mn^*)$ and $\text{Sym}^{n^*}(\nabla^\mu(E))$ has $\nabla^{\lambda^*}(E)$ as an irreducible constituent then

$$[\text{Sym}^{n+n^*}(\nabla^\mu(E)) : \nabla^{\lambda+\lambda^*}(E)] \geq [\text{Sym}^n(\nabla^\mu(E)) : \nabla^\lambda(E)].$$

Let $c = [\text{Sym}^n(\nabla^\mu(E)) : \nabla^\lambda(E)]$. Choose linearly independent highest-weight vectors $v_1, \dots, v_c \in \text{Sym}^n(\nabla^\mu(E))$ each of weight λ . By the hypothesis and Proposition 2.15, there is a highest-weight vector $w \in \text{Sym}^{n^*}(\nabla^\mu(E))$ of weight λ^* . Multiplying highest-weight vectors in the polynomial algebra $\bigoplus_{r=0}^\infty \text{Sym}^r(\nabla^\lambda(E))$, we see that $v_1 w, \dots, v_c w$ are c linearly independent highest-weight vectors in $\text{Sym}^{n+n^*}(\nabla^\mu(E))$ each of weight $\lambda + \lambda^*$. The result follows.

7. PROOF OF THEOREM 1.5

Let $d = \ell(\lambda)$ and recall that $E = \langle e_1, \dots, e_d \rangle$ is a d -dimensional complex vector space. Let $\mathcal{B} = \text{SSYT}_{\{1, \dots, d\}}(\mu)$. By Definition 1.4 the plethystic semistandard tableaux of shape μ^ν whose μ -tableau entries have entries from $\{1, \dots, d\}$ are precisely the elements of $\text{SSYT}_{\mathcal{B}}(\nu)$. By Lemma 3.1, $\nabla^\nu(\nabla^\mu(E))$ has $\{F(S) : S \in \text{SSYT}_{\mathcal{B}}(\nu)\}$ as a canonical basis. By Proposition 3.2, if $S \in \text{SSYT}_{\mathcal{B}}(\nu)$ has weight λ then $F(S) \in \nabla^\nu(\nabla^\mu(E))$ is a weight vector of weight λ .

We use this canonical basis to prove the following two results; the second is illustrated in Example 7.4 below.

Lemma 7.1. *If $[\nabla^\nu(\nabla^\mu(E)) : \nabla^\lambda(E)] \geq 1$ then there exists a plethystic semistandard tableau $T \in \text{SSYT}_{\mathcal{B}}(\nu)$ such that $\text{wt}(T) = \lambda$.*

Proof. Let $v \in \nabla^\nu(\nabla^\mu(E))$ be a highest-weight vector of weight λ . Let $v = \sum_{S \in \text{SSYT}_{\mathcal{B}}(\nu)} c_S F(S)$ be the expression of v in the canonical basis given by Lemma 3.1 of $\nabla^\nu(\nabla^\mu(E))$. By Proposition 3.2(i), each S such that $c_S \neq 0$ has weight λ . Take T to be any such S . \square

Proposition 7.2. *Suppose that λ is maximal in the dominance order on partitions such that there exists a plethystic semistandard tableau $T \in \text{SSYT}_{\mathcal{B}}(\nu)$ of weight λ . Then $F(T) \in \nabla^\nu(\nabla^\mu(E))$ is a highest-weight vector of weight λ .*

Proof. By Proposition 3.2(i), $F(T)$ is a weight vector of weight λ . Suppose, for a contradiction, that $F(T)$ is not highest-weight. Then there exists $c \in \{2, \dots, d\}$ such that $X^{(c)} \cdot F(T) \neq 0$. By Proposition 3.2(ii), $X^{(c)} \cdot F(T) = \sum F(U)$, where each U is obtained from T by changing a single c to $c - 1$ in a μ -tableau entry of

T . Thus each U has weight λ^* where

$$\lambda_b^* = \begin{cases} \lambda_b + 1 & \text{if } b = c - 1 \\ \lambda_b - 1 & \text{if } b = c \\ \lambda_b & \text{otherwise.} \end{cases}$$

Let

$$X^{(c)} \cdot F(T) = \sum_{S \in \text{SSYT}_{\mathcal{B}}(\nu)} c_S F(S)$$

be the expression of $\sum F(U)$ in the canonical basis of $\nabla^\nu(\nabla^\mu(E))$. Choose S such that $c_S \neq 0$. Then $\text{wt}(S) = \lambda^* \triangleright \lambda = \text{wt}(T)$. This contradicts the maximality of λ . \square

We are now ready to prove Theorem 1.5.

Proof of Theorem 1.5. By Lemma 7.1, if $[\nabla^\mu(\nabla^\mu(E)) : \nabla^\lambda(E)] \geq 1$ then there is a plethystic semistandard tableau $T \in \text{SSYT}_{\mathcal{B}}(\nu)$ of weight λ . Conversely, by Proposition 7.2, if λ is maximal in the dominance order such that there is a plethystic semistandard tableau $T \in \text{SSYT}_{\mathcal{B}}(\nu)$ then $F(T)$ is a highest-weight vector, and so $[\nabla^\mu(\nabla^\mu(E)) : \nabla^\lambda(E)] \geq 1$. Therefore the maximal partitions λ in the dominance order such that $[\nabla^\nu(\nabla^\mu(E)) : \nabla^\lambda(E)] \geq 1$ are precisely the maximal weights of the elements of $\text{SSYT}_{\mathcal{B}}(\nu)$. This proves the first part of the theorem. Now suppose that λ is maximal in the dominance order such that $\nabla^\lambda(E)$ appears in $\nabla^\nu(\nabla^\mu(E))$. Let $S_{(1)}, \dots, S_{(r)}$ be the plethystic semistandard tableaux of shape μ^ν and weight λ . By the canonical basis in Lemma 3.1 and Proposition 3.2(i), $F(S_{(1)}), \dots, F(S_{(r)})$ form a basis for the weight space $\nabla^\nu(\nabla^\mu(E))_\lambda$. By Proposition 7.2 these vectors are highest-weight. Therefore

$$r = [\nabla^\nu(\nabla^\mu(E)) : \nabla^\lambda(E)]$$

as required. \square

As a corollary we obtain the result mentioned in the introduction that is surprisingly non-trivial to prove entirely combinatorially.

Corollary 7.3. *If T is a plethystic semistandard tableau of maximal weight then $\text{wt}(T)$ is a partition.*

Proof. Since the weight of a highest-weight vector is a partition, this is immediate from Proposition 7.2. \square

By Proposition 1.9, if T is a plethystic semistandard tableau of shape $\mu^{(1^n)}$ and maximal weight then the set of μ -tableau entries of T is closed, in the sense of Definition 1.8. The converse does not hold: we show this in the following example, which makes constructive the proof of Proposition 7.2.

Example 7.4. For ease of notation we shall identify $\nabla^{(1^n)}(\nabla^\mu(E))$ with $\wedge^n(\nabla^\mu(E))$ via the map sending $F(T)$, where T is a plethystic semistandard tableau with μ -tableau entries t_1, \dots, t_n read from top to bottom, to $F(t_1) \wedge \dots \wedge F(t_n)$.

Let T be the plethystic semistandard tableau of shape $(2, 2)^{(1^{11})}$ and weight $(17, 11, 8, 8)$ whose $(2, 2)$ -tableau entries, read from top to bottom are

$$\begin{array}{|c|c|} \hline 1 & 1 \\ \hline 2 & 2 \\ \hline \end{array}, \begin{array}{|c|c|} \hline 1 & 1 \\ \hline 2 & 3 \\ \hline \end{array}, \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 2 & 3 \\ \hline \end{array}, \begin{array}{|c|c|} \hline 1 & 1 \\ \hline 3 & 3 \\ \hline \end{array}, \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & 3 \\ \hline \end{array}, \begin{array}{|c|c|} \hline 1 & 1 \\ \hline 2 & 4 \\ \hline \end{array}, \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 2 & 4 \\ \hline \end{array}, \begin{array}{|c|c|} \hline 1 & 1 \\ \hline 3 & 4 \\ \hline \end{array}, \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & 4 \\ \hline \end{array}, \begin{array}{|c|c|} \hline 1 & 1 \\ \hline 4 & 4 \\ \hline \end{array}, \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 4 & 4 \\ \hline \end{array}.$$

Observe that the set \mathcal{T} of these $(2, 2)$ -tableaux is closed. By Proposition 3.2(ii), $X^{(4)} \cdot F(T)$ has eight summands, each obtained by changing an entry of 4 in the final six $(2, 2)$ -tableaux above to 3. In all but two cases, the new $(2, 2)$ -tableau obtained is semistandard, and so present in the closed set \mathcal{T} ; under our agreed identification, the summand is of the form $\dots \wedge F(s) \wedge \dots \wedge F(s) \wedge \dots$, and so vanishes. Let u and u' denote the final two $(2, 2)$ -tableaux shown above. The corresponding summands of $X^{(4)} \cdot F(T)$ are $F(U)$ and $F(U')$ where U and U' are the plethystic semistandard tableaux of shape $(2, 2)^{(1^{11})}$ with sets of entries

$$\mathcal{T} \setminus \{u\} \cup \left\{ \begin{array}{|c|c|} \hline 1 & 1 \\ \hline 4 & 3 \\ \hline \end{array} \right\} \quad \text{and} \quad \mathcal{T} \setminus \{u'\} \cup \left\{ \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 4 & 3 \\ \hline \end{array} \right\},$$

respectively. By the snake relation defined in (2.5) with $A = \{(2, 1)\}$ and $B = \{(1, 2), (2, 2)\}$ we have, working in $\nabla^{(2,2)}(E)$,

$$\begin{aligned} F \left(\begin{array}{|c|c|} \hline 1 & 1 \\ \hline 4 & 3 \\ \hline \end{array} \right) &= F \left(\begin{array}{|c|c|} \hline 1 & 1 \\ \hline 3 & 4 \\ \hline \end{array} \right) + F \left(\begin{array}{|c|c|} \hline 1 & 4 \\ \hline 1 & 3 \\ \hline \end{array} \right) = F \left(\begin{array}{|c|c|} \hline 1 & 1 \\ \hline 3 & 4 \\ \hline \end{array} \right) \\ F \left(\begin{array}{|c|c|} \hline 1 & 2 \\ \hline 4 & 3 \\ \hline \end{array} \right) &= F \left(\begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & 4 \\ \hline \end{array} \right) + F \left(\begin{array}{|c|c|} \hline 1 & 4 \\ \hline 2 & 3 \\ \hline \end{array} \right) = F \left(\begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & 4 \\ \hline \end{array} \right) - F \left(\begin{array}{|c|c|} \hline 1 & 3 \\ \hline 2 & 4 \\ \hline \end{array} \right). \end{aligned}$$

Therefore $F(U) = 0$ and $F(U') = -F(T')$ where T' is the plethystic semistandard tableau of shape $(2, 2)^{(1^{11})}$ whose entries are the same as T , except for the final entry u' , which is replaced with $\begin{array}{|c|c|} \hline 1 & 3 \\ \hline 2 & 4 \\ \hline \end{array}$; since this new $(2, 2)$ -tableau is greater in the total order than all the tableaux in \mathcal{T} , no reordering within the column is necessary in order to make T' semistandard. Therefore $X^{(4)} \cdot F(T) \neq 0$, and so $F(T)$ is *not* a highest-weight vector. As expected from the proof of Proposition 7.2, we have obtained a plethystic semistandard tableau T' of more dominant weight, namely, $(17, 11, 9, 7)$, by expressing $X^{(4)} \cdot F(T)$ in the canonical basis of $\nabla^{(1^{11})}(\nabla^{(2,2)}(E))$. We leave it to the reader to show that $F(T')$ is a highest-weight vector, and correspondingly, T' has maximal weight for its shape.

Our final example shows that the converse of Proposition 7.2 is false.

Example 7.5. Let T be the plethystic semistandard tableau of shape $(2, 1)^{(1^4)}$ with entries

$$\begin{array}{|c|c|} \hline 1 & 1 \\ \hline 2 & \\ \hline \end{array}, \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 2 & \\ \hline \end{array}, \begin{array}{|c|c|} \hline 1 & 3 \\ \hline 2 & \\ \hline \end{array}, \begin{array}{|c|c|} \hline 1 & 4 \\ \hline 2 & \\ \hline \end{array}$$

read from top to bottom. Then $F(T) \in \bigwedge^4(\nabla^{(2,1)}(E))$ is a highest-weight vector of weight $(5, 5, 1, 1)$ and so $\langle s_{(1^4)} \circ s_{(2,1)}, s_{(5,5,1,1)} \rangle \geq 1$. However T is not of maximal weight for its shape since $(6, 4, 2) \supseteq (5, 5, 1, 1)$ and the plethystic semistandard tableau U of shape $(2, 1)^{(1^4)}$ with entries

$$\begin{array}{|c|c|} \hline 1 & 1 \\ \hline 2 & \\ \hline \end{array}, \begin{array}{|c|c|} \hline 1 & 1 \\ \hline 3 & \\ \hline \end{array}, \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 2 & \\ \hline \end{array}, \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & \\ \hline \end{array}$$

read from top to bottom has weight $(6, 4, 2)$. It is easily seen that U has maximal weight in the dominance order, and so $F(U)$ is a highest-weight vector. In fact there are two plethystic semistandard tableau of shape $(2, 1)^{(1^4)}$ and weight $(6, 4, 2)$, the second is obtained from U by swapping the 2 and 3 in the final $(2, 1)$ -tableau entry above. Thus, by Theorem 1.5, $\langle s_{(1^4)} \circ s_{(2,1)}, s_{(6,4,2)} \rangle = 2$; this is one of the smallest examples where the multiplicity of a maximal constituent is more than 1.

REFERENCES

1. Silvana Abeasis, *The $GL(V)$ -invariant ideals in $S(S^2V)$* , Rend. Mat. (6) **13** (1980), no. 2, 235–262.
2. Edward A. Bender and Donald E. Knuth, *Enumeration of plane partitions*, J. Combinatorial Theory Ser. A **13** (1972), 40–54.
3. Giandomenico Boffi, *On some plethysms*, Adv. Math. **89** (1991), no. 2, 107–126.
4. Michel Brion, *Stable properties of plethysm: on two conjectures of Foulkes*, Manuscripta Math. **80** (1993), no. 4, 347–371.
5. Winfried Bruns, Aldo Conca, and Matteo Varbaro, *Relations between the minors of a generic matrix*, Adv. Math. **244** (2013), 171–206.
6. C. Carré and B. Leclerc, *Splitting the square of a schur function into its symmetric and antisymmetric parts*, J. Alg. Comb. **4** (1995), 201–231.
7. Eun J. Choi, Young H. Kim, Hyoung J. Ko, and Seoung J. Won, *GL_n -decomposition of the Schur complex $S_r(\bigwedge^2 \phi)$* , Bull. Korean Math. Soc. **40** (2003), no. 1, 29–51.
8. Melanie de Boeck, *A study of Foulkes modules using semistandard homomorphisms*, arXiv:1409.0734 (2014), 10 pages.
9. Suzie Dent, *Incidence structure of partitions*, Ph.D. thesis, UEA, 1997.
10. Vlastimil Dlab and Claus Michael Ringel, *The module theoretical approach to quasi-hereditary algebras*, Representations of algebras and related topics (Kyoto, 1990), London Math. Soc. Lecture Note Ser., vol. 168, Cambridge Univ. Press, Cambridge, 1992, pp. 200–224.
11. Karin Erdmann and Mark J. Wildon, *Introduction to Lie algebras*, Springer Undergraduate Mathematics Series, Springer-Verlag London, Ltd., London, 2006.
12. H. O. Foulkes, *Concomitants of the quintic and sextic up to degree four in the coefficients of the ground form*, J. London Math. Soc. **25** (1950), 205–209.
13. William Fulton and Joe Harris, *Representation theory, a first course*, Graduate Texts in Mathematics, vol. 129, Springer, 1991.
14. J. A. Green, *Polynomial representations of gl_n , with an appendix on Schensted correspondence and Littelmann paths by K. Erdmann and M. Schocker*, 2nd ed., Lecture Notes in Mathematics, vol. 830, Springer, 2007.
15. David J. Hemmer and Daniel K. Nakano, *Specht filtrations for Hecke algebras of type A*, J. London Math. Soc. (2) **69** (2004), no. 3, 623–638.
16. James E. Humphreys, *Introduction to Lie algebras and representation theory*, Graduate Texts in Mathematics, vol. 9, Springer-Verlag, New York-Berlin, 1978, Second printing, revised.

17. Christian Ikenmeyer, *Geometric complexity theory, tensor rank, and Littlewood–Richardson coefficients*, Ph.D. thesis, Universität Paderborn, 2012.
18. G. D. James and A. Kerber, *The representation theory of the symmetric group*, Encyclopedia of Mathematics and its Applications, vol. 16, Addison-Wesley Publishing Co., Reading, Mass., 1981.
19. G. D. James, *The representation theory of the symmetric groups*, Lecture Notes in Mathematics, vol. 682, Springer, Berlin, 1978.
20. Thomas Kahle and Mateusz Michałek, *Plethysm and lattice point counting*, Found. Comput. Math. **16** (2016), no. 5, 1241–1261.
21. Nicholas A. Loehr and Jeffrey B. Remmel, *A computational and combinatorial exposé of plethystic calculus*, J. Algebraic Combin. **33** (2011), no. 2, 163–198.
22. I. G. Macdonald, *Symmetric functions and Hall polynomials*, second ed., Oxford Mathematical Monographs, The Clarendon Press Oxford University Press, New York, 1995, With contributions by A. Zelevinsky, Oxford Science Publications.
23. M. J. Newell, *A theorem on the plethysms of S -functions*, Quart. J. Math. Oxford **2** (1951), 161–166.
24. Rowena Paget, *A family of modules with Specht and dual Specht filtrations*, J. Algebra **312** (2007), no. 2, 880–890.
25. Rowena Paget and Mark Wildon, *Minimal and maximal constituents of twisted Foulkes characters*, J. Lond. Math. Soc. **93** (2016), 301–318.
26. Rowena Paget and Mark Wildon, *Generalized Foulkes modules and maximal and minimal constituents of plethysms of Schur functions*, arXiv:1608.04018 (August 2016), 44.
27. Richard P. Stanley, *Enumerative combinatorics. Vol. 2*, Cambridge Studies in Advanced Mathematics, vol. 62, Cambridge University Press, Cambridge, 1999, With a foreword by Gian-Carlo Rota and appendix 1 by Sergey Fomin.
28. ———, *Positivity problems and conjectures in algebraic combinatorics*, Mathematics: frontiers and perspectives, Amer. Math. Soc., Providence, RI, 2000, pp. 295–319.
29. Mark Wildon, *Vertices of Specht modules and blocks of the symmetric group*, J. Algebra **323** (2010), no. 8, 2243–2256.

SCHOOL OF MATHEMATICS, STATISTICS AND ACTUARIAL SCIENCE, UNIVERSITY OF KENT, CANTERBURY, CT2 7FS, U.K.

E-mail address: `melaniedeboeck@hotmail.com`

SCHOOL OF MATHEMATICS, STATISTICS AND ACTUARIAL SCIENCE, UNIVERSITY OF KENT, CANTERBURY, CT2 7FS, U.K.

E-mail address: `r.e.paget@kent.ac.uk`

DEPARTMENT OF MATHEMATICS, ROYAL HOLLOWAY, UNIVERSITY OF LONDON, EGHAM, TW20 0EX, U.K.

E-mail address: `mark.wildon@rhul.ac.uk`