COHOMOLOGICAL FINITENESS CONDITIONS IN BREDON COHOMOLOGY

D. H. KOCHLOUKOVA, C. MARTÍNEZ-PÉREZ, AND B. E. A. NUCINKIS

ABSTRACT. We show that soluble groups G of type Bredon-FP_{∞} with respect to the family of all virtually cyclic subgroups of G are always virtually cyclic. In such a group centralizers of elements are of type FP_{∞}. We show that this implies the group is polycyclic. Another important ingredient of the proof is that a polycyclic-by-finite group with finitely many conjugacy classes of maximal virtually cyclic subgroups is virtually cyclic. Finally we discuss refinements of this result: we only impose the property Bredon-FP_n for some $n \leq 3$ and restrict to abelian-by-nilpotent, abelian-by-polycyclic or (nilpotent of class 2)-by-abelian groups.

1. INTRODUCTION

In this paper we study soluble groups G of type Bredon-FP_{∞} with respect to the family \mathcal{VC} of all virtually cyclic subgroups of G. Bredon cohomology with respect to a family \mathcal{X} of subgroups plays a role in studying classifying spaces for families similar to the role played by ordinary cohomology in studying Eilenberg-Mac Lane spaces. A G-CW complex X is said to be a model for a classifying space for a family \mathcal{X} of subgroups of G, denoted by $E_{\mathcal{X}}G$, if X^H is contractible if $H \in \mathcal{X}$ and empty otherwise. For the family \mathcal{F} of finite subgroups, the classifying space for proper actions, also denoted $\underline{E}G$, has received a lot of attention. The classifying space for the family \mathcal{VC} , denoted $\underline{E}G$, has only recently been looked at [8, 17]. In [8] the following was conjectured:

Conjecture 1.1. [8, Juan-Pineda and Leary] Let G be a group admitting a finite model for $\underline{E}G$. Then G is virtually cyclic.

In [17] the dimension of the classifying spaces $\underline{E}G$ for virtually polycyclic groups is given in terms of the Hirsch length and in [5] Bredon cohomological dimension for soluble groups is considered.

In this note we shall address the above conjecture purely algebraically by considering finiteness conditions in Bredon cohomology. This is motivated by the fact that a group G has a finite type model for a classifying space with isotropy in a family if and only if the group is of type Bredon-FP_{∞} and there is a model for a classifying space with finite 2-skeleton [16].

Finiteness conditions for the family \mathcal{F} of all finite subgroups in soluble groups are by now very well understood. Recently it was shown that a soluble group of type FP_{∞} is of type Bredon- FP_{∞} for the class \mathcal{F} [18] and admits a finite model for <u>E</u>G

Date: August 13, 2008.

 $^{2000\} Mathematics\ Subject\ Classification.\ 20J05.$

This work was supported by EPSRC grant EP/F045395/1 and LMS Scheme 4 grant 4708. The second named author was also supported by Gobierno de Aragon and MTM2007-68010-C03-01.

 $\mathbf{2}$

[9]. The proofs there utilised the following description given by Lück [15]: A group G is of type Bredon-FP_{∞} with respect to the family \mathcal{F} if and only if G has finitely many conjugacy classes of finite subgroups and each centraliser $C_G(K)$ of a finite subgroup K is of type FP_{∞}. For an algebraic proof of this fact see also [9].

The main result, Theorem 5.1 gives an answer to Conjecture 1.1 for soluble groups. As we will also show, this result can easily be extended to elementary amenable groups.

Theorem 5.1. Soluble groups G of type Bredon-FP_{∞} with respect to the class of virtually cyclic groups are virtually cyclic.

In the next section we show a partial analogue to Lück's result for the family of all virtually cyclic subgroups reducing the proof of our main theorem 5.1 to an analysis of centralizers of virtually cyclic subgroups and of conjugacy classes of maximal virtually cyclic subgroups.

Part of the proof of Theorem 5.1 reduces then to the study of groups where centralizers of cyclic subgroups are finitely generated. This is not a new problem and some partial answers can be found in the work of J. Lennox [13]. In [12] J. Lennox asked whether a soluble group G is polycyclic if all centralizers of its finitely generated subgroups are finitely generated. A positive answer to this question was given in [13] when G is nilpotent-by-polycyclic and this result was strengthened further by showing that a group G with an abelian normal subgroup A such that G/A is nilpotent and the centralizer (in G) of every element of A is finitely generated, is polycyclic. In the same paper it was also conjectured that the same result holds for abelian-by-polycyclic groups. In Theorem 3.7 we give an affirmative answer of this problem for abelian-by-polycyclic groups G of homological type FP₃ using the recent result that such a group G is nilpotent-by-abelian-by-finite [6]. It is still an open question whether a soluble group G is polycyclic if the centralizer of any element of G is finitely generated.

The final step of the proof of Theorem 5.1 involves conjugacy classes of maximal virtually cyclic subgroups. More precisely we show that if a polycyclic-by-finite group has finitely many conjugacy classes of maximal virtually cyclic subgroups then it is virtually cyclic. We show a more general result: Every nilpotent-by-abelian-by-finite group with the maximal condition on virtually cyclic subgroups and finitely many conjugacy classes of maximal virtually cyclic subgroups is virtually cyclic, see Theorem 4.6.

2. Bredon cohomology with respect to virtually cyclic subgroups

Let us begin this section with recalling a few prerequisites from Bredon cohomology. Here we replace the group G by the orbit category $\mathcal{O}_{\mathcal{X}}G$, where, as before, \mathcal{X} denotes a family of subgroups of G. The category $\mathcal{O}_{\mathcal{X}}G$ has as objects the transitive Gsets with stabilizers in \mathcal{X} . Morphisms in this category are G-maps. In this note we shall mainly be concerned with the families \mathcal{F} of finite subgroups and \mathcal{VC} of virtually cyclic subgroups and we write $\mathcal{O}_{\mathcal{F}}G$ and $\mathcal{O}_{\mathcal{VC}}G$ for the orbit categories. Modules over the orbit category, or $\mathcal{O}_{\mathcal{X}}G$ -modules, are contravariant functors from the orbit category to the category of abelian groups. Exactness is defined pointwise: A sequence $A \to B \to C$ of $\mathcal{O}_{\mathcal{X}}G$ -modules is exact at B if and only if $A(\Delta) \to$ $B(\Delta) \to C(\Delta)$ is exact at $B(\Delta)$ for every transitive G-set Δ . The category of $\mathcal{O}_{\mathcal{X}}G$ -modules has enough projectives, which are constructed as follows: For any G-sets Δ and Ω we denote by $[\Delta, \Omega]$ the set of G-maps from Δ to Ω . Let $\mathbb{Z}[\Delta, \Omega]$ be the free abelian group on $[\Delta, \Omega]$. Now fix Ω and let Δ range over the transitive G-sets with stabilizers in \mathcal{X} . This gives rise to an $\mathcal{O}_{\mathcal{X}}$ G-module $\mathbb{Z}[\ ,\Omega]$. Let G/K be a transitive G-set with $K \in \mathcal{X}$. Then $\mathbb{Z}[\ ,G/K]$ is a free module. Projective modules are defined analogously to the ordinary case. The trivial $\mathcal{O}_{\mathcal{X}}$ G-module, denoted $\mathbb{Z}(\)$, is the constant functor from $\mathcal{O}_{\mathcal{X}}G$ to the category of abelian groups. Bredon cohomology is now defined via projective resolutions of $\mathbb{Z}(\)$. The notions of type Bredon-FP, Bredon-FP_n $(n \geq 0$ an integer) and Bredon-FP_{∞} are defined in terms of projective resolutions of $\mathbb{Z}(\)$ over $\mathcal{O}_{\mathcal{X}}G$ analogously the classical notions of type FP, FP_n and FP_{∞}.

To stay in line with notation previously used, we say a module is of type \underline{FP}_{∞} if it is of type Bredon- FP_{∞} with respect to the family of all finite subgroups and is of type \underline{FP}_{∞} if it is of type Bredon- FP_{∞} with respect to the family of all virtually cyclic subgroups. We also denote by $\underline{\mathbb{Z}}$ and $\underline{\mathbb{Z}}$ the respective constant modules.

We shall now give a description of groups of type $\underline{\underline{FP}}_n$ and $\underline{\underline{FP}}_\infty$ in terms of cohomological finiteness conditions of centralisers of virtually cyclic subgroups and conjugacy in \mathcal{VC} . Let us consider the following two conditions on a group G:

- (1) G has the maximal condition on virtually cyclic subgroups (max-vc) and has finitely many conjugacy classes of maximal virtually cyclic subgroups.
- (2) There are finitely many virtually cyclic groups H_1, \ldots, H_n of G such that every virtually cyclic subgroup is subconjugated to one of the H_i s.

Obviously, (1) implies (2) but the converse is not true in general.

Example 2.1. Let G be the semidirect product of $A = \mathbb{Z}[\frac{1}{p}]$ by an infinite cyclic subgroup generated by t, which acts by multiplication by $\frac{1}{p}$. Then G satisfies (2) but not (1).

We have, however:

Lemma 2.2. If G has property (2) and max-vc then it has property (1).

Proof. We only have to prove that there are finitely many conjugacy classes of maximal virtually cyclic subgroups. But this is obvious as the subgroups H_1, \ldots, H_n of condition (2) can in this case be taken to be maximal virtually cyclic. And for any other maximal virtually cyclic T, we have $T^x \leq H_i$ for some $x \in T$, some i. Then T^x is also maximal virtually cyclic and therefore $T^x = H_i$.

The following Lemma is an analogue to [9, Lemma 3.1]

Lemma 2.3. Let G be a group. Then G is of type $\underline{\underline{FP}}_0$ if and only if G satisfies condition (2).

Proof. Let G be of type $\underline{\underline{FP}}_0$. Then the trivial module $\underline{\mathbb{Z}}$ has a finitely generated free Bredon module mapping onto it. Hence there is a G-finite G-set Ω with virtually cyclic stabilizers and an epimorphism

 $\mathbb{Z}[,\Omega] \twoheadrightarrow \underline{\mathbb{Z}}.$

Now let K be an arbitrary virtually cyclic subgroup of G. Evaluating the above epimorphism at G/K we obtain an epimorphism $\mathbb{Z}\Omega^K \twoheadrightarrow \mathbb{Z}$ and therefore Ω^K is non-empty. Hence K is subconjugated to a representative of the finitely many conjugacy classes of subgroups which have fixed points in Ω . Conversely, if G satisfies condition (2), then we can take $\Omega = \bigsqcup_{H_i} G/H_i$ where H_i runs through a set of conjugacy class representatives as above, and the obvious augmentation map $\mathbb{Z}[\ ,\Omega] \twoheadrightarrow \mathbb{Z}$ is an epimorphism. \Box

We say a Bredon module M is finitely generated if there is a finite $\mathcal{O}_{\mathcal{X}}G$ -set Σ in the sense of Lück [14, 9.16, 9.19] such that there is a free Bredon module F on Σ mapping onto M. In particular, a projective Bredon-module P is finitely generated if there is a G-finite G-set Ω with stabilisers in \mathcal{X} such that P is a direct summand of $\mathbb{Z}[-,\Omega]$.

Lemma 2.4. Let $P(\)$ be a finitely generated projective $\mathcal{O}_{\mathcal{VC}}G$ -module, and let K be a virtually cyclic subgroup. Then, after evaluation, P(G/K) is a module of type FP_{∞} over the Weyl-group $WK = N_G(K)/K$.

Proof. By the remarks above, it suffices to show that for each virtually cyclic group H, the module $\mathbb{Z}(G/H)^K \cong \mathbb{Z}[G/K, G/H]$ is of type $\operatorname{FP}_{\infty}$ as a WK-module. Furthermore, $(G/H)^K$ is a WK-set in which each xH is stabilized by $(N_G(K) \cap H^{x^{-1}})K/K = WK_x$. Now

$$\mathbb{Z}(G/H)^K = \oplus \mathbb{Z}(WK/WK_x)$$

is a finitely generated WK-module. An argument analogous to [19, Proposition 6.3] reduces to showing that each $\mathbb{Z}(WK/WK_x)$ is of type FP_{∞} . But this follows from the fact that WK_x is virtually cyclic so it is a group of type FP_{∞} .

Corollary 2.5. Let M() be a Bredon-module of type $\underline{\text{FP}}_n$. Then, for each virtually cyclic subgroup K, the WK-module M(G/K) is of type FP_n .

Proof. This is a straight forward dimension shift applying [3, Proposition 1.4]. \Box

Proposition 2.6. Let G be a group of type $\underline{\underline{FP}}_n$. Then the following two equivalent statements hold:

- (i) G satisfies condition (2) and for each virtually cyclic subgroup K, the Weyl-group WK is of type FP_n.
- (ii) G satisfies condition (2) and for each virtually cyclic subgroup K, the centralizer $C_G(K)$ is of type FP_n .

Proof. Statement (i) follows directly from Lemmas 2.3 and 2.4. Since K is virtually cyclic and hence of type FP_{∞} , it follows that $N_G(K)$ is of type FP_n if and only $WK = N_G(K)/K$ is [3, Proposition 2.7]. Aut(K) is a finite group and hence the index $|N_G(K) : C_G(K)|$ is finite and therefore $N_G(K)$ is of type FP_n if and only $C_G(K)$ is of type FP_n .

Let us also remark on an obvious link with \underline{FP}_{∞} .

Corollary 2.7. Let G be a group of type $\underline{\text{FP}}_{\infty}$. Then G is of type $\underline{\text{FP}}_{\infty}$.

Proposition 2.8. [17, Corollary 5.4] Let G be a group admitting a finite (finite type) model for $\underline{E}G$. Then G admits a finite (finite type) model for $\underline{E}G$.

3. Soluble groups

Lemma 3.1. Let Q be a finitely generated abelian group and V be a finitely generated $\mathbb{Z}Q$ -module. Then V is finitely generated as an additive group if and only if $\mathbb{Z}Q/P_i$ is finitely generated as an additive group for every minimal associated prime P_i of V.

Proof. Suppose that $\mathbb{Z}Q/P_i$ is finitely generated for $1 \leq i \leq s$, where P_1, \ldots, P_s are the minimal associated primes of V. Let I be the annihilator $ann_{\mathbb{Z}Q}(V)$. Then $\sqrt{I} = P_1 \cap \ldots \cap P_s$ and $\mathbb{Z}Q/P_1 \cap \ldots \cap P_s$ embeds in $\mathbb{Z}Q/P_1 \oplus \ldots \oplus \mathbb{Z}Q/P_s$ via the diagonal map, so $\mathbb{Z}Q/\sqrt{I}$ is finitely generated as an additive group. Let s be a natural number such that $\sqrt{I}^s \subseteq I$. Note that every quotient $\sqrt{I}^j/\sqrt{I}^{j+1}$ is a finitely generated as additive groups. Since V is finitely generated as a $\mathbb{Z}Q/I$ -module, V is finitely generated as an additive group.

Conversely suppose that V is finitely generated as an additive group. Since $\mathbb{Z}Q/P_1 \oplus \ldots \oplus \mathbb{Z}Q/P_s$ embeds in V it follows that every $\mathbb{Z}Q/P_i$ is finitely generated.

Lemma 3.2. Let $N_2 \rightarrow G_2 \rightarrow Q_2$ be a short exact sequence of groups with N_2 nilpotent, Q_2 abelian and G_2 of type FP₂. Suppose that G_2 and N_1 are subgroups of a group G, N_1 is nilpotent, $G_2 \subseteq N_G(N_1)$ and N_2 is a normal subgroup in N_1 such that N_1/N_2 is finitely generated and abelian. Then $G_1 = N_1G_2$ is of type FP₂.

Proof. The lemma is obvious if G_2 is normal in G_1 but we cannot assume this. Since N_1/N_2 is a finitely generated abelian group, it is a direct sum of cyclic groups. Let T be a finite subset of N_1 such that the images of the elements of T in N_1/N_2 are the generators of the direct cyclic summands. We split T as a disjoint union $T = T_0 \cup T_1$, where the images in N_1/N_2 of the elements of T_0 have finite order and the images in N_1/N_2 of the elements of T_1 have infinite order. Write $T_0 = \{t_1, \ldots, t_m\}$, a_i is the order of the image of t_i in N_1/N_2 , $T_1 = \{t_{m+1}, \ldots, t_s\}$. Fix a generating set $Y = \{g_1, \ldots, g_s\}$ of G_2 . Note that every element g of G_1 can be written in a unique way as

(1)
$$g = t_1^{z_1} \dots t_s^{z_s} h$$
 where $h \in G_2, z_i \in \mathbb{Z}$ and $0 \le z_i < a_i$ for $i \le m$.

Then for some reduced words $\alpha_{i,j}, \beta_i$ on $Y \cup Y^{-1}$ we have the following equations in G_1

$$t_i t_j = t_j t_i \alpha_{i,j}$$
 for $i > j, t_i^{a_i} = \beta_i$ for $1 \le i \le m$

and for $y \in Y \cup Y^{-1}$ there are reduced words $\gamma_{i,y}$ on $Y \cup Y^{-1}$ such that

$$yt_iy^{-1} = t_1^{z_{1,i,y}} \dots t_s^{z_{s,i,y}} \gamma_{i,y}$$
 where $z_{j,i,y} \in \mathbb{Z}$ and $0 \le z_{j,i,y} < a_j$ for $j \le m$

Let $\langle Y | R \rangle$ be a presentation of G_2 that shows it is of type FP₂ i.e. Y is as above, R might be infinite but the corresponding relation module is finitely generated as a $\mathbb{Z}G_2$ -module. Then G_1 has a presentation

$$\langle Y \cup T \mid R \cup S \rangle$$

where S is

 $\{t_j^{-1}t_i^{-1}t_jt_i\alpha_{i,j}\}_{1\leq j< i\leq s} \cup \{t_i^{a_i}\beta_i^{-1}\}_{m+1\leq i\leq s} \cup \{yt_i^{-1}y^{-1}t_1^{z_{1,i,y}}\dots t_s^{z_{s,i,y}}\gamma_{i,y}\}_{y\in Y\cup Y^{-1}, 1\leq i\leq s}.$ Indeed the group H defined by the presentation $\langle Y\cup T \mid R\cup S\rangle$ maps surjectively to G_2 and the elements of H satisfy the normal form condition (1). Finally the presentation $\langle Y \cup T \mid R \cup S \rangle$ is obtained from the presentation $\langle Y \mid R \rangle$ of G_2 by adding finitely many relations and generators i.e. G_1 is of type FP₂.

Theorem 3.3. Let $N \rightarrow G \rightarrow Q$ be a short exact sequence of groups with Nnilpotent and Q abelian, G finitely generated such that every quotient of G/Z(N) is of type FP₂ and for every $g \in N$ the centraliser $C_G(g)$ is finitely generated and the image of the centralizer $C_G(g)$ in any quotient of G/Z(N) is of type FP₂. Then Gis polycyclic.

Proof. We show first that Z(N) is finitely generated as a group. By assumption G/Z(N) is of type FP₂, thus the centre Z(N) of N is finitely generated as a $\mathbb{Z}Q$ -module, where Q acts via conjugation. Let P_1, \ldots, P_s be the minimal associated primes for the $\mathbb{Z}Q$ -module Z(N), thus $M = \mathbb{Z}Q/P_1 \oplus \ldots \oplus \mathbb{Z}Q/P_s$ embeds in Z(N) as a $\mathbb{Z}Q$ -submodule (i.e. if v_i is an element of Z(N) with $Ann_{\mathbb{Z}Q}(v_i) = P_i$ then M is isomorphic to $\sum_{1 \le i \le s} \mathbb{Z}Qv_i$) and we think of M as a submodule of Z(N). Let a be the element of M whose projection to $\mathbb{Z}Q/P_i$ is the unity element for every $1 \le i \le s$.

By assumption the centralizer $C_G(a)$ is finitely generated. Note that $N \subseteq C_G(a)$ and define $Q_0 = C_G(a)/N \subseteq Q$. Since $C_G(a)/Z(N)$ is of type FP₂, Z(N) is finitely generated as a $\mathbb{Z}Q_0$ -module. Since $\mathbb{Z}Q_0$ is Noetherian, every $\mathbb{Z}Q_0$ -submodule of Z(N) is finitely generated. In particular M is finitely generated as a $\mathbb{Z}Q_0$ -module. Since Q_0 acts trivially on a we deduce that Q_0 acts trivially on M. Thus M is finitely generated as an additive abelian group and by Lemma 3.1 Z(N) is finitely generated.

We prove the theorem by induction on the nilpotency class of N. We aim to prove that the assumptions of the theorem hold for the quotient group $\overline{G} = G/Z(N)$ and then the proof can be completed by induction. Write \overline{g} for the image of $g \in G$ in \overline{G} and define $\overline{N} = N/Z(N)$. For some fixed $g \in N$ define the homomorphism

$$\varphi: C_{\overline{N}}(\overline{g}) \to Z(N)$$

that sends \overline{n} to [n, g]. Note that $Ker(\varphi) = C_N(g)/Z(N)$ and that $C_{\overline{N}}(\overline{g})/Ker(\varphi)$ is a subgroup of Z(N), hence is a finitely generated abelian group. Then $G_0 = C_{\overline{G}}(\overline{g})$ has a filtration of subgroups

$$G_2 = C_G(g)/Z(N) \subseteq G_1 = G_2 C_{\overline{N}}(\overline{g}) \subseteq G_0.$$

Let $N_2 = C_N(g)/Z(N)$ and $N_1 = C_{\overline{N}}(\overline{g})$, so N_1/N_2 is finitely generated and abelian. Then by Lemma 3.2 G_1 is of type FP₂. Since G_1 is normal in G_0 and G_0/G_1 is finitely generated abelian, G_0 is of type FP₂.

By assumption the image of $C_G(g)$ in any quotient of G/Z(N) is of type FP₂. Then the above argument applied for the images of the groups G_2, G_1, G_0, N_1, N_2 in a quotient of G/Z(N) shows that the image of G_0 in any quotient of G/Z(N) is of type FP₂. Thus we can apply the inductive argument for G/Z(N) and deduce that G/Z(N) is polycyclic.

Corollary 3.4. Let $N \rightarrow G \rightarrow Q$ be a short exact sequence of groups such that N is nilpotent of class 2, Q is abelian, G is of type FP_2 and for every $g \in N$ the centralizer $C_G(g)$ is of type FP_2 . Then G is polycyclic.

Proof. Every metabelian quotient of a group of type FP_2 not containing non-cyclic free subgroups is of type FP_2 [4]. Then the previous theorem applies.

There are known examples of groups G that are split extensions of a nilpotent of class 3 group N by a finite rank free abelian group Q such that G is finitely presented, Z(N) = Z(G) is infinitely generated as a group and thus G/Z(N) is not of type FP₂ [1]. Though these examples show that the first condition of Theorem 3.3 is quite restrictive, at the same time these examples do not satisfy the second type conditions about the centralizer, i.e. there is an element $g \in N$ such that $C_G(g)$ is not even finitely generated.

Lemma 3.5. Let G be a soluble group of type FP_{∞} . Then every quotient of G is of type FP_{∞} .

Proof. This follows directly by [2, Thm. 4] and the classification of soluble FP_{∞} groups as constructible [10], [11].

Corollary 3.6. Let G be a soluble group of type FP_{∞} such that the centraliser $C_G(g)$ is of type FP_{∞} for every element $g \in G$. Then G is polycyclic.

Proof. It follows directly from Theorem 3.3 and Lemma 3.5.

7

In the following theorem a stronger condition on G is imposed at the expense of relaxing the conditions on the centralizers. Note that we have not included that every quotient of G is of type FP₃ but this holds by one of the main results of [6].

Theorem 3.7. Let G be an abelian-by-polycyclic group of type FP_3 such that the centraliser $C_G(g)$ is finitely generated for every element $g \in A$, where A is a normal abelian subgroup of G with G/A polycyclic. Then G is polycyclic.

Proof. We aim to prove that A is finitely generated as a group. We view A as a $\mathbb{Z}H$ -module with H-action induced by conjugation, where H = G/A. By [6] G is nilpotent-by-abelian-by-finite. By going down to a subgroup of finite index we can assume that G is nilpotent-by-abelian, so the commutator H' acts nilpotently on A i.e. there is some natural number s such that $A\Omega^s = 0$ where Ω is the augmentation ideal of $\mathbb{Z}H'$. Then A has a filtration of $\mathbb{Z}H$ -modules

$$0 = A_s = A\Omega^s \subseteq \ldots \subseteq A_{i+1} = A\Omega^{i+1} \subseteq A_i = A\Omega^i \subseteq \ldots \subseteq A_0 = A$$

where H' acts trivially on A_i/A_{i+1} via conjugation i.e. A_i/A_{i+1} is a $\mathbb{Z}Q$ -module, where Q = H/H'. Since A is a finitely generated module over a Noetherian ring $\mathbb{Z}H$, we have that every A_i/A_{i+1} is a finitely generated as a $\mathbb{Z}Q$ -module.

Suppose that A_{i+1} is finitely generated as an additive group. We will prove that A_i is finitely generated as an additive group. We follow the method used in the proof of Theorem 3.4. Let P_1, \ldots, P_s be the minimal associated primes for the $\mathbb{Z}Q$ -module A_i/A_{i+1} , thus $M = \mathbb{Z}Q/P_1 \oplus \ldots \oplus \mathbb{Z}Q/P_s$ is a submodule of A_i/A_{i+1} .

Let \tilde{a} be the element of M whose projection to $\mathbb{Z}Q/P_j$ is the unity element for every $1 \leq j \leq s$ and a be a preimage of \tilde{a} in A. Note that $A \subseteq C_G(a)$ and define $H_0 = C_G(a)/A \subseteq H$. Since $C_G(a)$ is finitely generated A is finitely generated as a $\mathbb{Z}H_0$ -module, so A_i/A_{i+1} is finitely generated as a $\mathbb{Z}Q_0$ -module and so M is finitely generated as a $\mathbb{Z}Q_0$ -module, where $Q_0 = H_0/(H_0 \cap H')$. By the choice of a we get that Q_0 acts trivially on M, thus M is finitely generated as an additive group and by Lemma 3.1 A_i/A_{i+1} is finitely generated as an additive group.

4. Conjugacy classes

Let T and G be groups with T acting on G. We consider the following condition:

(1T) G has the maximal condition for virtually cyclic subgroups (max-vc) and only finitely many T-orbits of maximal virtually cyclic subgroups.

If G = T acts on itself by conjugation, then (1T) is equivalent to condition (1) of Section 2. Note also that max-vc is subgroup closed and that for any $x \in T$ and any $H \leq G$ maximal virtually cyclic, the group H^x is also maximal virtually cyclic.

Lemma 4.1. Let $h(x) \in \mathbb{Z}[x]$ be an integer polynomial. There are infinitely many primes p such that there exists an integer n_p with

$$p|h(n_p).$$

<u>Proof:</u> Consider prime factors of h(n) when $n \in \mathbb{Z}$. If we had only a finite set $\{p_1, \ldots, p_r\}$, then for each *i* let n_i such that

$$p_i^{n_i-1}|h(0), p_i^{n_i} \not| h(0).$$

Note that for any m with $\prod_{i=1}^{r} p_i^{n_i} | m$,

 $p_i^{n_i} \not| h(m).$

So we may choose m big enough so that h(m) has some other prima factor q not in the finite set above.

Lemma 4.2. Let A be an abelian group and T be an infinite cyclic group acting on A. Suppose A has property (1T). Then A is finite.

Proof. Note first that the condition (1T) implies that the order of the torsion elements of A is bounded. Therefore there are only finitely many primes p for which the p-primary component of A is non-trivial, so by splitting A in its torsion free part and its p-primary components we may assume A is either torsion free or that all its elements have p-power order for a fixed prime p.

If we denote a_1, \ldots, a_s for representatives of the *T*-orbits of maximal cyclic subgroups in *A*, then there is an epimorphism

$$\mathbb{Z}T \oplus . :. \oplus \mathbb{Z}T \twoheadrightarrow A$$

and each $\mathbb{Z}Ta_i \leq A$ has also (1T), so we may assume

$$A = \mathbb{Z}Ta \cong \mathbb{Z}T/I$$

for certain ideal $I \triangleleft \mathbb{Z}T$. Let $m\mathbb{Z} = I \cap \mathbb{Z}$ (the assumption above implies that m is either 0 or a power of p). As $A = \mathbb{Z}T/I$ has only finitely many T-orbits of maximal virtually cyclic subgroups, $I \neq m\mathbb{Z}T$. Therefore we may choose some polynomial

$$f(t) \in I \cap \mathbb{Z}[t] \setminus m\mathbb{Z}[t]$$

of minimal degree k where t is a fixed generator of T. Note that k > 0. Then the family $\{1, t, t^2, \ldots, t^{k-1}\}$ yields a maximal independent subset of the quotient $A = \mathbb{Z}T/I$. Therefore A has finite Prüfer rank so max-vc implies that it is a finitely generated abelian group. In the torsion case we get the result so from now on we assume A is torsion free.

There is a principal ideal $I_1 \triangleleft \mathbb{Z}T$ such that $I \leq I_1$ and I_1/I is torsion. Indeed $I \otimes_{\mathbb{Z}} \mathbb{Q}$ is an ideal of the principal ideal domain $\mathbb{Q}T$, hence is generated by some element h(t). In fact h(t) can be chosen to be a primitive polynomial in the variable

9

t. A localisation version of Gauss' lemma implies $\mathbb{Z}T \cap (\mathbb{Q}Th(t)) = \mathbb{Z}Th(t)$, and we can define $I_1 = \mathbb{Z}Th(t)$. So our assumption that A is torsion free implies that $I = \mathbb{Z}Th(t)$.

Now, by Lemma 4.1 there is an infinite set of primes Ω such that for each $p \in \Omega$ there exists an integer n_p with

$$t - n_p | h(t) \mod p.$$

Any of the elements $t - n_p + I$ generates a maximal cyclic subgroup of A. So our assumption on A implies that there are only finitely many T-orbits between them. And therefore there are only finitely many distinct ideals in the set

$$\{I_p = \mathbb{Z}T(t - n_p, h(t)) : p \in \Omega\}.$$

Moreover,

$$\mathbb{Z}T/I_p \cong \mathbb{Z}/h(n_p)\mathbb{Z}$$

is finite so there are only finitely many maximal ideals containing some ideal in the set above.

But for each prime $p \in \Omega$,

$$\mathbb{Z}T(t - n_p, h(t)) \le M_p = \mathbb{Z}T(t - n_p, h(t), p)$$

which is a maximal ideal and obviously $M_p \neq M_q$ for $p \neq q$. This is a contradiction.

From now on let G = T act on itself by conjugation and we shall consider the conditions (1) and (2) of Section 2.

Lemma 4.3. Let $H \triangleleft G$ be a finite index normal subgroup. If G has property (1), then so does H.

Proof. Clearly, H has max-vc. Moreover for any $C \leq H$ maximal virtually cyclic there is some $D \leq G$ also maximal virtually cyclic with $C \leq D$ and from this follows that $D \cap H = C$. As the index $|G : N_G(D)H|$ is finite, the G-conjugacy class of D splits after intersecting with H as finitely many H-conjugacy classes of maximal virtually cyclic subgroups of H.

Lemma 4.4. Assume that G has (1) and there is some $N \triangleleft G$ with G/N finitely generated abelian. Then G/N is virtually cyclic.

Proof. By Lemma 4.3 we may assume that G/N is torsion free. Note that for any $\overline{C} = \langle \overline{x} \rangle \leq G/N$ maximal cyclic we may choose an $x \in G$ with $xN = \overline{x}$ and a maximal virtually cyclic C_x of G with $x \in C_x$. And $C_x^g = C_y$ implies $x^g N = yN$. Therefore there are finitely many conjugacy classes of maximal cyclic subgroups of G/N. As this is torsion free abelian we deduce that G/N is cyclic.

Lemma 4.5. Assume that G has property (1) and there is $N \triangleleft G$ finite. Then G/N has property (1).

Proof. For any $N \leq C \leq G$ such that C/N is virtually cyclic, C itself is virtually cyclic. This yields the result.

Theorem 4.6. Let G be finitely generated nilpotent-by-abelian-by-finite with property (1). Then G is virtually cyclic.

10

Proof. By Lemma 4.3 we may assume G is nilpotent-by-abelian. Let $N \triangleleft G$ be nilpotent with G/N abelian. As T = G/N is finitely generated by Lemma 4.4 it must be cyclic. Let A = Z(N). Then G/A acts on A and N/A is in the kernel of the action. Moreover the condition (1) implies that A has property (1T) with respect to the group T = G/N. So by Lemma 4.2 A must be finite. And Lemma 4.5 implies that G/A also has (1) so an induction on the nilpotency length of N yields the result.

Corollary 4.7. Let G be an abelian-by-polycyclic-by-finite group of type FP_3 and with property (1). Then G is virtually cyclic.

Proof. By [6] abelian-by-polycyclic groups of type FP_3 are nilpotent-by-abelian-by-finite and we can apply the previous Theorem.

Corollary 4.8. Let G be polycyclic-by-finite group with property (2). Then G is virtually cyclic.

Proof. As G is polycyclic-by-finite, it has max and hence max-vc. Therefore by Lemma 2.2 G has (1) so it suffices to apply the previous Theorem. \Box

5. The main results

Theorem 5.1. Soluble groups of type \underline{FP}_{∞} are virtually cyclic.

Proof. Proposition 2.6 and Corollary 3.6 imply that the group is polycyclic and satisfies property (2). Now apply Corollary 4.8. \Box

Theorem 5.2. Let G be an abelian-by-nilpotent group of type $\underline{\underline{FP}}_1$. Then G is virtually cyclic.

Proof. By Proposition 2.6 centralizers of elements are finitely generated. [13, Thm. C] implies G is polycyclic. Now apply Corollary 4.8. \Box

Theorem 5.3. Let G be a (nilpotent of class 2)-by-abelian group of type $\underline{\underline{FP}}_2$. Then G is virtually cyclic.

Proof. This follows from Proposition 2.6, Corollary 3.4 and Theorem 4.6. \Box

Theorem 5.4. Let G be an abelian-by-polycyclic group of types FP_3 and $\underline{\text{FP}}_1$. Then G is virtually cyclic.

Proof. This follows from Proposition 2.6, Theorem 3.7 and Corollary 4.7. \Box

Corollary 5.5. Let G be an abelian-by-polycyclic group of type $\underline{\underline{FP}}_{3}$. Then G is virtually cyclic.

Proof. By Proposition 2.6 every group of type $\underline{\text{FP}}_3$ is of type FP_3 .

We shall conclude with a remark on virtual notions. Let C be a class of groups. We say a group is virtually C if G has a finite index subgroup belonging to C.

Lemma 5.6. Let G_1 be a group of type $\underline{\underline{FP}}_n$, $n \ge 0$. Then any subgroup G of finite index in G_1 is of type $\underline{\underline{FP}}_n$.

Proof. Any $\mathcal{O}_{\mathcal{VC}}G_1$ -projective resolution of $\underline{\mathbb{Z}}$ can be viewed as an $\mathcal{O}_{\mathcal{VC}}G$ -projective resolution. We therefore need to show that every finitely generated $\mathcal{O}_{\mathcal{VC}}G_1$ -projective is finitely generated as $\mathcal{O}_{\mathcal{VC}}G$ -module. This reduces to showing that $\mathbb{Z}[\ ,G_1/K]$ is a finitely generated $\mathcal{O}_{\mathcal{VC}}G$ -projective for all virtually cyclic subgroups K of G_1 . G_1/K viewed as a G-set is the G-finite set $\Omega = \sqcup G/G \cap K^{x^{-1}}$, where x runs through a set of representatives of the finite number of cosets of G in G_1 . \Box

Remark 5.7. In the statements of Theorem 5.1 through to Corollary 5.5 above we can let G be virtually-C, where C stands for soluble, abelian-by-nilpotent, (nilpotent of class 2)-by-abelian or abelian-by-polycyclic respectively.

As mentioned in the introduction, we can also strengthen Theorem 5.1 to apply to the class of elementary amenable groups:

Corollary 5.8. Elementary amenable groups of type \underline{FP}_{∞} are virtually cyclic.

Proof. These groups are of type FP_{∞} by 2.6 and hence have finite Hirsch length and a bound on the orders of their finite subgroups [11]. Combining this with [7], we might assume that the group is virtually soluble. Now apply Lemma 5.6.

References

- H. Abels. An example of a finitely presented solvable group. In Homological group theory (Proc. Sympos., Durham, 1977), volume 36 of London Math. Soc. Lecture Note Ser., pages 205–211. Cambridge Univ. Press, Cambridge, 1979.
- [2] G. Baumslag and R. Bieri. Constructable solvable groups. Math. Z., 151(3):249–257, 1976.
- [3] R. Bieri. Homological dimension of discrete groups. Queen Mary College Mathematical Notes. Queen Mary College Department of Pure Mathematics, London, second edition, 1981.
- [4] R. Bieri and R. Strebel. Valuations and finitely presented metabelian groups. Proc. London Math. Soc. (3), 41(3):439-464, 1980.
- [5] M. Fluch. Dimensions in bredon cohomology. Ph.D Thesis, University of Southampton, in preparation.
- [6] J. Groves, D. Kochloukova, and C. Rodrigues. Abelian-by-polycyclic groups of homological type FP₃. To appear in J. Group Theory
- [7] J. A. Hillman and P. A. Linnell. Elementary amenable groups of finite Hirsch length are locally-finite by virtually-solvable. J. Austral. Math. Soc. Ser. A, 52(2):237–241, 1992.
- [8] D. Juan-Pineda and I. J. Leary. On classifying spaces for the family of virtually cyclic subgroups. Ádem, Alejandro (ed.) et al., Recent developments in algebraic topology. A conference to celebrate Sam Gitler's 70th birthday, San Miguel de Allende, México, December 3–6, 2003. Providence, RI: American Mathematical Society (AMS). Contemporary Mathematics 407, 135-145 (2006).
- [9] P. Kropholler, C. Martínez-Pérez, and B. E. A. Nucinkis. Cohomological finiteness conditions for elementary amenable groups. preprint, 14 pp., 2008.
- [10] P. H. Kropholler. Cohomological dimension of soluble groups. J. Pure Appl. Algebra, 43(3):281–287, 1986.
- [11] P. H. Kropholler. On groups of type $(FP)_{\infty}$. J. Pure Appl. Algebra, 90(1):55–67, 1993.
- [12] J. C. Lennox. Centrality in soluble groups. In *Infinite groups 1994 (Ravello)*, pages 153–162. de Gruyter, Berlin, 1996.
- [13] J. C. Lennox. On soluble groups in which centralizers are finitely generated. Rend. Sem. Mat. Univ. Padova, 96:131–135, 1996.
- [14] W. Lück. Transformation groups and algebraic K-theory, volume 1408 of Lecture Notes in Mathematics. Springer-Verlag, Berlin, 1989., Mathematica Gottingensis.
- [15] W. Lück. The type of the classifying space for a family of subgroups. J. Pure Appl. Algebra, 149(2):177–203, 2000.
- [16] W. Lück and D. Meintrup. On the universal space for group actions with compact isotropy. Grove, Karsten (ed.) et al., Geometry and topology, Aarhus. Proceedings of the conference

on geometry and topology, Aarhus, Denmark, August 10-16, 1998. Providence, RI: American Mathematical Society (AMS). Contemp. Math. 258, 293-305 (2000).

- [17] W. Lück and M. Weiermann. On the classifying space of the family of virtually cyclic subgroups. arXiv:math.AT/0702646v1, 2007.
- [18] C. Martínez-Pérez and B. E. A. Nucinkis. Virtually soluble groups of type FP_{∞} . to appear Comment. Math. Helv.
- [19] B. E. A. Nucinkis. Cohomology relative to a G-set and finiteness conditions. Topology Appl., 92(2):153–171, 1999.

Dessislava H. Kochloukova, Department of Mathematics, State University of Campinas, Cx. P. 6065, 13083-970 Campinas, SP, Brazil

 $E\text{-}mail\ address: \texttt{desiQunicamp.br}$

Conchita Martínez-Pérez, Departamento de Matemáticas, Universidad de Zaragoza, 50009 Zaragoza, Spain

 $E\text{-}mail\ address: \texttt{conmarQunizar.es}$

BRITA E. A. NUCINKIS, SCHOOL OF MATHEMATICS, UNIVERSITY OF SOUTHAMPTON, SOUTHAMPTON, SO17 1BJ, UNITED KINGDOM

 $E\text{-}mail\ address: \verb"bean@soton.ac.uk"$